

Tameness and Homogeneity

Dedicated to Professor C.M. Ringel on the occasion of
his 60'th birthday

Zhang Yingbo

Xu Yunge

Contents

1	Introduction	1
1.1	AR-sequences and AR-quivers	1
1.2	Tameness and wildness	2
1.3	The main theorem: tameness and homogeneity	3
1.4	The outline of the proof	3
1.5	Notations.	5
2	Bimodule problems	6
2.1	The category $P_1(\Lambda)$	6
2.2	The bimodule problem $(\mathcal{K}, \mathcal{M}, H)$	9
2.3	The triangular basis of $(\mathcal{K}, \mathcal{M})$	13
2.4	An order on triangular basis	14
2.5	Weyr matrices	16
2.6	Reductions	18
3	Induced bimodule problems	22
3.1	Induced bimodule problems	22
3.2	Deletions and size changes	25
3.3	Reduction functors and reduction sequences	26
3.4	Canonical forms	28
3.5	Krull-Schmidt property	30
3.6	Corresponding bocses	36
4	Freely parameterized bimodule problems	41
4.1	Parameters	41
4.2	Freely parameterized bimodule Problems	41
4.3	The differential of the first arrow	44
4.4	Valuation matrices	48
4.5	The reduction sequence towards a valuation matrix	49
4.6	Valuation representation categories	53
5	Minimally wild bimodule problems	56
5.1	The wild theorem	56
5.2	The tame theorem	60
5.3	Local wild bimodule problems	62
5.4	Triangular formulae	64
5.5	The minimal size assumption	66
5.6	The classification of minimally wild bimodule problems	68
6	Exact categories and almost split conflations	70
6.1	Some basic concepts	70
6.2	Exact categories $P(\Lambda)$ and $P_1(\Lambda)$	71
6.3	Morphisms in $R(\mathfrak{A})$	73

6.4	An exact structure on $R(\mathfrak{A})$	76
6.5	Homogeneous conflations	78
6.6	Minimal bocses	79
7	Almost split conflations in reductions	81
7.1	Projective and injective objects	81
7.2	Homogeneous property in reductions	82
7.3	Critically non-homogeneous bocses $\mathfrak{B}_1$ and $\mathfrak{B}_2$	85
7.4	A critically non-homogeneous local boc $\mathfrak{B}_3$	88
7.5	A critically non-homogeneous local boc $\mathfrak{B}_4$	89
7.6	An example in case of MW5	91
8	One-sided differentials	96
8.1	Matrix problems	96
8.2	One-sided differentials I	101
8.3	One-sided differentials II	103
8.4	The structure of loop $\bar{b}$	106
8.5	Some lemmas in algebra	111
8.6	Some lemmas in category theory	113
9	Added columns	117
9.1	The structure of added columns	117
9.2	The reduction sequence $(\tilde{\Delta})$	119
9.3	The differentials in partial boc $\tilde{\mathfrak{B}}^s$	123
9.4	The calculation of $\delta(c^0)$	126
9.5	The induction on (α^η, l^η)	127
9.6	Non-homogeneous partial boc ${}^p\mathfrak{B}^s$	131
10	Bipartite bimodule Problems	135
10.1	Bipartite property	135
10.2	Locations	136
10.3	Links	137
10.4	The non-homogeneous property in case of MW5	138
10.5	The main theorem	141
	References	143

Abstract

Let Λ be a finite-dimensional algebra over an algebraically closed field, then Λ is either tame or wild. Is there any homological description in terms of AR-translations on tameness? Or equivalently, is there any combinatorial description in terms of AR-quivers? The answer is yes. In the present paper we prove the following main theorem: “ Λ is tame if and only if almost all modules are isomorphic to their Auslander-Reiten translations, if and only if they lie in homogeneous tubes”.

The method used in the paper is bimodule problem, which has been studied by several authors. We treat bimodule problems and their reductions in terms of matrices and generalized Jordan forms respectively. In particular, we introduce a concept of freely parameterized bimodule problems corresponding to layered bocses in order to define the reductions of bimodule problems. Moreover, we list all the possibilities of the differential of the first arrow of a layered boc such that the induced boc is still layered and preserves all the free parameters, especially in wild case.

Keywords bimodule problem, layered boc, tameness, wildness, almost split conflation, homogeneity,

2000 Mathematics Subject Classification: 15A21, 16G20, 16G60, 16G70

1 Introduction

1.1 AR-sequences and AR-quivers

Let Λ be a finite-dimensional algebra (associative, with 1) over an algebraically closed field k , and $\Lambda\text{-mod}$ be the category of finite-dimensional left Λ -modules. Auslander and Reiten have defined a concept of almost split sequence (i.e. AR-sequence) in a module category [AR]. If we take any non-projective indecomposable Λ -module N (resp. non-injective module M), there exists an almost split sequence ending at N (resp. starting at M):

$$(e) \quad 0 \longrightarrow M \xrightarrow{\iota} E \xrightarrow{\pi} N \longrightarrow 0$$

where $M \cong DTr(N)$, the Auslander-Reiten translation of N , or AR-translation of N for short. $DTr : \Lambda\text{-mod} \rightarrow \Lambda\text{-mod}$ is a functor, which maps indecomposable non-projective Λ -modules to non-injective Λ -modules [AR]. AR-sequences yield a combinatorial description of $\Lambda\text{-mod}$. Namely, we draw a vertex $[M]$ corresponding to an iso-class of indecomposable module M , and an arrow $[L] \xrightarrow{[\zeta]} [M]$ representing that there is an irreducible map $\zeta : L \rightarrow M$, then the obtained quiver is called an Auslander-Reiten quiver of Λ , or AR-quiver of Λ for short. If we have

$$E = \bigoplus_{i=1}^l E_i, \quad \iota = (\iota_1, \iota_2, \dots, \iota_l) \text{ and } \pi = (\pi_1, \pi_2, \dots, \pi_l)^T$$

in the sequence (e), where E_i are indecomposable, T stands for the transpose of a matrix, then $\iota_i : M \rightarrow E_i$ and $\pi_i : E_i \rightarrow N$ are irreducible maps between indecomposables. Conversely, if $M \xrightarrow{\iota'} E'$ (resp. $E' \xrightarrow{\pi'} N$) with E' being indecomposable is an irreducible map, there exists some module E'' and morphism $M \xrightarrow{\iota''} E''$ (resp. $E'' \xrightarrow{\pi''} N$) such that the sequence

$$(e') \quad 0 \longrightarrow M \xrightarrow{(\iota', \iota'')} E' \oplus E'' \xrightarrow{(\pi', \pi'')} N \longrightarrow 0$$

is equivalent to the almost split sequence (e).

Furthermore, the shape of stable components of AR-quivers has been completely described in [Rie], [HPR] and [Z]. The simplest stable components which mostly deserve to pay attention to are so-called homogeneous tubes consisting of vertices $[M_i]$ and arrows $[M_i] \xrightleftharpoons[\pi_i]{\iota_i} [M_{i+1}]$, where i is any positive integer. Thus almost split sequences lying in a homogeneous tube have the shape of

$$0 \longrightarrow M_1 \xrightarrow{\iota_1} M_2 \xrightarrow{\pi_1} M_1 \longrightarrow 0$$

$$0 \longrightarrow M_i \xrightarrow{(\pi_{i-1}, \iota_i)} M_{i-1} \oplus M_{i+1} \xrightarrow{(\iota_{i-1}, \pi_i)} M_i \longrightarrow 0$$

for $i \geq 2$, and $DTr(M_i) \cong M_i$ for $i = 1, 2, \dots$.

1.2 Tameness and wildness

The well known Drozd's theorem tells us that a finite-dimensional algebra Λ over an algebraically closed field k is either of tame representation type or of wild representation type [D].

Definition 1.2.1 [D, CB1, DS] A finite-dimensional k -algebra Λ is of tame representation type, if for any positive integer d , there are a finite number of localizations $R_i = k[x, f_i(x)^{-1}]$ of $k[x]$ and Λ - R_i -bimodules T_i which are free as right R_i -modules, such that almost all (except finitely many) indecomposable Λ -modules of dimension at most d are isomorphic to

$$T_i \otimes_{R_i} R_i / (x - \lambda)^m,$$

for some $\lambda \in k$, $f_i(\lambda) \neq 0$, and some positive integer m .

Definition 1.2.2 [D, CB1] A finite-dimensional k -algebra Λ is of wild representation type if there is a finitely generated Λ - $k\langle x, y \rangle$ -bimodule T , which is free as a right $k\langle x, y \rangle$ -module, such that the functor

$$T \otimes_{k\langle x, y \rangle} - : k\langle x, y \rangle\text{-mod} \rightarrow \Lambda\text{-mod}$$

preserves indecomposability and isomorphism classes.

Crawley-Boevey once proposed to consider generic modules instead of dealing with families of modules [CB4]. A Λ -module G is called generic, if G is of infinite length over Λ , but of finite endlength (the length over its own endomorphism ring) and G is indecomposable. Then Λ is tame, if and only if Λ is generically tame. Generically tame type means for any positive integer d , there are only finitely many generic modules of endlength at most d . Based on Crawley-Boevey's work on tame algebras, Krause presents a new description of tameness in terms of Ziegler spectrum, functor category and model theory. More precisely, he proves that a finite-dimensional algebra over an algebraically closed field is of tame representation type, if and only if every generic Λ -module appears as the only generic module on the Ziegler closure of a homogeneous tube [Kr].

In summary, the definition of tameness given by Drozd, and the description in terms of generic modules by Crawley-Boevey, as well as the description in terms of Ziegler spectrum by Krause, all involve infinite-dimensional modules outside $\Lambda\text{-mod}$. A natural question arises: is there any internal description of tameness, which only involves finite-dimensional modules? Moreover, is there any combinatorial description of tameness in terms of AR-quivers?

1.3 The main theorem: tameness and homogeneity

A result due to Crawley-Boevey may lead to an internal description of tameness.

Theorem 1.3.1 [CB1]: Let Λ be a finite-dimensional algebra over an algebraically closed field. If Λ is of tame representation type, then almost all Λ -modules are isomorphic to their Auslander-Reiten translations. And then lie in homogeneous tubes.

Here “almost all modules” means “for any fixed positive integer d , all but a finite number of isomorphism classes of indecomposable Λ -modules of dimension at most d .”

The proof was based on the method of bocses. In fact, a minimal bocs possesses such a property. It is also conjectured in the same paper that the property may possibly describe tameness. Since then, several experts have tried to prove the converse of the theorem, expecting that the same method of bocses would work, that is, none of the minimally wild bocses would have such a property. Unfortunately, this was proved to be wrong. A counterexample was constructed in [ZLB], and later a great number of wild categories having such a property were given in [BCLZ] and [V].

Another approach to testify the converse of Crawley-Boevey’s theorem is to prove Ringel’s conjecture. A wild algebra Λ is called τ -wild if there exist infinitely many non-isomorphic τ -variant (i.e. non-homogeneous) indecomposable modules of dimension d , for a certain positive integer d , where $\tau = DTr$. Han Yang introduces the concept of controlled wild algebras (see [H] and [R3]). And Ringel conjectures that all wild algebras in the meaning of Drozd are controlled wild with finite controlled index. Nagase shows that all controlled wild algebras with finite controlling index are τ -wild [N2]. Therefore, if Ringel’s conjecture holds, so does the converse of Crawley-Boevey’s theorem. The covering criterion for an algebra to be controlled wild, given by Han Yang in [H] is very effective. It seems to be impossible to find a concrete wild algebra which is not controlled wild. However, Ringel’s conjecture is still open up to now.

In this paper we will prove the converse of Crawley-Boevey’s theorem: if almost all Λ -modules are isomorphic to their AR -translations, then Λ is tame.

Main theorem 1.3.2 *Let Λ be a finite-dimensional algebra over an algebraically closed field. Then Λ is tame, if and only if almost all Λ -modules are isomorphic to their AR -translations, and then if and only if they lie in homogeneous tubes.*

1.4 The outline of the proof

Our argument relies on the methods of bimodule problems [S] and bocses [Ro, D, CB1]. The notion of bocses was introduced by Rojter in [Ro] in order to apply formulation of

the matrix problems to representation theory. There are several formulations of matrix problems [ZZ] such as bocses, differential graded categories [Ro], differential biquivers, relatively projective categories [BK], bimodule problems [CB2], lift categories [CB3], etc. The following picture is suggested by Crawley-Boevey.

$$\begin{aligned}
& \{\text{matrix problems given by “layered” or “almost free” bocses [D, CB1]}\} \\
& \supset \{\text{matrix problems of the form } (\mathcal{K}, \mathcal{M}, d), \text{ where } \mathcal{K} \text{ is an algebra, } \mathcal{M} \text{ an} \\
& \quad \mathcal{K}\text{-}\mathcal{K}\text{-bimodule, } d \text{ a derivation, which is called a bimodule problem [CB3]}\} \\
& = \{\text{matrix problems of the form } (\mathcal{K}, \mathcal{M}, H), \text{ where } \mathcal{K} \text{ is an upper triangular} \\
& \quad \text{matrix algebra, } \mathcal{M} \text{ a matrix } \mathcal{K}\text{-}\mathcal{K}\text{-bimodule, } H \text{ is a fixed matrix, which} \\
& \quad \text{determines a derivation (see 2.2)}\} \\
& \supset \{\text{matrix problem given by bipartite bimodule problems (see 10.1)}\} \\
& \supset \{\text{matrix problems of the form } P_1(\Lambda) \text{ (see 2.1)}\}.
\end{aligned}$$

Throughout the paper, we concentrate on the third set of matrix problems, which are convenient for the calculations in the proof of the main theorem. On the other hand, such matrix problems are helpful to understand the notion of bocses, since their structure is very concrete.

The reductions of matrix problems given by layered bocses are well-established by the same authors, which seem to be elegant for theoretical purposes. Among all the reductions, Belitskii’s reduction algorithm is very effective for reducing an individual matrix to a canonical form, which may be considered as a generalized Jordan form, under some admissible transformations. In this paper, we will describe the reductions in terms of generalized Jordan forms for a bimodule problem given in the third set, which is based only on linear algebra.

The idea of our proof of the main theorem is as follows. Suppose an algebra Λ is of wild representation type and has homogeneous property, which means that the bimodule problem $(\mathcal{K}, \mathcal{M})$, and equivalently the Drozd boc $\mathfrak{A}$, corresponding to Λ is also wild and homogeneous. The homogeneous property is inherited to any induced structure given by reductions. Then we are restricted to dealing with one of those configurations listed in the wild theorem 5.1 obtained at some stage of reductions. For some easy cases, using the method proposed by Bautista in [B], we can directly construct infinitely many iso-classes of non-homogeneous modules of a fixed dimension. For the most difficult case, we will construct a non-homogeneous parameterized matrix based on the bipartite property of $(\mathcal{K}, \mathcal{M})$. And thereby we show that $(\mathcal{K}, \mathcal{M})$, as well as Λ , has no homogeneous property, which is a contradiction to the assumption.

1.5 Notations.

In this paper k represents an algebraically closed field. By an algebra we mean an associative finite-dimensional k -algebra with an identity, which we may assume (without loss of generality) to be basic and connected. By a module over an algebra Λ we mean a finite-dimensional left Λ -module. And by $\Lambda\text{-mod}$ we denote the category of finite-dimensional left Λ -modules.

Let $Q = (Q_0, Q_1)$ be a quiver, where Q_0 is the set of vertices and Q_1 is the set of arrows. For any arrow $\alpha : i \rightarrow j$ in Q_1 , $s(\alpha)$ stands for the starting vertex of α and $e(\alpha)$ for the ending one. Let p be any path in Q , $s(p)$ and $e(p)$ stand also for the starting and ending vertices of p respectively. We write the composition of paths from left to right, i.e.

$$p \cdot q = \begin{cases} pq & \text{if } e(p) = s(q), \\ 0 & \text{if } e(p) \neq s(q). \end{cases}$$

According to a theorem due to Gabriel, for any basic finite-dimensional algebra Λ there exists some quiver Q and some admissible ideal I of kQ , such that $\Lambda \cong kQ/I$.

Let us denote the set of natural numbers by $\mathbb{N}$, the set of $m \times n$ matrices over k by $M_{m \times n}(k)$, or $M_n(k)$ whenever $m = n$. We denote by E_{ij} the matrix with the (i, j) -entry 1 and others zero, and by I_n the $n \times n$ identity matrix.

We assume that all the categories and functors under the consideration are k -linear.

2 Bimodule problems

This section is devoted to describing a special class of bimodule problems of the form $(\mathcal{K}, \mathcal{M}, H)$ listed in the third set of subsection 1.4. We will give the definition of $(\mathcal{K}, \mathcal{M}, H)$, calculate the triangular bases of $\mathcal{M}$ and $\text{rad}\mathcal{K}$, define Weyr matrices, and give the reductions for a triple of $(\mathcal{K}, \mathcal{M}, H)$.

2.1 The category $P_1(\Lambda)$

Let Λ be a finite-dimensional k -algebra, which is basic, and $\Lambda\text{-proj}$ be the full subcategory of $\Lambda\text{-mod}$ consisting of projective Λ -modules. Let us recall the category

$$P(\Lambda) = \{P_1 \xrightarrow{\alpha} P_0 \mid P_1, P_0 \in \Lambda\text{-proj}, \alpha \in \text{Hom}_\Lambda(P_1, P_0)\},$$

whose objects are morphisms between Λ -projectives. If $(P_1 \xrightarrow{\alpha} P_0), (Q_1 \xrightarrow{\beta} Q_0) \in P(\Lambda)$, the morphisms from $(P_1 \xrightarrow{\alpha} P_0)$ to $(Q_1 \xrightarrow{\beta} Q_0)$ are pairs (f_1, f_0) with $f_i \in \text{Hom}_\Lambda(P_i, Q_i)$, $i = 1, 0$, such that the following diagram commutes:

$$\begin{array}{ccc} P_1 & \xrightarrow{\alpha} & P_0 \\ f_1 \downarrow & & \downarrow f_0 \\ Q_1 & \xrightarrow{\beta} & Q_0 \end{array}$$

The compositions and additions of the morphisms are given componentwise.

There are a full subcategory

$$P_1(\Lambda) = \{(P_1 \xrightarrow{\alpha} P_0) \mid \text{Im}(\alpha) \subseteq \text{rad}(P_0)\} \subset P(\Lambda),$$

and a full subcategory

$$P_2(\Lambda) = \{(P_1 \xrightarrow{\alpha} P_0) \in P_1(\Lambda) \mid \text{Ker}(\alpha) \subseteq \text{rad}(P_1)\} \subset P_1(\Lambda).$$

It is well known that the functor $\text{Cok} : P_2(\Lambda) \rightarrow \Lambda\text{-mod}$ is a representation equivalence, i.e. it is dense, full and reflects isomorphisms.

The category $P_1(\Lambda)$ can be described explicitly in terms of matrices. Let J be the Jacobson radical of Λ with $J^m \neq 0$, $J^{m+1} = 0$, and $\Lambda/J \cong ke_1 \times \cdots \times ke_s$ with $\{e_1, e_2, \dots, e_s\}$ being a complete set of orthogonal primitive idempotents of Λ . Let us first take a k -basis of $e_i J^m e_j$, $1 \leq i, j \leq s$. Suppose we have already had a basis of J^l , and then extend it to a k -basis of J^{l-1} , such that the images of the added new elements form a k -basis of $e_i(J^{l-1}/J^l)e_j$, for $1 \leq i, j \leq s$. The obtained k -basis of J is denoted by

$$B = \{\zeta_1, \zeta_2, \dots, \zeta_{t-s}\}$$

with a fixed order given above, where $t = \dim \Lambda$. Moreover, by adding the primitive idempotents $e_1, e_2, \dots, e_s$ after ζ_{t-s} , we obtain an ordered k -basis of Λ .

Given any $a \in \Lambda$, we define a map $\bar{a} : \Lambda \rightarrow \Lambda$ given by $\bar{a}(b) = ab$, which is a k -linear transformation of the k -vector space Λ . Then ζ_i and e_j will correspond to some $t \times t$ matrices $\bar{\zeta}_i$ and $\bar{e}_j$ under the ordered basis respectively, which are upper triangular because of the order of the basis. Thus a matrix algebra $\tilde{\Lambda}$ follows, which is upper triangular and isomorphic to Λ , with a k -basis $\{\bar{\zeta}_1, \dots, \bar{\zeta}_{t-s}, \bar{e}_1, \dots, \bar{e}_s\}$. And $\tilde{\Lambda}$ is usually called a *left regular representation* of Λ .

Example 1. Let $Q = a \begin{array}{c} \circ \\ \bullet \\ \circ \end{array} c$ be a quiver, kQ be the path algebra of Q ,

$$I = \langle a^2, b^2, c^2, ab, ba, bc, cb, ca, ac \rangle$$

be an ideal of kQ generated by the elements in the bracket, and $\Lambda = kQ/I$. We denote the residue classes of $\{e, a, b, c\}$ in Λ still by $\{e, a, b, c\}$ respectively. Then an ordered k -basis $\{c, b, a, e\}$ of Λ yields

$$\begin{aligned} \tilde{\Lambda} &= \left\{ \begin{pmatrix} s_1 & 0 & 0 & s_4 \\ & s_1 & 0 & s_3 \\ & & s_1 & s_2 \\ & & & s_1 \end{pmatrix} \mid \forall s_i \in k \right\}, \\ \text{rad} \tilde{\Lambda} &= \left\{ \begin{pmatrix} 0 & 0 & 0 & m_c \\ & 0 & 0 & m_b \\ & & 0 & m_a \\ & & & 0 \end{pmatrix} \mid \forall m_a, m_b, m_c \in k \right\}. \end{aligned}$$


Example 2. Let $Q = \begin{array}{ccccc} & a & & c & \\ & \bullet & \xrightarrow{\quad} & \bullet & \\ 1 & \xleftarrow{b} & 2 & & 3 \end{array}$ be a quiver and $\Lambda = kQ$ be the path algebra of Q . If we choose an ordered k -basis $\{c, b, a, e_1, e_2, e_3\}$ of Λ , then


$$\begin{aligned} \tilde{\Lambda} &= \left\{ \begin{pmatrix} s_2 & 0 & 0 & 0 & 0 & s_6 \\ & s_2 & 0 & s_5 & 0 & 0 \\ & & s_2 & s_4 & 0 & 0 \\ & & & s_1 & 0 & 0 \\ & & & & s_2 & 0 \\ & & & & & s_3 \end{pmatrix} \mid \forall s_i \in k \right\}; \\ \text{rad} \tilde{\Lambda} &= \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & m_c \\ & 0 & 0 & m_b & 0 & 0 \\ & & 0 & m_a & 0 & 0 \\ & & & 0 & 0 & 0 \\ & & & & 0 & 0 \\ & & & & & 0 \end{pmatrix} \mid \forall m_a, m_b, m_c \in k \right\}. \end{aligned}$$

Example 3. [R1, D] Let $Q = a \circ \bullet \circ b$ be a quiver, $I = \langle a^2, ba - \alpha ab, ab^2, b^3 \rangle$ be an ideal of kQ with $\alpha \in k$ being a fixed non-zero constant, and let algebra $\Lambda = kQ/I$. Denote the residue classes of e, a, b in Λ still by e, a, b respectively. Moreover let us set $c = b^2, d = ab$. Then an ordered k -basis $\{d, c, b, a, e\}$ of Λ yields

$$\tilde{\Lambda} = \left\{ \begin{pmatrix} s_1 & 0 & s_2 & \alpha s_3 & s_5 \\ & s_1 & s_3 & 0 & s_4 \\ & & s_1 & 0 & s_3 \\ & & & s_1 & s_2 \\ & & & & s_1 \end{pmatrix} \mid \forall s_i \in k \right\},$$

$$\text{rad} \tilde{\Lambda} = \left\{ \begin{pmatrix} 0 & 0 & m_a & \alpha m_b & m_d \\ & 0 & m_b & 0 & m_c \\ & & 0 & 0 & m_b \\ & & & 0 & m_a \\ & & & & 0 \end{pmatrix} \mid \forall m_a, m_b, m_c, m_d \in k \right\}.$$



Example 4. Let $Q =$  be a quiver and $\Lambda = kQ$ be the path algebra of Q . Then an k -basis $\{f, d, c, b, a, e_1, e_2, e_3, e_4, e_5, e_6\}$ of Λ yields

$$\tilde{\Lambda} = \left\{ \begin{pmatrix} s_5 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & s_{11} \\ & s_4 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & s_{10} \\ & & s_3 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & s_9 \\ & & & s_2 & 0 & 0 & 0 & 0 & 0 & 0 & s_8 \\ & & & & s_1 & 0 & 0 & 0 & 0 & 0 & s_7 \\ & & & & & s_1 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & s_2 & 0 & 0 & 0 & 0 \\ & & & & & & & s_3 & 0 & 0 & 0 \\ & & & & & & & & s_4 & 0 & 0 \\ & & & & & & & & & s_5 & 0 \\ & & & & & & & & & & s_6 \end{pmatrix} \mid \forall s_i \in k \right\};$$

$$\text{rad}\tilde{\Lambda} = \left\{ \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_f \\ & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_d \\ & & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_c \\ & & & 0 & 0 & 0 & 0 & 0 & 0 & 0 & m_b \\ & & & & 0 & 0 & 0 & 0 & 0 & 0 & m_a \\ & & & & & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & 0 & 0 & 0 & 0 & 0 \\ & & & & & & & 0 & 0 & 0 & 0 \\ & & & & & & & & 0 & 0 & 0 \\ & & & & & & & & & 0 & 0 \\ & & & & & & & & & & 0 \end{pmatrix} \mid \forall m_a, m_b, m_c, m_d, m_f \in k \right\}.$$

2.2 The bimodule problem $(\mathcal{K}, \mathcal{M}, H)$

Definition 2.2.0 [CB2] By a bimodule problem we mean a triple (K, M, d) , where K is a k -algebra, M is a K - K -bimodule, and $d : K \rightarrow M$ is a derivation. $\square$

Proposition 2.2.0 (given by Crawley-Boevey) Let (K, M, d) be a bimodule problem, then there exist

- (1) a positive integer t ;
- (2) an upper triangular matrix algebra $\mathcal{K} \subset M_t(k)$ and an algebra isomorphism $\varphi : K \rightarrow \mathcal{K}$;
- (3) a $\mathcal{K}$ - $\mathcal{K}$ -bimodule $\mathcal{M} \subset M_t(k)$, and an isomorphism $\psi : M \rightarrow \mathcal{M}$, such that $\psi(xm) = \varphi(x)\psi(m)$, $\psi(mx) = \psi(m)\varphi(x)$ for any $x \in K, m \in M$;
- (4) a fixed matrix $H \in M_t(k)$, such that $\psi(d(x)) = \varphi(x)H - H\varphi(x)$ for any $x \in M$.

Proof. Let $N = M \oplus K$ as vector spaces. We define a K - K -bimodule structure on N given by $y(m, x) = (ym, yx)$, and $(m, x)y = (my - xd(y), xy)$ for any $m \in M, x, y \in K$. If we choose $h = (0, 1) \in N$, then

$$xh - hx = x(0, 1) - (0, 1)x = (0, x) - (-d(x), x) = (d(x), 0) \in M.$$

Let $A = K \oplus N$ be an algebra with multiplication $(x, n)(x', n') = (xx', xn' + nx')$. It is clear that $M \subset N \subset A$ and $K \subset A$. Since $N^2 = 0$, the idea $N \subset \text{rad}(A)$. Thus any complete set of orthogonal primitive idempotents of K is also that of A . Suppose that $\dim(A) = t$, choose a suitable k -basis of $\text{rad}(A) = \text{rand}(K) \oplus N$ according to the method given in 2.1, we can embed A as a upper triangular subalgebra $\tilde{A} \subset M_t(k)$, i.e there is an algebra isomorphism $\theta : A \rightarrow \tilde{A}$. Denote by φ the restriction of θ on K , by ψ that on M . And let $\mathcal{K} = \varphi(K)$, $\mathcal{M} = \psi(M)$, $H = \theta(h)$, the proof is completed. $\square$

Proposition 2.2.0 suggests the following definition of bimodule problems in terms of matrices in order to do some calculations.

Definition 2.2.1 A bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ given by matrices consists of the following datum:

- I. A linearly ordered set of integers $T = \{1, 2, \dots, t\}$ and an equivalent relation $\sim$ on T , where

$$T/\sim = \{\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_s\}.$$

- II. An upper triangular matrix algebra $\mathcal{K} = \{(s_{ij})_{t \times t} \in M_t(k)\}$, where $s_{ii} = s_{jj}$ when $i \sim j$; s_{ij} , when $i < j$, satisfy the following system of homogeneous linear equations in indeterminates x_{ij} :

$$\sum_{\mathcal{I} \ni i < j \in \mathcal{J}} c_{ij}^l x_{ij} = 0,$$

where the coefficients $c_{ij}^l \in k$, the equations are indexed by $1 \leq l \leq q_{\mathcal{I}\mathcal{J}}$ for some $q_{\mathcal{I}\mathcal{J}} \in \mathbb{N}$, and for each pair

$$(\mathcal{I}, \mathcal{J}) \in (T/\sim) \times (T/\sim).$$

- III. A $\mathcal{K}$ - $\mathcal{K}$ -bimodule $\mathcal{M} = \{(m_{ij})_{t \times t} \in M_t(k)\}$, where m_{ij} satisfy the following system of homogeneous linear equations in indeterminates z_{ij} :

$$\sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} d_{ij}^l z_{ij} = 0,$$

where the coefficients $d_{ij}^l \in k$, the equations are indexed by $1 \leq l \leq q'_{\mathcal{I}\mathcal{J}}$ for some $q'_{\mathcal{I}\mathcal{J}} \in \mathbb{N}$, and for each pair

$$(\mathcal{I}, \mathcal{J}) \in (T/\sim) \times (T/\sim).$$

- IV. A fixed matrix $H = (h_{ij})_{t \times t} \in M_t(k)$, where $h_{ij} = 0$ when $i \not\sim j$. And a derivation $d : \mathcal{K} \rightarrow \mathcal{M}$ is given by

$$d(S) = SH - HS$$

for any $S \in \mathcal{K}$. □

It is obvious that $(\mathcal{K}, \mathcal{M}, H)$ is a bimodule problem in the sense of Definition 2.2.0. We stress that definition 2.2.1 is a modification of definition 1.1 of [S]. Sometimes we write $(\mathcal{K}, \mathcal{M})$ instead of $(\mathcal{K}, \mathcal{M}, H)$ for short.

Corollary 2.2.1 $\{E_{\mathcal{I}} = \sum_{i \in \mathcal{I}} E_{ii} \mid \forall \mathcal{I} \in T/\sim\}$ is a complete set of primitive idempotents of $\mathcal{K}$. And $d(E_{\mathcal{I}}) = 0, \forall \mathcal{I} \in T/\sim$. □

Sometimes we write $s_{\mathcal{I}} = s_{ii}$ for any $i \in \mathcal{I}$.

Proposition 2.2.1 Let Λ be a k -algebra of dimension t having s pairwise orthogonal primitive idempotents, let $\tilde{\Lambda}$ be the left regular representation of Λ defined in 2.1. Set

$$\mathcal{K} = \begin{pmatrix} \tilde{\Lambda} & 0 \\ 0 & \tilde{\Lambda} \end{pmatrix}, \quad \mathcal{M} = \begin{pmatrix} 0 & \text{rad}\tilde{\Lambda} \\ 0 & 0 \end{pmatrix}, \quad H = (0).$$

and

$$T = \{1, 2, \dots, t; t+1, t+2, \dots, 2t\},$$

is the set of row (or column) indices of $\mathcal{K}$,

$$T/\sim = \{\mathcal{I}_1, \dots, \mathcal{I}_s; \mathcal{I}_{s+1}, \dots, \mathcal{I}_{2s}\},$$

where $\mathcal{I}_l = \{i | s_{ii} = s_{i_l i_l} \text{ for a fixed } i_l \in T\}$. Then $(\mathcal{K}, \mathcal{M}, H)$ is a bimodule problem. $\square$

The rest of this subsection is devoted to illustrating the representation category $\text{Mat}(\mathcal{K}, \mathcal{M})$ of a bimodule problem $(\mathcal{K}, \mathcal{M}, H)$, whose objects and morphisms are both given by matrices. The category is coincide with that given in [CB2].

A vector $\underline{n} = (n_1, n_2, \dots, n_t) \in \mathbb{N}^t$ is called a *size vector* of $(T, \sim)$ provided that $n_i = n_j$ when $i \sim j$ in T . And $n = \sum_{i=1}^t n_i$ is called the *size of* $\underline{n}$. On the other hand, if we denote n_i by $n_{\mathcal{I}} \forall i \in \mathcal{I}$, then $\underline{d} = (n_{\mathcal{I}_1}, n_{\mathcal{I}_2}, \dots, n_{\mathcal{I}_s})$ is called a *dimension vector* of $(T, \sim)$, and $d = \sum_{\mathcal{I} \in T/\sim} n_{\mathcal{I}}$ is called the *dimension of* $\underline{d}$.

Let $\underline{n}$ be any size vector,

$$\mathcal{M}_{\underline{n}} = \{(N_{ij})_{t \times t} \mid N_{ij} \text{ are } n_i \times n_j \text{ blocks satisfying the equation system III}\}.$$

Then any partitioned matrix $N \in \mathcal{M}_{\underline{n}}$ is called a *representation*, or a *matrix* over $(\mathcal{K}, \mathcal{M})$ of size vector $\underline{n}$.

If $\underline{m}$ is also a size vector of $(T, \sim)$, $M \in \mathcal{M}_{\underline{m}}$. Let

$$\mathcal{K}_{\underline{m} \times \underline{n}} = \{(S_{ij})_{t \times t} \mid S_{ij} \text{ are } m_i \times n_j \text{ blocks}\},$$

where $S_{ij} = 0$ if $i > j$; $S_{ii} = S_{jj}$ if $i \sim j$; S_{ij} satisfy the equation system II if $i < j$. Then a partitioned matrix $S \in \mathcal{K}_{\underline{m} \times \underline{n}}$ is called a *morphism* from M to N , provided that

$$MS - SN = d(S),$$

or equivalently

$$(M + H_{\underline{m}})S = S(N + H_{\underline{n}}),$$

where if $H = (h_{ij})_{t \times t}$, then $H_{\underline{n}} = (H_{ij})_{\underline{n} \times \underline{n}}$ is a partitioned matrix with $H_{ij} = 0$ if $i \not\sim j$, and $H_{ij} = h_{ij} I_{n_i}$ if $i \sim j$. $H_{\underline{m}}$ is defined similarly for the size vector $\underline{m}$. We denote the partitioned diagonal part of S by S_0 .

A morphism $S : M \rightarrow N$ is called an *isomorphism* provided that S is invertible. In this case, M and N are said to be *isomorphic*, and denoted by $M \simeq N$.

Corollary 2.2.2 (1) If $S \in \mathcal{K}_{\underline{m} \times \underline{n}}$, and $S' \in \mathcal{K}_{\underline{n} \times \underline{m}}$ such that $SS' = I$, $S'S = I$, then $S_0 S'_0 = I$, $S'_0 S_0 = I$.

(2) If $M \simeq N$, then $\underline{m} = \underline{n}$.

Proof. (1) The reason is that S and S' are both partitioned upper triangular.

(2) If $S : M \rightarrow N$ and $S' : N \rightarrow M$ are morphisms, such that $SS' = id_M$, and $S'S = id_N$, then $S_{\mathcal{I}} S'_{\mathcal{I}} = I_{m_{\mathcal{I}}}$ and $S'_{\mathcal{I}} S_{\mathcal{I}} = I_{n_{\mathcal{I}}}$ for any $\mathcal{I} \in T/\sim$. Thus $m_{\mathcal{I}} = n_{\mathcal{I}}$, consequently $\underline{m} = \underline{n}$ as desired. $\square$

Finally we fix some notations of various indices used in this paper by an example. The equivalent classes in $T/\sim$ are denoted by italic English letters $\mathcal{I}, \mathcal{J}, \mathcal{P}, \mathcal{Q}, \dots$; the positive integers in T by lower-case English $i, j, p, q, \dots$; the usual row and column indices of a matrix by lower-case Greek $\alpha, \beta, \gamma, \dots$. And we write $\alpha \in i, \beta \in j$ whenever the (α, β) -entry is sitting at the (i, j) -block.

Example. See Example 2 of 2.1 and Proposition 2.2.1. In the corresponding bimodule problem of Λ , $T = \{1, 2, 3, \dots, 12\}$,

$$\mathcal{I}_1 = \{1, 2, 3, 5\}, \mathcal{I}_2 = \{4\}, \mathcal{I}_3 = \{6\}, \mathcal{I}_4 = \{7, 8, 9, 11\}, \mathcal{I}_5 = \{10\}, \mathcal{I}_6 = \{12\}.$$

Let

$$\underline{m} = (2, 2, 2, 0, 2, 0; 0, 0, 0, 2, 0, 1)$$

be a size vector of $(T, \sim)$. Let us take

$$i_l = l, \quad l = 1, 2, \dots, 12; \quad \text{and} \quad \alpha_j = j, \quad j = 1, \dots, 11.$$

Then $\alpha_1, \alpha_2 \in i_1$; $\alpha_3, \alpha_4 \in i_2$; $\alpha_5, \alpha_6 \in i_3$; $\alpha_7, \alpha_8 \in i_5$; $\alpha_9, \alpha_{10} \in i_{10}$; $\alpha_{11} \in i_{12}$. The following is an 11×11 matrix $M \in \text{Mat}(\mathcal{K}, \mathcal{M})$ of size vector $\underline{m}$, and a morphism $S \in \text{End}_{(\mathcal{K}, \mathcal{M})}(M)$.

$$M = \begin{array}{c|cccc|c} \hline & & & & & 1 \\ & & & & & 0 \\ \hline & & & & 0 & \lambda \\ & & & & 1 & 0 \\ \hline & & & & 0 & 1 \\ & & & & 0 & 0 \\ \hline & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ \hline \end{array}, \quad S = \begin{array}{c|cccc|c} \hline s_1 & 0 & & & & i_1 \\ 0 & s_2 & & & & \\ \hline & s_1 & 0 & & & i_2 \\ & 0 & s_2 & & & \\ \hline & & s_1 & 0 & & i_3 \\ & & 0 & s_2 & & \\ \hline & & & s_1 & 0 & i_5 \\ & & & 0 & s_2 & \\ \hline & & & & s_2 & 0 \\ & & & & 0 & s_1 \\ \hline & & & & & s_1 \\ & & & & & i_{12} \\ \hline \end{array}$$

2.3 The triangular basis of $(\mathcal{K}, \mathcal{M})$

Definition 2.2.1 shows that $\mathcal{M}$ and $\text{rad}\mathcal{K}$ are solution spaces of the equation systems III and II respectively. Some nice triangular bases of the spaces will be chosen in this subsection.

Definition 2.3.1 Let $M = (m_{ij})_{t \times t}$ be a matrix. An order on the indices is defined as follows: $(i, j) \prec (i', j')$ provided that $i > i'$ or $i = i', j < j'$. The order is also valid on the indices of blocks of a $t \times t$ partitioned matrix.

In fact there are many possibilities to define an order on the indices of a $t \times t$ matrix. For example, $(i, j) \prec (i', j')$ if $j < j'$, or $j = j', i > i'$. Another example: $(i, j) \prec (i', j')$ if

$$i - j > i' - j', \text{ or } i - j = i' - j', \quad i > i'.$$

The principle is to ensure that (the index of $m_{ij}s_{jj'}$) $\succ$ (that of m_{ij}) when $j < j'$, and (the index of $s_{i'i}m_{ij}$) $\succ$ (that of m_{ij}) when $i' < i$ for any entry m_{ij} of M in $\mathcal{M}$, and $s_{jj'}$, $s_{i'i}$ of S in $\mathcal{K}$. In this paper we use mainly the order given by Definition 2.3.1, unless otherwise stated.

Lemma 2.3.1 Let $\sum_{j=1}^n d_{ij}z_j = 0, i = 1, 2, \dots, m$, be a system of homogeneous linear equations. Then there exists a unique choice of free indeterminates $z_{p_1}, \dots, z_{p_r}$ such that

$$\begin{cases} z_{j_1} & = \sum_{l=1}^r a_{1l}z_{p_l} \\ z_{j_2} & = \sum_{l=1}^r a_{2l}z_{p_l} \\ \ddots & \\ z_{j_{n-r}} & = \sum_{l=1}^r a_{n-r,l}z_{p_l} \end{cases}$$

where $p_1 < p_2 < \dots < p_r$, $j_1 < j_2 < \dots < j_{n-r}$, and $p_l < j_k$, whenever $a_{kl} \neq 0$ for each $1 \leq k \leq n - r$.

Proof. $n = 1$ is trivial. Suppose that the assertion is true for $(n - 1)$ indeterminates, and we are in the case of n indeterminates. If $z_1 = 0$, then replacing z_1 by 0, we obtain a new system with $(n - 1)$ indeterminates. If $z_1 \neq 0$, then z_1 is taken as a free indeterminate. Suppose that $z_1, z_2, \dots, z_{j_1-1}$ are free, but $z_{j_1} = \sum_{l=1}^{j_1-1} a_{1l}z_l$, then substituting z_{j_1} for $\sum_{l=1}^{j_1-1} a_{1l}z_l$ we obtain a new system with $(n - 1)$ indeterminates. Thus the assertion follows by induction. $\square$

The lemma allows us to choose some nice basis for $\mathcal{M}$ (resp. $\text{rad}\mathcal{K}$) according to the order given in Definition 2.3.1. Let

$$z_{p_{\mathcal{I}\mathcal{J}}, q_{\mathcal{I}\mathcal{J}}^1}, \dots, z_{p_{\mathcal{I}\mathcal{J}}^{r_{\mathcal{I}\mathcal{J}}}, q_{\mathcal{I}\mathcal{J}}^{r_{\mathcal{I}\mathcal{J}}}}$$

be all the free indeterminates of the equation system III of Definition 2.2.1 for any fixed

$(\mathcal{I}, \mathcal{J})$, such that $(p_{\mathcal{I}\mathcal{J}}^1, q_{\mathcal{I}\mathcal{J}}^1) \prec \cdots \prec (p_{\mathcal{I}\mathcal{J}}^{r_{\mathcal{I}\mathcal{J}}}, q_{\mathcal{I}\mathcal{J}}^{r_{\mathcal{I}\mathcal{J}}})$ and

$$z_{ij} = \sum_{w=1}^{r_{\mathcal{I}\mathcal{J}}} a_{\mathcal{I}\mathcal{J}}^w(i, j) z_{p_{\mathcal{I}\mathcal{J}}^w} q_{\mathcal{I}\mathcal{J}}^w,$$

where $(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w) \prec (i, j)$ whenever $a_{\mathcal{I}\mathcal{J}}^w(i, j) \neq 0$, for $i \in \mathcal{I}, j \in \mathcal{J}$. Define a set

$$\begin{aligned} A_{\mathcal{I}\mathcal{J}} &= \{\rho_{\mathcal{I}\mathcal{J}}^w = E_{p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w} + \sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} a_{\mathcal{I}\mathcal{J}}^w(i, j) E_{ij} \mid 1 \leq w \leq r_{\mathcal{I}\mathcal{J}}\}, \text{ and} \\ A &= \bigcup_{\mathcal{I}, \mathcal{J} \in T/\sim} A_{\mathcal{I}\mathcal{J}}. \end{aligned} \quad (1)$$

Then $A_{\mathcal{I}\mathcal{J}}$ is a k -basis of the solution space $E_{\mathcal{I}}\mathcal{M}E_{\mathcal{J}}$ of equation system III of Definition 2.2.1 for any fixed pair $(\mathcal{I}, \mathcal{J})$. Moreover A is a k -basis of $\mathcal{M}$.

Similarly we can choose a k -basis $B_{\mathcal{I}\mathcal{J}}$ of the solution space $E_{\mathcal{I}}(\text{rad}\mathcal{K})E_{\mathcal{J}}$ of the equation system II of Definition 2.2.1 for any fixed pair $(\mathcal{I}, \mathcal{J})$, and obtain a k -basis of $\text{rad}\mathcal{K}$:

$$\begin{aligned} B_{\mathcal{I}\mathcal{J}} &= \{\zeta_{\mathcal{I}\mathcal{J}}^w = E_{\bar{p}_{\mathcal{I}\mathcal{J}}^w, \bar{q}_{\mathcal{I}\mathcal{J}}^w} + \sum_{\mathcal{I} \ni i < j \in \mathcal{J}} b_{\mathcal{I}\mathcal{J}}^w(i, j) E_{ij} \mid 1 \leq w \leq \bar{r}_{\mathcal{I}\mathcal{J}}\}, \text{ and} \\ B &= \bigcup_{\mathcal{I}, \mathcal{J} \in T/\sim} B_{\mathcal{I}\mathcal{J}}. \end{aligned} \quad (2)$$

The pair (A, B) is said to be a *triangular basis* of bimodule problem $(\mathcal{K}, \mathcal{M})$.

2.4 An order on triangular basis

Definition 2.4.1 There is a natural linear order on $A : \rho_{\mathcal{I}\mathcal{J}}^w \prec \rho_{\mathcal{I}'\mathcal{J}'}^{w'}$ provided $(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w) \prec (p_{\mathcal{I}'\mathcal{J}'}^{w'}, q_{\mathcal{I}'\mathcal{J}'}^{w'})$. Let us take

$$(p, q) = (p_{\mathcal{P}\mathcal{Q}}^{w_0}, q_{\mathcal{P}\mathcal{Q}}^{w_0}) = \min\{(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w) \mid w = 1, \dots, r_{\mathcal{I}\mathcal{J}}, \forall (\mathcal{I}, \mathcal{J}) \in (T/\sim) \times (T/\sim)\},$$

and $\rho = \rho_{\mathcal{P}\mathcal{Q}}^{w_0}$, then $p \in \mathcal{P}, q \in \mathcal{Q}$. Similarly there is also a linear order on B .

Proposition 2.4.1 Let (A, B) be the triangular basis of a bimodule problem $(\mathcal{K}, \mathcal{M}, H)$. If the left and right module actions are

$$\begin{cases} \zeta_{\mathcal{I}\mathcal{L}}^u \rho_{\mathcal{L}\mathcal{J}}^v = \sum_w {}_l c_{\binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}}}^{\binom{w}{\mathcal{I}\mathcal{J}}} \rho_{\mathcal{I}\mathcal{J}}^w, \\ \rho_{\mathcal{I}\mathcal{L}}^u \zeta_{\mathcal{L}\mathcal{J}}^v = \sum_w {}_r c_{\binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}}}^{\binom{w}{\mathcal{I}\mathcal{J}}} \rho_{\mathcal{I}\mathcal{J}}^w \end{cases}$$

respectively, with the structure constants ${}_l c, {}_r c \in k$, then $\rho_{\mathcal{I}\mathcal{J}}^w \succ \rho_{\mathcal{L}\mathcal{J}}^v$ for the first case, and $\rho_{\mathcal{I}\mathcal{J}}^w \succ \rho_{\mathcal{I}\mathcal{L}}^u$ for the second case.

Proof. We will only prove the first one, and the second one can be obtained similarly.

$$\zeta_{\mathcal{IL}}^u \rho_{\mathcal{LJ}}^v = (E_{\bar{p}_{\mathcal{IL}}^u \bar{q}_{\mathcal{IL}}^u} + \sum_{\mathcal{I} \ni i < l \in \mathcal{L}} b_{\mathcal{IL}}^u(i, l) E_{il}) (E_{p_{\mathcal{LJ}}^v q_{\mathcal{LJ}}^v} + \sum_{(l', j) \in \mathcal{L} \times \mathcal{J}} a_{\mathcal{LJ}}^v(l', j) E_{l'j})$$

by Formulae (1) and (2) in 2.3. We may write

$$\zeta_{\mathcal{IL}}^u = \sum_{\mathcal{I} \ni i < l \in \mathcal{L}} b_{\mathcal{IL}}^u(i, l) E_{il} \quad \text{with} \quad b_{\mathcal{IL}}^u(\bar{p}_{\mathcal{IL}}^u, \bar{q}_{\mathcal{IL}}^u) = 1$$

and

$$\rho_{\mathcal{LJ}}^v = \sum_{(l', j) \in \mathcal{L} \times \mathcal{J}} a_{\mathcal{LJ}}^v(l', j) E_{l'j} \quad \text{with} \quad a_{\mathcal{LJ}}^v(p_{\mathcal{LJ}}^v, q_{\mathcal{LJ}}^v) = 1$$

for simplicity. Then

$$E_{il} E_{l'j} = \begin{cases} E_{ij} & \text{when } l = l', \\ 0 & \text{otherwise,} \end{cases}$$

since $\zeta_{\mathcal{IL}}^u$ is an upper triangular nilpotent matrix, $i < l$. Thus $(i, j) \succ (l', j) \succeq (p_{\mathcal{LJ}}^v, q_{\mathcal{LJ}}^v)$, i.e. the indices of E_{ij} in the expression of $\rho_{\mathcal{LJ}}^w$ in the first formula are all greater than $(p_{\mathcal{LJ}}^v, q_{\mathcal{LJ}}^v)$. Therefore $\rho_{\mathcal{ILJ}}^w \succ \rho_{\mathcal{LJ}}^v$ as required. $\square$

Let C, D be two partitioned matrices. We write

$$C \equiv_{\prec(p,q)} D \quad (\text{resp. } C \equiv_{\preceq(p,q)} D)$$

if (i, j) -blocks of C, D are the same for all $(i, j) \prec (p, q)$ (resp. $(i, j) \preceq (p, q)$).

Corollary 2.4.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, with a pair (p, q) given by Definition 2.4.1. Then

$$SH \equiv_{\prec(p,q)} HS$$

for any $S \in \mathcal{K}$.

Proof. Since $d(S) = SH - HS \in \mathcal{M}$ by IV of Definition 2.2.1, and all the entries, which are smaller than (p, q) in a matrix of $\mathcal{M}$, equal zero, so that $SH - HS \equiv_{\prec(p,q)} 0$, i.e. $SH \equiv_{\prec(p,q)} HS$. $\square$

Let $D = \text{Hom}_k(-, k)$ be the usual dual functor. Let $\mathcal{M}^* = D\mathcal{M}$, and $(\text{rad}\mathcal{K})^* = D(\text{rad}\mathcal{K})$ be the dual space of $\mathcal{M}$ and $\text{rad}\mathcal{K}$ respectively. Then we write their dual bases respectively by

$$A^* = \{(\rho_{\mathcal{ILJ}}^w)^* \mid 1 \leq w \leq r_{\mathcal{ILJ}}, \forall (\mathcal{I}, \mathcal{J}) \in (T/\sim) \times (T/\sim)\} \quad (3)$$

$$B^* = \{(\zeta_{\mathcal{ILJ}}^w)^* \mid 1 \leq w \leq \bar{r}_{\mathcal{ILJ}}, \forall (\mathcal{I}, \mathcal{J}) \in (T/\sim) \times (T/\sim)\} \quad (4)$$

thus $(\rho_{\mathcal{ILJ}}^w)^*$ can be regarded as the coefficient functions of $\rho_{\mathcal{ILJ}}^w$ in $\mathcal{M}$ and $(\zeta_{\mathcal{ILJ}}^w)^*$ as those of $\zeta_{\mathcal{ILJ}}^w$ in $\text{rad}\mathcal{K}$, which yield a dual structure of the bimodule problem (see 3.6 below).

The linear order on A is transferred to that on A^* , namely, $(\rho_{I\mathcal{J}}^w)^* \prec (\rho_{I'\mathcal{J}'}^{w'})^*$ provided $\rho_{I\mathcal{J}}^w \prec \rho_{I'\mathcal{J}'}^{w'}$. And B^* is also ordered such that $(\zeta_{I\mathcal{J}}^w)^* \prec (\zeta_{I'\mathcal{J}'}^{w'})^*$ provided $\zeta_{I\mathcal{J}}^w \prec \zeta_{I'\mathcal{J}'}^{w'}$.

Example. See Example 3 of 2.1, and Proposition 2.2.1. In the corresponding bimodule problem of Λ ,

$$\rho^1 = E_{4,10} + E_{18} = a, \quad \rho^2 = E_{3,10} + E_{28} + \alpha E_{19} - b, \quad \rho^3 = E_{2,10} = c, \quad \rho^4 = E_{1,10} = d;$$

and

$$(\rho^1)^* = a^*, \quad (\rho^2)^* = b^*, \quad (\rho^3)^* = c^*, \quad (\rho^4)^* = d^*,$$

if the k -basis of $\text{rad}(\Lambda)$, a, b, c, d still stand for the k -basis of $\text{rad}(\tilde{\Lambda})$, as well as $\begin{pmatrix} 0 & \text{rad}\tilde{\Lambda} \\ 0 & 0 \end{pmatrix}$ with $a \prec b \prec c \prec d$, and a^*, b^*, c^*, d^* are dual basis.

2.5 Weyr matrices

Let

$$J(\lambda) = J_d(\lambda)^{e_d} \oplus J_{d-1}(\lambda)^{e_{d-1}} \oplus \cdots \oplus J_1(\lambda)^{e_1}$$

be a direct sum of Jordan blocks with a common eigenvalue λ , where e_j stands for the number of the summands of $J_j(\lambda)$, write

$$m_j = e_d + e_{d-1} + \cdots + e_j$$

for $j = 1, 2, \dots, d$. Then $m_1 \geq m_2 \geq \cdots \geq m_d$, and $e_j = m_j - m_{j+1}$, $m_{d+1} = 0$. The following partitioned matrix W_λ is called a *Weyr matrix of eigenvalue λ* :

$$W_\lambda = \begin{pmatrix} \lambda I_{m_1} & W_{12} & 0 & \cdots & 0 & 0 \\ & \lambda I_{m_2} & W_{23} & \cdots & 0 & 0 \\ & & \lambda I_{m_3} & \ddots & 0 & 0 \\ & & & \ddots & \ddots & \vdots \\ & & & & \lambda I_{m_{d-1}} & W_{d-1,d} \\ & & & & & \lambda I_{m_d} \end{pmatrix}_{d \times d}, \quad W_{j,j+1} = \begin{pmatrix} I_{m_{j+1}} \\ 0 \end{pmatrix}_{m_j \times m_{j+1}}.$$

Lemma 2.5.1 W_λ is obtained from $J(\lambda)$ by a series of elementary transformations of exchanging rows and columns simultaneously. $\square$

The following is called a vector of W_λ :

$$\underline{e} = \begin{pmatrix} e_d & e_{d-1} & e_{d-2} & \cdots & e_3 & e_2 & e_1 \\ e_d & e_{d-1} & e_{d-2} & \cdots & e_3 & e_2 & \\ & \cdots & & \cdots & & & \\ e_d & e_{d-1} & e_{d-2} & & & & \\ e_d & e_{d-1} & & & & & \\ e_d & & & & & & \end{pmatrix}. \quad (5)$$

For example, given a Jordan form $J_4(\lambda)^2 \oplus J_2(\lambda)^3$, we have $m_1 = 2 + 0 + 3 + 0 = 5$, $m_2 = 2 + 0 + 3 = 5$, $m_3 = 2 + 0 = 2$, $m_4 = 2$,

$$\underline{e} = \begin{pmatrix} 2 & 0 & 3 & 0 \\ 2 & 0 & 3 & \\ 2 & 0 & & \\ 2 & & & \end{pmatrix}, \quad W_\lambda = \begin{pmatrix} \boxed{\lambda I_5} & \boxed{I_5} & & \\ & \boxed{\lambda I_5} & \boxed{I_2} & \\ & & \boxed{\lambda I_2} & \boxed{I_2} \\ & & & \boxed{\lambda I_2} \end{pmatrix}.$$

Let $\lambda^1, \lambda^2, \dots, \lambda^\alpha$ be a set of eigenvalues which are pairwise different, then

$$W = \text{diag}(W_{\lambda^1}, W_{\lambda^2}, \dots, W_{\lambda^\alpha})$$

is called a *Weyr matrix* with a vector

$$\underline{e} = (\underline{e}^1, \underline{e}^2, \dots, \underline{e}^\alpha),$$

where $\underline{e}^l$ is a vector of W_{λ^l} defined in Formula (5). From now on we will fix an order on the base field k , such that

$$\lambda^1 < \lambda^2 < \dots < \lambda^\alpha.$$

For example, if k is the field of complex numbers, we may use the lexicographic order: $a + bi < c + di$ provided $a < c$ or $a = c$ but $b < d$.

Corollary 2.5.1 Any square matrix over k is similar to a unique Weyr matrix under a fixed order of k . $\square$

Let W be a Weyr matrix, X be a square matrix having the same size as W . Then an equation $WX = XW$ gives

$$X = \text{diag}(X_1, \dots, X_l, \dots, X_\alpha).$$

Denote X_l by Y , and d_l by d , then $Y = (Y_{ij})_{d \times d}$, where $Y_{ij} = 0$ when $i > j$, and

$$Y_{ij} = \begin{pmatrix} U_{11}^h & U_{12}^h & \cdots & U_{1j'}^h \\ & \cdots & & \cdots \\ U_{h1}^h & U_{h2}^h & \cdots & U_{hj'}^h \\ & U_{h+1,2}^h & \cdots & U_{h+1,j'}^h \\ & & \ddots & \vdots \\ & & & U_{i'j'}^h \end{pmatrix}_{i' \times j'}$$

when $i \leq j$, where $h = j - i + 1$, $i' = d - i + 1$, $j' = d - j + 1$, and U_{gr}^h are $e_{g'} \times e_{r'}$ free matrices, where $g' = d - g + 1$, $r' = d - r + 1$. For example, if $W = W_\lambda$ is given above,

then

$$X = \begin{pmatrix} \boxed{\begin{matrix} U_{11}^1 & U_{13}^1 \\ & U_{33}^1 \end{matrix}} & \boxed{\begin{matrix} U_{11}^2 & U_{13}^2 \\ & 0 \end{matrix}} & \boxed{\begin{matrix} U_{11}^3 \\ U_{31}^3 \end{matrix}} & \boxed{\begin{matrix} U_{11}^4 \\ U_{31}^4 \end{matrix}} \\ & \boxed{\begin{matrix} U_{11}^1 & U_{13}^1 \\ & U_{33}^1 \end{matrix}} & \boxed{\begin{matrix} U_{11}^2 \\ 0 \end{matrix}} & \boxed{\begin{matrix} U_{11}^3 \\ U_{31}^3 \end{matrix}} \\ & & \boxed{\begin{matrix} U_{11}^1 \\ U_{11}^2 \end{matrix}} & \\ & & & \boxed{U_{11}^1} \end{pmatrix}_{4 \times 4}.$$

2.6 Reductions

Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, (A, B) be the triangular basis, and (A^*, B^*) the dual basis. Define two matrices

$$R_0 = \sum_{\mathcal{I}} (E_{\mathcal{I}})^* \otimes_k E_{\mathcal{I}} + \sum_{(\mathcal{I}, \mathcal{J})} \sum_{w=1}^{\bar{r}_{\mathcal{I}\mathcal{J}}} (\zeta_{\mathcal{I}\mathcal{J}}^w)^* \otimes_k \zeta_{\mathcal{I}\mathcal{J}}^w,$$

$$N_0 = \sum_{(\mathcal{I}, \mathcal{J})} \sum_{w=1}^{r_{\mathcal{I}\mathcal{J}}} (\rho_{\mathcal{I}\mathcal{J}}^w)^* \otimes \rho_{\mathcal{I}\mathcal{J}}^w.$$

where the tensor product is the usual tensor product of matrices, i.e if

$$C = (c_{ij})_{p \times q}, \quad D = (d_{ij})_{t \times s}, \quad \text{then} \quad C \otimes_k D = (C \cdot d_{ij})_{t \times s}.$$

And we regard the coefficient functions $(E_{\mathcal{I}})^*$, $(\zeta_{\mathcal{I}\mathcal{J}}^w)^*$, $(\rho_{\mathcal{I}\mathcal{J}}^w)^*$ as indeterminates. Clearly, if the values of

$$((E_{\mathcal{I}})^*, (\zeta_{\mathcal{I}\mathcal{J}}^w)^* \mid \forall \mathcal{I}, 1 \leq w \leq \bar{r}_{\mathcal{I}\mathcal{J}}, (\mathcal{I}, \mathcal{J}) \in T / \sim \times T / \sim)$$

range over $k^{\dim \mathcal{K}}$, we obtain the algebra $\mathcal{K}$; and if the values of

$$((\rho_{\mathcal{I}\mathcal{J}}^w)^* \mid 1 \leq w \leq r_{\mathcal{I}\mathcal{J}}, (\mathcal{I}, \mathcal{J}) \in T / \sim \times T / \sim)$$

run over $k^{\dim \mathcal{M}}$, we obtain the bimodule $\mathcal{M}$. The $(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w)$ -th elements of N_0 are said to be *free*, and the others are said to be *dependent* (see Formula (1) of 2.3).

Next we fix a size vector $\underline{n}$ of $(T, \sim)$, consider the partitioned upper triangular matrix algebra $\mathcal{K}_{\underline{n} \times \underline{n}}$, bimodule $\mathcal{M}_{\underline{n}}$ and matrix $H_{\underline{n}}$. it is obvious that $\mathcal{K}_{\underline{n} \times \underline{n}}$ is Morita equivalent

to $\mathcal{K}$ if $n_{\mathcal{I}} \neq 0, \forall \mathcal{I} \in T/\sim$; and $\mathcal{K}_{\underline{n} \times \underline{n}}$ is not basic if there exists some $\mathcal{I} \in T/\sim$, such that $n_{\mathcal{I}} > 1$. Let

$$\begin{aligned} R &= \sum_{\mathcal{I}} (\overline{E_{\mathcal{I}}})^* \otimes E_{\mathcal{I}} + \sum_{(\mathcal{I}, \mathcal{J})} \sum_{w=1}^{\bar{r}_{\mathcal{I}\mathcal{J}}} (\overline{\zeta_{\mathcal{I}\mathcal{J}}^w})^* \otimes \zeta_{\mathcal{I}\mathcal{J}}^w, \\ N &= \sum_{(\mathcal{I}, \mathcal{J})} \sum_{w=1}^{r_{\mathcal{I}\mathcal{J}}} (\overline{\rho_{\mathcal{I}\mathcal{J}}^w})^* \otimes \rho_{\mathcal{I}\mathcal{J}}^w, \end{aligned} \quad (6)$$

where $(\overline{E_{\mathcal{I}}})^*$, $(\overline{\zeta_{\mathcal{I}\mathcal{J}}^w})^*$ and $(\overline{\rho_{\mathcal{I}\mathcal{J}}^w})^*$ are $n_{\mathcal{I}} \times n_{\mathcal{I}}$, $n_{\mathcal{I}} \times n_{\mathcal{J}}$ and $n_{\mathcal{I}} \times n_{\mathcal{J}}$ matrices with all the entries being indeterminates respectively. It is clear that if the values of $(\overline{E_{\mathcal{I}}})^*$ and $(\overline{\zeta_{\mathcal{I}\mathcal{J}}^w})^*$ range over

$$\prod_{\mathcal{I} \in T/\sim} M_{n_{\mathcal{I}}}(k) \times \prod_{\mathcal{I}, \mathcal{J} \in T/\sim} M_{n_{\mathcal{I}} \times n_{\mathcal{J}}}(k)^{\bar{r}_{\mathcal{I}\mathcal{J}}},$$

we obtain $\mathcal{K}_{\underline{n} \times \underline{n}}$; and if the values of $(\overline{\rho_{\mathcal{I}\mathcal{J}}^w})^*$ run over

$$\prod_{\mathcal{I}, \mathcal{J} \in T/\sim} M_{n_{\mathcal{I}} \times n_{\mathcal{J}}}(k)^{r_{\mathcal{I}\mathcal{J}}},$$

we obtain $\mathcal{M}_{\underline{n}}$. And $(N, R, \underline{n})$ is called a *triple of bimodule problem* $(\mathcal{K}, \mathcal{M}, H)$. The $(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w)$ -th blocks of N are said to be *free*, and the others are said to be *dependent* (see Formula (1) of 2.3).

The matrix N_0 may be regarded as a “universal matrix” of $\mathcal{M}$, and R_0 a “universal matrix” of $\mathcal{K}$ or a “universal endomorphism” of N_0 . Similarly, N may be regarded as a “universal matrix” of $\mathcal{M}_{\underline{n}}$, and R a “universal matrix” of $\mathcal{K}_{\underline{n}}$ or a “universal endomorphism” of N . Consider the matrix equation

$$(N + H_{\underline{n}})R = R(N + H_{\underline{n}}).$$

Let (p, q) be as in Definition 2.4.1, then N_{pq} is the first non-zero block of N . Then Corollary 2.4.1 ensures that $H_{\underline{n}}R \equiv_{\prec(p, q)} RH_{\underline{n}}$, i.e. the blocks before (p, q) are really equal on both sides. And the (p, q) -block of the equation is

$$\begin{aligned} & (H_{\underline{n}})_{p1}X_{1q} + \cdots + (H_{\underline{n}})_{p, q-1}X_{q-1, q} + (H_{\underline{n}})_{pq}X_{qq} + N_{pq}X_{qq} \\ &= X_{pp}N_{pq} + X_{pp}(H_{\underline{n}})_{pq} + X_{p, p+1}(H_{\underline{n}})_{p+1, q} + \cdots + X_{pt}(H_{\underline{n}})_{tq}. \end{aligned}$$

Since $(H_{\underline{n}})_{ij} = 0$ when $i \not\sim j$, $(H_{\underline{n}})_{ij} = h_{ij}I_{n_i}$ when $i \sim j$ by IV of Definition 2.2.1, and $(H_{\underline{n}})_{pq}X_{qq} = X_{pp}(H_{\underline{n}})_{pq}$ in both cases $p \approx q$ and $p \sim q$, the above equation is equivalent to the following:

$$N_{pq}X_{qq} - X_{pp}N_{pq} = \sum_{l=p+1}^t X_{pl}h_{lq} - \sum_{l=1}^{q-1} h_{pl}X_{lq}. \quad (7)$$

Let us define a new equation

$$\sum_{l=p+1}^t x_{pl} h_{lq} - \sum_{l=1}^{q-1} h_{pl} x_{lq} = 0. \quad (8)$$

And define a constant matrix $\overline{N}_{pq}$ with the same size of N_{pq} in the following 3 cases [D, CB1, S]:

Regularization. If the equation system II of Definition 2.2.1 at $(\mathcal{P}, \mathcal{Q})$ do not imply equation (8), then let $\overline{N}_{pq} = 0$. And denote zero here by $\emptyset$, i.e. $\overline{N}_{pq} = \emptyset$, to distinguish 0's coming from regularization or from other cases.

If the equation system II of Definition 2.2.1 at $(\mathcal{P}, \mathcal{Q})$ implies equation (8), then

$$N_{pq} X_{qq} = X_{pp} N_{pq}. \quad (9)$$

Edge reduction. When $p \approx q$, let

$$\overline{N}_{pq} = \begin{pmatrix} 0 & I_r \\ 0 & 0 \end{pmatrix}_{n_p \times n_q}$$

for some $r \leq \min\{n_p, n_q\}$.

Loop reduction. When $p \sim q$, then $X_{pp} = X_{qq}$, let

$$\overline{N}_{pq} = W$$

for some Weyr matrix W .

As soon as $\overline{N}_{pq}$ has been fixed, we obtain a new triple $(N', R', \underline{n}')$ as follows. We first define a new vector $\underline{n}'$ according to the 3 cases respectively. Regularization: $\underline{n}' = \underline{n}$; Edge reduction: $\underline{n}'$ is obtained from $\underline{n}$ by splitting each component $n_i, \forall i \in \mathcal{P}$, into (n_{i_1}, n_{i_2}) when $n_{i_1} = r, n_{i_2} = n_{\mathcal{P}} - r$; and $n_j, \forall j \in \mathcal{Q}$, into (n_{j_1}, n_{j_2}) when $n_{j_1} = n_{\mathcal{Q}} - r, n_{j_2} = r$; Loop reduction: $\underline{n}'$ is obtained from $\underline{n}$ by splitting each component $n_i, \forall i \in \mathcal{P}$, into $\underline{e} = (\underline{e}^1, \dots, \underline{e}^l, \dots, \underline{e}^\alpha)$ given in 2.5. Let

$$\begin{cases} (H_{\underline{n}})' = H_{\underline{n}} + \overline{N}_{pq} \otimes \rho, \\ N' = N - \overline{N}_{pq} \otimes \rho, \\ R' \text{ is a restriction of } R \text{ given by } R'(H_{\underline{n}})' \equiv_{\preceq(p,q)} (H_{\underline{n}})' R' \end{cases} \quad (10)$$

Then R' satisfies equation system II of Definition 2.2.1 and equation (7), if we replace N_{pq} by $\overline{N}_{pq}$. R' determines a partitioned upper triangular matrix algebra $(\mathcal{K}_{\underline{n}})'$, which is not basic in general. Similarly N' determines a $(\mathcal{K}_{\underline{n}})'$ -($\mathcal{K}_{\underline{n}}')'$ -bimodule $(\mathcal{M}_{\underline{n}})'$. The procedure

of regularization, edge reduction and loop reduction is called a *reduction for the triple* $(N, R, \underline{n})$.

We will construct an induced bimodule problem $(\mathcal{K}', \mathcal{M}', H')$ from $(\mathcal{K}, \mathcal{M}, H)$ given by each reduction for $(N, R, \underline{n})$, such that $\mathcal{K}'$ is basic in 3.1. we will also show in 3.1 that

$$(\mathcal{K}_{\underline{n}})' = \mathcal{K}'_{\underline{n}'}, \quad (\mathcal{M}_{\underline{n}})' = \mathcal{M}'_{\underline{n}'} \quad \text{and} \quad (H_{\underline{n}})' = H'_{\underline{n}'}.$$

Thus $(N', R', \underline{n}')$ is a triple of $(\mathcal{K}', \mathcal{M}', H')$.

3 Induced bimodule problems

In this section we will describe the induced bimodule problem $(\mathcal{K}', \mathcal{M}', H')$ given by one of three reductions respectively, and give a nice functor $\vartheta : \text{Mat}(\mathcal{K}', \mathcal{M}') \rightarrow \text{Mat}(\mathcal{K}, \mathcal{M})$. In the end, we will present an explicit correspondence between bimodule problems and their dual structure — bocses. In particular we show the differentials of the arrows of a bocs in terms of the multiplication of matrices.

3.1 Induced bimodule problems

Now we construct an induced bimodule problem $(\mathcal{K}', \mathcal{M}', H')$ from the bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ after one of the three reductions given in 2.6 respectively.

Regularization. I. Let $T' = T$, $\sim' = \sim$. Then $\underline{n}'$ is a size vector of $(T', \sim')$.

II. $\mathcal{K}' = \{S' \in \mathcal{K} \mid S' \text{ satisfies the equation (8) of 2.6}\}$.

III. $\mathcal{M}' = \{M' \in \mathcal{M} \mid M' \text{ satisfies the equation } z_{pq} = 0\}$, which is a $\mathcal{K}'$ - $\mathcal{K}'$ -bimodule by Proposition 2.4.1.

IV. $H' = H + \emptyset \otimes \rho$ which implies $H'_{\underline{n}'} = (H_{\underline{n}})'$ and $(H'_{\underline{n}'})_{pq} = \overline{N}_{pq}$.

Edge reduction. The equation at the (p, q) -block of $R'(H_{\underline{n}})' \equiv_{\preceq(p, q)} (H_{\underline{n}})'R'$ is as follows:

$$\begin{pmatrix} 0 & I_r \\ 0 & 0 \end{pmatrix} X'_{qq} = X'_{pp} \begin{pmatrix} 0 & I_r \\ 0 & 0 \end{pmatrix},$$

which yields

$$X'_{pp} = \begin{pmatrix} X_{p_1 p_1} & X_{p_1 p_2} \\ 0 & X_{p_2 p_2} \end{pmatrix}, \quad X'_{qq} = \begin{pmatrix} X_{q_1 q_1} & X_{q_1 q_2} \\ 0 & X_{q_2 q_2} \end{pmatrix},$$

and $X_{p_1 p_1} = X_{q_2 q_2}$.

I. Define a size vector $\tilde{n} \in (T, \sim)$ by $\tilde{n}_{\mathcal{I}} = 1, \forall \mathcal{I} \in T/\sim \setminus \{\mathcal{P}, \mathcal{Q}\}$; and (i) $\tilde{n}_{\mathcal{P}} = \tilde{n}_{\mathcal{Q}} = 2$ when $\min\{n_{\mathcal{P}}, n_{\mathcal{Q}}\} > r > 0$; (ii) $\tilde{n}_{\mathcal{P}} = 2, \tilde{n}_{\mathcal{Q}} = 1$ when $n_{\mathcal{P}} > r, n_{\mathcal{Q}} = r > 0$; (iii) $\tilde{n}_{\mathcal{P}} = 1, \tilde{n}_{\mathcal{Q}} = 2$ when $n_{\mathcal{P}} = r > 0, n_{\mathcal{Q}} > r$; (iv) $\tilde{n}_{\mathcal{P}} = \tilde{n}_{\mathcal{Q}} = 1$ when $n_{\mathcal{P}} = n_{\mathcal{Q}} = r > 0$; (v) $\tilde{n}_{\mathcal{P}} = \tilde{n}_{\mathcal{Q}} = 1$ when $r = 0$. Let T' be obtained from T by splitting $i, \forall i \in \mathcal{P}$, into i_1, i_2 in cases (i) and (ii), into i_1 in cases (iii) and (iv), into i_2 in case (v), and splitting $j, \forall j \in \mathcal{Q}$, into j_1, j_2 in cases (i) and (iii), into j_2 in cases (ii) and (iv), into j_1 in case (v). Define

$$(\mathcal{P} \cup \mathcal{Q})' = \{i_1, j_2 \mid \forall i \in \mathcal{P}, \forall j \in \mathcal{Q}\},$$

$$\mathcal{P}' = \{i_2 \mid \forall i \in \mathcal{P}\} \text{ and } \mathcal{Q}' = \{j_1 \mid \forall j \in \mathcal{Q}\}.$$

Then

$$T'/\sim' = (T/\sim \setminus \{\mathcal{P}, \mathcal{Q}\}) \cup \{\mathcal{P}', \mathcal{Q}', (\mathcal{P} \cup \mathcal{Q})'\}.$$

$\underline{n}'$ is a size vector of $(T', \sim')$.

II. $\mathcal{K}' = \{S' \in \mathcal{K}_{\tilde{n} \times \tilde{n}} \mid S' \text{ satisfies the following additional equations}\}$:

$$x_{i_1 i_2} - x_{p_1 p_2} = 0, \quad \forall i \in \mathcal{P} \text{ and } x_{j_1 j_2} - x_{q_1 q_2} = 0, \quad \forall j \in \mathcal{Q}.$$

III. $\mathcal{M}' = \{M' \in \mathcal{M}_{\tilde{n}} \mid M' \text{ satisfies } z_{pq} = 0\}$, which is a $\mathcal{K}'$ - $\mathcal{K}'$ -bimodule by Proposition 2.4.1.

IV. $H' = H_{\tilde{n}} + U \otimes \rho$, where

$$U = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (0 \quad 1), \quad (1), \quad (0)$$

according to cases (i)–(v) of $\tilde{n}$ respectively. $H'_{\underline{n}'} = (H_{\underline{n}})'$ and $(H'_{\underline{n}'})_{pq} = \overline{N}_{pq}$.

Loop reduction. The equation

$$W X'_{qq} = X'_{pp} W$$

gives a new index set. Namely we define a linearly ordered set of indices according to $\underline{e} = (\underline{e}^1, \dots, \underline{e}^l, \dots, \underline{e}^\alpha)$ in order to define T' :

$$\underline{\delta}^{il} = \begin{pmatrix} \delta_{1,d}^{il} & \delta_{1,d-1}^{il} & \delta_{1,d-2}^{il} & \cdots & \delta_{1,3}^{il} & \delta_{1,2}^{il} & \delta_{1,1}^{il} \\ \delta_{2,d}^{il} & \delta_{2,d-1}^{il} & \delta_{2,d-2}^{il} & \cdots & \delta_{2,3}^{il} & \delta_{2,2}^{il} & \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \\ \delta_{d-2,d}^{il} & \delta_{d-2,d-1}^{il} & \delta_{d-2,d-2}^{il} & & & & \\ \delta_{d-1,d}^{il} & \delta_{d-1,d-1}^{il} & & & & & \\ \delta_{d,d}^{il} & & & & & & \end{pmatrix}$$

for each $i \in \mathcal{P}$, $1 \leq l \leq \alpha$. For example, the row and column indices of the blocks of X given in example of 2.5 can be expressed by the elements of δ in the table below:

$\begin{matrix} \text{column} \\ h \\ \text{row} \end{matrix}$	δ_{14}	δ_{12}	δ_{24}	δ_{22}	δ_{34}	δ_{44}
δ_{14}	1	1	2	2	3	4
δ_{12}		1	2	2	3	4
δ_{24}			1	1	2	3
δ_{22}				1	2	3
δ_{34}					1	2
δ_{44}						1

I. T' is obtained from T by splitting $i \in \mathcal{P}$ into

$$\{\delta_{gr}^{il} \mid e_r^l \neq 0, 0 \leq g \leq r; r = d_l, d_l - 1, \dots, 2, 1; l = 1, 2, \dots, \alpha\}.$$

where e_r^l is the $(d_l - r + 1)$ -th component of $\underline{e}^l$. And

$$T'/\sim' = (T/\sim \setminus \mathcal{P}) \cup \{\mathcal{P}_r^l \mid 1 \leq r \leq d_l, l = 1, 2, \dots, \alpha\},$$

where $\mathcal{P}_r^l = \{\delta_{gr}^{il} \in T' \mid 1 \leq g \leq r, \forall i \in \mathcal{P}\}$. Then $\underline{n}'$ is a size vector of $(T', \sim')$.

Moreover we define a size vector $\tilde{\underline{n}}$ of $(T, \sim)$ with $\tilde{n}_{\mathcal{I}} = 1, \forall \mathcal{I} \in T/\sim \setminus \{\mathcal{P}\}$, and

$$\tilde{n}_{\mathcal{P}} = \#\{\delta_{gr}^{pl} \in T'\} = \sum_{l=1}^{\alpha} \sum_{r=1}^{d_l} \delta_r^l \cdot r,$$

where $\delta_r^l = 1$, if $e_r^l \neq 0$, and $\delta_r^l = 0$, if $e_r^l = 0$.

II. $\mathcal{K}' = \{S' \in \mathcal{K}_{\tilde{\underline{n}} \times \tilde{\underline{n}}} \mid S' \text{ satisfies the following additional equations}\}$:

$$x_{\delta_{g_1' r}^{il} \delta_{g_2' s}^{il}} - x_{\delta_{g_1' r}^{pl} \delta_{g_2' s}^{pl}} = 0,$$

where $g_2' - g_1' = g_2 - g_1 = h - 1$, $s - r < h$, $g_1, g_1' \leq r$, $g_2, g_2' \leq s$, $1 \leq r, s \leq d_l$, $l = 1, 2, \dots, \alpha$, $\forall i \in \mathcal{P}$; and

$$x_{\delta_{g_1' r}^{il} \delta_{g_2' s}^{il}} = 0$$

where $g_2 - g_1 = h - 1$, $s - r \geq h$, $g_1 \leq r$, $g_2 \leq s$, $1 \leq r, s \leq d_l$, $l = 1, 2, \dots, \alpha$, $\forall i \in \mathcal{P}$.

III. $\mathcal{M}' = \{M' \in \mathcal{M}_{\tilde{\underline{n}}} \mid M' \text{ satisfies } z_{pq} = 0\}$ which is a $\mathcal{K}'$ - $\mathcal{K}'$ -bimodule by Proposition 2.4.1.

IV. $H' = H_{\tilde{\underline{n}}} + \widetilde{W} \otimes \rho$ with $\widetilde{W}$ being a Weyr matrix similar to $\bigoplus_{l=1}^{\alpha} \bigoplus_{r=d_l}^1 J_r(\lambda_l)^{\delta_r^l}$, where δ_r^l are as in the end of I. Thus $H'_{\underline{n}'} = (H_{\underline{n}})'$ and $(H'_{\underline{n}'})_{pq} = \overline{N}_{pq}$.

Corollary 3.1.1 $n'_{\mathcal{P}_r^l} = e_r^l$ where $1 \leq r \leq d_l$, $l = 1, \dots, \alpha$. □

Proposition 3.1.1. In the above 3 procedures, H' satisfies IV of Definition 2.2.1. Let $d'(S') = S'H' - H'S'$ for any $S' \in \mathcal{K}'$, we obtain a derivation from $\mathcal{K}'$ to $\mathcal{M}'$. Thus $(\mathcal{K}', \mathcal{M}', H')$ is a bimodule problem, and $(N', R', \underline{n}')$ is a triple of $(\mathcal{K}', \mathcal{M}', H')$.

Proof. $S'H' \equiv_{\leq(p,q)} H'S'$ implies that $d'(S') \in \mathcal{M}'$. □

The procedure of regularization, edge reduction or loop reduction given above is called a *reduction of bimodule problem* $(\mathcal{K}, \mathcal{M}, H)$, and $(\mathcal{K}', \mathcal{M}', H')$ is called an *induced bimodule problem* of $(\mathcal{K}, \mathcal{M}, H)$. There is a map

$$(\vartheta_1, \vartheta_0) : (\mathcal{K}', \mathcal{M}') \rightarrow (\mathcal{K}_{\tilde{\underline{n}} \times \tilde{\underline{n}}}, \mathcal{M}_{\tilde{\underline{n}}}),$$

such that the algebra homomorphism $\vartheta_1 : \mathcal{K}' \rightarrow \mathcal{K}_{\tilde{\underline{n}} \times \tilde{\underline{n}}}$, and the $\mathcal{K}'$ - $\mathcal{K}'$ -bimodule homomorphism $\vartheta_0 : \mathcal{M}' \rightarrow \mathcal{M}_{\tilde{\underline{n}}}$ are both natural embeddings.

3.2 Deletions and size changes

In this subsection we first define the 4-th reduction algorithm, so called *deletion*. Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, $\underline{n}$ a size vector of $(T, \sim)$ with $\tilde{n}_{\mathcal{I}} = 1$ or 0. Let

$$T' = \{i \mid \tilde{n}_i = 1\}, \quad \sim' = \sim|_{T'}, \quad \mathcal{K}' = \mathcal{K}_{\underline{n} \times \underline{n}}, \quad \mathcal{M}' = \mathcal{M}_{\underline{n}}, \quad H' = H_{\underline{n}}.$$

Then we obtain an induced bimodule problem $(\mathcal{K}', \mathcal{M}', H')$ from $(\mathcal{K}, \mathcal{M}, H)$ by *deleting a subset of equivalent classes* $\{\mathcal{I} \mid \tilde{n}_{\mathcal{I}} = 0\}$ from $T/\sim$. And we have a map

$$(\vartheta_1, \vartheta_0) : (\mathcal{K}', \mathcal{M}') \rightarrow (\mathcal{K}, \mathcal{M}),$$

such that the algebra homomorphism $\vartheta_1 : \mathcal{K}' \rightarrow \mathcal{K}$, and the $\mathcal{K}'$ - $\mathcal{K}'$ -bimodule homomorphism $\vartheta_0 : \mathcal{M}' \rightarrow \mathcal{M}$ are both natural embeddings.

The following observations are straightforward.

(1) Given any size vector $\underline{n}$ of $(T, \sim)$, we define a size vector $\underline{\tilde{n}}$ according to $\underline{n}$ with $\tilde{n}_{\mathcal{I}} = 1$ when $n_{\mathcal{I}} \neq 0$, and $\tilde{n}_{\mathcal{I}} = 0$ when $n_{\mathcal{I}} = 0$. Then we obtain an induced bimodule problem $(\mathcal{K}', \mathcal{M}', H')$. Let $\underline{n}' \in (T', \sim')$ with $n'_i = n_i, \forall i \in T'$, and a triple $(N', R', \underline{n}')$ over $(\mathcal{K}', \mathcal{M}', H')$ be given by Formula (6) of 2.6. If $(N, R, \underline{n})$ is a triple of $(\mathcal{K}, \mathcal{M})$, then

$$N' = N, \quad R' = R, \quad H' = H,$$

i.e. the two triples are essentially the same.

(2) A size vector $\underline{n}$ of $(T, \sim)$ is said to be *sincere* if $n_{\mathcal{I}} \neq 0$ for all $\mathcal{I} \in T/\sim$. In most considerations, a size vector is usually assumed to be sincere by a deletion if necessary.

(3) If $\underline{n}$ is sincere, then after doing anyone of the 3 reductions given in 3.1, the induced size vector $\underline{n}'$ is still sincere.

Next we make some size changes of triples over the induced bimodule problem.

Proposition 3.2.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, $(N, R, \underline{n})$ be a triple of $(\mathcal{K}, \mathcal{M}, H)$, and $(N', R', \underline{n}')$ be induced from $(N, R, \underline{n})$ by one of the 3 reductions in 3.1. Then for any size vector $\underline{\hat{n}}'$ of $(T', \sim')$, there exist a size vector $\underline{\hat{n}}$ of $(T, \sim)$ thus a triple $(\hat{N}, \hat{R}, \underline{\hat{n}})$ of $(\mathcal{K}, \mathcal{M}, H)$, such that

(1) If $\hat{n}_{\mathcal{P}} = 0$ or $\hat{n}_{\mathcal{Q}} = 0$, then $(\hat{N}', \hat{R}', \underline{\hat{n}}')$ is obtained from $(\hat{N}, \hat{R}, \underline{\hat{n}})$ by identity.

(2) If $\hat{n}_{\mathcal{P}} \neq 0$ and $\hat{n}_{\mathcal{Q}} \neq 0$, then the equation system II of Definition 2.2.1 and equation (8) of 2.6 for $(\mathcal{K}, \mathcal{M}, H)$ at the pair $(\mathcal{P}, \mathcal{Q})$ determine the same type of reductions from

$(N, R, \underline{n})$ to $(N', R', \underline{n}')$ and from $(\hat{N}, \hat{R}, \hat{\underline{n}})$ to $(\hat{N}', \hat{R}', \hat{\underline{n}}')$, where

$$\widehat{N}_{pq} = (H'_{\hat{\underline{n}}})_{pq} = \begin{cases} \emptyset, & \text{in regularization;} \\ \begin{pmatrix} 0 & \hat{I} \\ 0 & 0 \end{pmatrix}, & \text{in edge reduction;} \\ \hat{W}, & \text{in loop reduction.} \end{cases}$$

Proof. The size vector $\hat{\underline{n}}$ is defined from $\hat{\underline{n}}'$ according to the 3 reductions respectively.

$$\begin{cases} \text{regularization:} & \hat{\underline{n}} = \hat{\underline{n}}'; \\ \text{edge reduction:} & \hat{n}_{\mathcal{I}} = \hat{n}'_{\mathcal{I}}, \forall \mathcal{I} \in T/\sim \setminus \{\mathcal{P}, \mathcal{Q}\}, \\ & \hat{n}_{\mathcal{P}} = \hat{n}'_{\mathcal{P}'} + \hat{n}'_{(\mathcal{P} \cup \mathcal{Q})'}, \hat{n}_{\mathcal{Q}} = \hat{n}'_{(\mathcal{P} \cup \mathcal{Q})'} + \hat{n}'_{\mathcal{Q}'}; \\ \text{Loop reduction:} & \hat{n}_{\mathcal{I}} = \hat{n}'_{\mathcal{I}}, \forall \mathcal{I} \in T/\sim \setminus \{\mathcal{P}\}, \hat{n}_{\mathcal{P}} = \sum_{l,r} r \hat{n}'_{\mathcal{P}_r^l}. \end{cases} \quad (11)$$

Let $C = \{\mathcal{I} \in T/\sim \mid \hat{n}_{\mathcal{I}} = 0\}$. When $\mathcal{P} \in C$, or $\mathcal{Q} \in C$, we set $\hat{\underline{n}} = \hat{\underline{n}}'$, then

$$\hat{H}_{\hat{\underline{n}}} = \hat{H}'_{\hat{\underline{n}}'}, \quad \hat{N} = \hat{N}', \quad \hat{R} = \hat{R}'.$$

Thus the procedure from $(\hat{N}, \hat{R}, \hat{\underline{n}})$ to $(\hat{N}', \hat{R}', \hat{\underline{n}}')$ is an identity. When $\mathcal{P} \notin C$ and $\mathcal{Q} \notin C$, the equation system II of Definition 2.2.1 and (8) of 2.6 for $(\mathcal{K}, \mathcal{M}, H)$ at the pair $(\mathcal{P}, \mathcal{Q})$ allow us to make a reduction from $(\hat{N}, \hat{R}, \hat{\underline{n}})$ to $(\hat{N}', \hat{R}', \hat{\underline{n}}')$ in the same way as that from $(N, R, \underline{n})$ to $(N', R', \underline{n}')$. $\square$

The procedure given in the proposition is called a *size change*.

3.3 Reduction functors and reduction sequences

Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem and $(\mathcal{K}', \mathcal{M}', H')$ be an induced bimodule problem given by one of the 4 reductions. There is a nice functor

$$\vartheta : \text{Mat}(\mathcal{K}', \mathcal{M}') \longrightarrow \text{Mat}(\mathcal{K}, \mathcal{M})$$

which is induced from $(\vartheta_1, \vartheta_0)$ given in 3.1 and 3.2 and is defined as follows. For any size vector $\underline{m}'$ of $(T', \sim')$ and any matrix $M' \in \mathcal{M}'_{\underline{m}'}$, let

$$\vartheta(M') = M' + (H'_{\underline{m}'})_{pq} \otimes \rho$$

in case of regularization, edge or loop reductions. We write $\hat{\underline{n}}' = \underline{m}'$, and let $\hat{\underline{n}} = \underline{m}$ be calculated by Formula (11) of 3.2. In case of deletion let

$$\vartheta(M') = M', \quad \underline{m} = \underline{m}'.$$

Then $\underline{m}$ is a size vector of $(T, \sim)$ in all the cases and $\vartheta(M') \in \mathcal{M}_{\underline{m}}$. On the other hand, the action of ϑ on morphisms is always identity.

Lemma 3.3.1 $M' + H'_{\underline{m}'} = \vartheta(M') + H_{\underline{m}}$.

Proof. Because of $H'_{\underline{m}'} = (H'_{\underline{m}'})_{pq} \otimes \rho + H_{\underline{m}}$, and the definition of $\vartheta(M')$. $\square$

Proposition 3.3.1 ϑ is a functor, which yields an equivalence between category $Mat(\mathcal{K}', \mathcal{M}')$ and a full subcategory of $Mat(\mathcal{K}, \mathcal{M})$.

More precisely, $\varphi' : M' \rightarrow L'$ is a morphism in $Mat(\mathcal{K}', \mathcal{M}')$ if and only if $\vartheta(\varphi') = \varphi' : M \rightarrow L$ is a morphism in $Mat(\mathcal{K}, \mathcal{M})$, where $M = \vartheta(M')$, $L = \vartheta(L')$.

Proof. Suppose that the size vectors of M', L' are $\underline{m}', \underline{l}'$, and the size vectors of M, L are $\underline{m}, \underline{l}$, calculated by Formula (11) of 3.2 respectively. Then Lemma 3.3.1 tells us that

$$M' + H'_{\underline{m}'} = M + H_{\underline{m}}, \quad L' + H'_{\underline{l}'} = L + H_{\underline{l}},$$

therefore

$$(M' + H'_{\underline{m}'})\varphi' = \varphi'(L' + H'_{\underline{l}'}) \text{ iff } (M + H_{\underline{m}})\varphi' = \varphi'(L + H_{\underline{l}}).$$

Thus if $\varphi' \in \mathcal{K}'_{\underline{m}' \times \underline{l}'}$, then $\varphi' \in \mathcal{K}_{\underline{m} \times \underline{l}}$, since $\mathcal{K}'_{\underline{m}' \times \underline{l}'} \subseteq \mathcal{K}_{\underline{m} \times \underline{l}}$. Conversely, if $\varphi' \in \mathcal{K}_{\underline{m} \times \underline{l}}$ is a morphism from M to N , then the first formula ensures that $H'_{\underline{m}'}\varphi' \equiv_{\leq(p,q)} \varphi'H'_{\underline{l}'}$ i.e. $\varphi' \in \mathcal{K}'_{\underline{m}' \times \underline{l}'}$. $\square$

The functor ϑ defined above is called a *reduction functor*.

Let us go back to 2.6 and 3.1. Suppose that a size vector $\underline{n}$ of $(T, \sim)$ is sincere, and a triple $(N, R, \underline{n})$ is given by Formula (6) of 2.6, moreover $(N', R', \underline{n}')$ is obtained from $(N, R, \underline{n})$ by one of the 3 reductions. Then if we change superscript “ r ” to “1”, and continue the reduction for $(N^1, R^1, \underline{n}^1)$, we will obtain again a triple $(N^2, R^2, \underline{n}^2)$. Thus a *reduction sequence* of triples follows by induction:

$$\begin{array}{ccccccc}
 (N, R, \underline{n}), & (N^1, R^1, \underline{n}^1), & \dots, & (N^r, R^r, \underline{n}^r), & \dots, & (N^{s-1}, R^{s-1}, \underline{n}^{s-1}), & (N^s, R^s, \underline{n}^s) \\
 \\
 (*) \quad & N_{pq} & & N_{p_1 q_1}^1 & \dots & N_{p_r q_r}^r & \dots & N_{p_{s-1} q_{s-1}}^{s-1} \\
 & \ddots & & \ddots & & \ddots & & \ddots \\
 & \overline{N}_{pq} & & \overline{N}_{p_1 q_1}^1 & \dots & \overline{N}_{p_r q_r}^r & \dots & \overline{N}_{p_{s-1} q_{s-1}}^{s-1}
 \end{array}$$

where $p_r \in \mathcal{P}_r$, $q_r \in \mathcal{Q}_r$, $\mathcal{P}_r, \mathcal{Q}_r \in T^r / \sim^r$ with $N_{p_r q_r}^r$ being the first non-zero block of N^r , $\overline{N}_{p_r q_r}^r = (H_{\underline{n}^{r+1}}^{r+1})_{p_r q_r}$, $r = 0, 1, \dots, s-1$.

There is also a corresponding *reduction sequence of induced bimodule problems*:

$$(**) \quad (\mathcal{K}, \mathcal{M}, H), (\mathcal{K}^1, \mathcal{M}^1, H^1), \dots, (\mathcal{K}^r, \mathcal{M}^r, H^r), \dots, (\mathcal{K}^{s-1}, \mathcal{M}^{s-1}, H^{s-1}), (\mathcal{K}^s, \mathcal{M}^s, H^s)$$

Furthermore, the sequence $(**)$ yields a *sequence of reduction functors*

$$\vartheta_{r-1, r} : Mat(\mathcal{K}^r, \mathcal{M}^r) \rightarrow Mat(\mathcal{K}^{r-1}, \mathcal{M}^{r-1}), \quad r = 1, 2, \dots, s,$$

where $(\mathcal{K}^0, \mathcal{M}^0, H^0) = (\mathcal{K}, \mathcal{M}, H)$. A composition of reduction functors is also called a *reduction functor*. In particular, let

$$\vartheta_{0,r} = \vartheta_{0,1}\vartheta_{1,2}\cdots\vartheta_{r-1,r} : \text{Mat}(\mathcal{K}^r, \mathcal{M}^r) \rightarrow \text{Mat}(\mathcal{K}, \mathcal{M}),$$

$r = 1, \dots, s$. Then $\vartheta_{0,r}$ are reduction functors.

Corollary 3.3.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem with a reduction sequence $(*)$, and let $\hat{n}^s$ be a size vector of $(T^s, \sim^s)$. Then

$$(\hat{*}) \quad (\hat{N}, \hat{R}, \hat{n}), (\hat{N}^1, \hat{R}^1, \hat{n}^1), \dots, (\hat{N}^r, \hat{R}^r, \hat{n}^r), \dots, (\hat{N}^{s-1}, \hat{R}^{s-1}, \hat{n}^{s-1}), (\hat{N}^s, \hat{R}^s, \hat{n}^s)$$

is also a reduction sequence, provided that $(\hat{N}^{r-1}, \hat{R}^{r-1}, \hat{n}^{r-1})$ is obtained from $(\hat{N}^r, \hat{R}^r, \hat{n}^r)$ by size change given in Proposition 3.2.1 inductively for $r = s, s-1, \dots, 1$. $\square$

3.4 Canonical forms

Our special interests in the sequence $(*)$ lead to a case of $N^s = (0)$, then $\mathcal{M}^s = \{(0)\}$. If that is the case, we shall refer to $(N^s, R^s, \underline{n}^s)$ as $(N^\infty, R^\infty, \underline{n}^\infty)$, $(\mathcal{K}^s, \mathcal{M}^s, H^s)$ as $(\mathcal{K}^\infty, \mathcal{M}^\infty, H^\infty)$, and $\vartheta_{0,s}$ as $\vartheta_{0,\infty}$. The canonical form of an individual matrix of $\text{Mat}(\mathcal{K}, \mathcal{M})$ will emerge from this special case in the present subsection.

Lemma 3.4.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, $M \in \mathcal{M}_{\underline{m}}$, and a triple $(N, R, \underline{m})$ be given by Formula (6) of 2.6. Then there exists a unique choice of $\overline{N}_{pq}$ given by one of the 3 reductions of 3.1, such that $M \simeq \vartheta(M')$ for some $M' \in \text{Mat}(\mathcal{K}', \mathcal{M}')$.

Proof. Quote the matrix equation (7) of 2.6, where we change the first block N_{pq} to the fixed block M_{pq} of M , and the second N_{pq} to $\overline{N}_{pq}$:

$$M_{pq}X_{qq} - X_{pp}\overline{N}_{pq} = \sum_{l=p+1}^t X_{pl}h_{lq} - \sum_{l=1}^{q-1} h_{pl}X_{lq} \quad (12)$$

We will determine $\overline{N}_{pq}$ based on the 3 reductions respectively.

Regularization. If the equation system II of Definition 2.2.1 does not imply equation (8) of 2.6, set $\overline{N}_{pq} = \emptyset$. Define a matrix $S \in \mathcal{K}_{\underline{m} \times \underline{m}}$ having the identities as the diagonal blocks, and satisfying

$$M_{pq} = \sum_{l=p+1}^t S_{pl}h_{lq} - \sum_{l=1}^{q-1} h_{pl}S_{lq}.$$

Suppose now that the equation system II implies (8).

Edge reduction. Set $\overline{N}_{pq} = \begin{pmatrix} 0 & I_r \\ 0 & 0 \end{pmatrix}$ with $r = \text{rank}(M_{pq})$. Define a diagonal block matrix $S \in \mathcal{K}_{\underline{m} \times \underline{m}}$, where $S_{\mathcal{I}} = I_{n_{\mathcal{I}}}$ for any $\mathcal{I} \in T/\sim \setminus \{\mathcal{P}, \mathcal{Q}\}$ and $S_{\mathcal{P}}, S_{\mathcal{Q}}$ satisfy $M_{pq}S_{qq} = S_{pp} \begin{pmatrix} 0 & I_r \\ 0 & 0 \end{pmatrix}$.

Loop reduction. Set $\overline{N}_{pq} = W$, the Weyr matrix similar to M_{pq} . Define a diagonal block matrix $S \in \mathcal{K}_{\underline{m} \times \underline{m}}$ with $S_{\mathcal{I}} = I_{n_{\mathcal{I}}}$ for any $\mathcal{I} \in T/\sim \setminus \{\mathcal{P}\}$ and $S_{\mathcal{P}}$ satisfying $M_{pq}S_{qq} = S_{pp}W$.

In all the 3 cases, let

$$M' = S^{-1}(M + H_{\underline{m}})S - (\overline{N}_{pq} \otimes \rho + H_{\underline{m}}). \quad (13)$$

Then $M' \in \mathcal{M}_{\underline{m}'}$. Thus

$$(M + H_{\underline{m}})S = S(M' + H'_{\underline{m}'}),$$

i.e. $M \simeq \vartheta(M')$ in $Mat(\mathcal{K}, \mathcal{M})$. $\square$

Lemma 3.4.2 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, $M, L \in Mat(\mathcal{K}, \mathcal{M})$. If $M \simeq L$, then there exists an induced bimodule problem $(\mathcal{K}', \mathcal{M}', H')$ obtained by one of 3 reductions of 3.1, such that $M \simeq \vartheta(M')$, $L \simeq \vartheta(L')$ for some $M', L' \in Mat(\mathcal{K}', \mathcal{M}')$ and $M' \simeq L'$.

Proof. Since $M \simeq L$, M and N have the same size vector $\underline{m}$ of $(T, \sim)$ by Corollary 2.2.1. Suppose $S \in \mathcal{K}_{\underline{m} \times \underline{m}}$ is invertible such that $(M + H_{\underline{m}})S = S(L + H_{\underline{m}})$ with the (p, q) -block

$$M_{pq}S_{qq} - S_{pp}L_{pq} = \sum_{l=p+1}^t S_{pl}h_{lq} - \sum_{l=1}^{q-1} h_{pl}S_{lq}.$$

If the equation system II of 2.2 does not imply (8) of 2.6, set $\overline{N}_{pq} = \emptyset$.

Otherwise, the equality becomes $M_{pq}S_{qq} = S_{pp}L_{pq}$. Set $\overline{N}_{pq} = \begin{pmatrix} 0 & I_r \\ 0 & 0 \end{pmatrix}$ whenever $p \approx q$, where $r = rank(M_{pq}) = rank(L_{pq})$; or $\overline{N}_{pq} = W$ whenever $p \sim q$, where W is a Weyr matrix similar to M_{pq} and L_{pq} . Therefore $M \simeq \vartheta(M')$, $L \simeq \vartheta(L')$ for some $M', L' \in \mathcal{M}_{\underline{m}'}$ by Lemma 3.4.1, i.e.

$$(M + H_{\underline{m}})U = U(M' + H'_{\underline{m}'}) \quad \text{and} \quad (L + H_{\underline{m}})V = V(L' + H'_{\underline{m}'})$$

for some invertible $U, V \in \mathcal{K}_{\underline{m} \times \underline{m}}$. Finally

$$(M' + H'_{\underline{m}'})U^{-1}SV = U^{-1}SV(L' + H'_{\underline{m}'})$$

with $U^{-1}SV \in \mathcal{K}'_{\underline{m}' \times \underline{m}'}$, i.e. $M' \simeq L'$ in $Mat(\mathcal{K}', \mathcal{M}')$. $\square$

Theorem 3.4.1[S]. Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem, $M \in \mathcal{M}_{\underline{m}}$. Then there exists a unique reduction sequence $(*)$ starting from $(N, R, \underline{m})$, ending at $(N^{\infty}, R^{\infty}, \underline{m}^{\infty})$, such that

$$M \simeq \vartheta_{0, \infty}(M^{\infty})$$

for some $M^\infty \in \text{Mat}(\mathcal{K}^\infty, \mathcal{M}^\infty)$ under both the fixed orders of matrices and of the base field (see Definition 2.3.1 and 2.5).

Proof. Suppose we have already had a unique sequence $(*)$ up to $(N^r, R^r, \underline{m}^r)$, and some $M^r \in \mathcal{M}_{\underline{m}^r}^r$, such that $\vartheta_{0,r}(M^r) \simeq M$. Lemma 3.4.1 shows a unique induced triple $(N^{r+1}, R^{r+1}, \underline{m}^{r+1})$ with $\vartheta_{r,r+1}(M^{r+1}) \simeq M^r$ for some $M^{r+1} \in \mathcal{M}_{\underline{m}^{r+1}}^{r+1}$. On the other hand, Lemma 3.4.2 ensures that any choice of M^r in an isomorphism class of $\text{Mat}(\mathcal{K}^r, \mathcal{M}^r)$ does not influence the determination of $\overline{N}_{p_r q_r}$. Therefore the induced triple $(N^{r+1}, R^{r+1}, \underline{m}^{r+1})$ is uniquely determined. Because of the finiteness of the size of N , we finally reach to $N^s = (0)$ for some positive integer s . $\square$

Remarks. From now on, we assume that the original bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ satisfies the condition $H = 0$.

(1) If that is the case, $\vartheta_{0,\infty}(M^\infty) \simeq M$ implies that $(M + 0)S = S(0 + H_{\underline{m}^\infty}^\infty)$ for some invertible $S \in \mathcal{K}_{\underline{m} \times \underline{m}}$, or equivalently,

$$H_{\underline{m}^\infty}^\infty = S^{-1}MS.$$

The matrix $H_{\underline{m}^\infty}^\infty$ is called the *canonical form* of M , which can be regarded as a generalized Jordan form. It is clear that $\mathcal{K}_{\underline{m}^\infty \times \underline{m}^\infty}^\infty$ is the endomorphism ring of $H_{\underline{m}^\infty}^\infty$.

Without $H = 0$ the trouble is that there is no guarantee such that $H_{\underline{m}^\infty}^\infty$ is similar to M , which is not convenient.

(2) The restriction $H = 0$ is natural, since the most bimodule problems considered in practice, including our main subject $P_1(\Lambda)$, have such a property.

Corollary 3.4.1 Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem, $M, L \in \text{Mat}(\mathcal{K}, \mathcal{M})$. Then $M \simeq L$ if and only if M and L have the same canonical forms. $\square$

3.5 Krull-Schmidt property

Following Ringel [R2], an k -additive category with the finite dimensional objects is called a *Krull-Schmidt category*, provided it has finite direct sums and split idempotents.

Proposition 3.5.1 [CB2]. Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, then $\text{Mat}(\mathcal{K}, \mathcal{M})$ is a Krull-Schmidt category.

Proof. Clearly, $\text{Mat}(\mathcal{K}, \mathcal{M})$ has finite direct sums.

Now suppose $\varphi : M \rightarrow M$ is an idempotent with $M \in \mathcal{M}_{\underline{m}}$ and $\varphi \in \mathcal{K}_{\underline{m} \times \underline{m}}$. Since idempotents split in the finite dimensional algebra $\mathcal{K}$, there exists a size vector $\underline{l}$ and some matrices $\pi \in \mathcal{K}_{\underline{m} \times \underline{l}}$, $\iota \in \mathcal{K}_{\underline{l} \times \underline{m}}$ such that $\pi\iota = \varphi$ and $\iota\pi = id_{\mathcal{K}_{\underline{l} \times \underline{l}}}$. Let $L = \iota M \pi - \iota d(\pi)$, we will show that $\pi : M \rightarrow L$ and $\iota : L \rightarrow M$ are morphisms of $\text{Mat}(\mathcal{K}, \mathcal{M})$.

In fact, since $\pi = \varphi\pi$, $d(\pi) = d(\varphi)\pi + \varphi d(\pi)$, $M\pi - \pi L = M\pi - \varphi M\pi + \varphi d(\pi) = M\varphi\pi - \varphi M\pi + d(\pi) - d(\varphi)\pi = (M\varphi - \varphi M - d(\varphi))\pi + d(\pi) = d(\pi)$. Similarly, since $id_L = \iota\pi$, $\iota = \iota\varphi$, we have $d(\iota)\pi + \iota d(\pi) = 0$ and $d(\iota) = d(\iota)\varphi + \iota d(\varphi)$. Thus $L\iota - \iota M = \iota M\varphi - \iota d(\pi)\iota - \iota M = \iota M\varphi + d(\iota)\varphi - \iota M = \iota M\varphi - \iota\varphi M + d(\iota) - \iota d(\varphi) = \iota(M\varphi - \varphi M - d(\varphi)) + d(\iota) = d(\iota)$. $\square$

Lemma 3.5.1. Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, and $\underline{n}$ be a size vector of $(T, \sim)$. If $\varphi \in \mathcal{K}_{\underline{n} \times \underline{n}}$ is an idempotent, there exists some transformation matrix $\chi \in \mathcal{K}_{\underline{n} \times \underline{n}}$, such that $\overline{\varphi} = \chi^{-1}\varphi\chi$ is a diagonal idempotent with

$$(\overline{\varphi})_{\mathcal{L}} = \begin{pmatrix} I_{d_{\mathcal{L}}} & 0 \\ 0 & 0 \end{pmatrix}_{n_{\mathcal{L}} \times n_{\mathcal{L}}},$$

for some $0 \leq d_{\mathcal{L}} \leq n_{\mathcal{L}}$, and for each $\mathcal{L} \in T/\sim$.

Proof. (Given by Hu Yongjian) Suppose that $\varphi = (S_{ij})_{t \times t}$ is an upper triangular partitioned matrix with $S_{ij} (i < j)$ being $n_i \times n_j$ matrices satisfying the equation system II of Definition 2.2.1.

(1) It is clear that $\forall \mathcal{L} \in T/\sim$, $S_{\mathcal{L}}^2 = S_{\mathcal{L}}$. Thus there exists some invertible matrix $U_{\mathcal{L}}$, such that $U_{\mathcal{L}}^{-1}S_{\mathcal{L}}U_{\mathcal{L}} = \begin{pmatrix} I_{d_{\mathcal{L}}} & 0 \\ 0 & 0 \end{pmatrix}_{n_{\mathcal{L}} \times n_{\mathcal{L}}}$, since our base field k is algebraically closed. Let $\chi_0 = \text{diag}(U_{11}, U_{22}, \dots, U_{tt})$, and $\varphi_1 = \chi_0^{-1}\varphi\chi_0$

(2) Denote φ_1 by A , then

$$A = \begin{pmatrix} \boxed{\begin{matrix} I_{d_1} & 0 \\ 0 & 0 \end{matrix}} & \boxed{S_{12}^1} & \boxed{S_{12}^2} & \cdots & \boxed{S_{1l}^1} & \boxed{S_{1l}^2} & \cdots & \boxed{S_{1t}^1} & \boxed{S_{1t}^2} \\ & \boxed{\begin{matrix} I_{d_2} & 0 \\ 0 & 0 \end{matrix}} & \cdots & & \boxed{S_{2l}^1} & \boxed{S_{2l}^2} & \cdots & \boxed{S_{2t}^1} & \boxed{S_{2t}^2} \\ & & \ddots & & \vdots & \vdots & \cdots & \vdots & \vdots \\ & & & \boxed{\begin{matrix} I_{d_t} & 0 \\ 0 & 0 \end{matrix}} & \cdots & & \boxed{S_{lt}^1} & \boxed{S_{lt}^2} \\ & & & & \ddots & & \vdots & \vdots \\ & & & & & \boxed{\begin{matrix} I_{d_t} & 0 \\ 0 & 0 \end{matrix}} \end{pmatrix},$$

Let A_l be the partitioned submatrix of A consisting of the intersections of 1- l block-rows

and block columns. Let

$$\xi_l = \begin{pmatrix} S_{1l}^1 \\ S_{2l}^1 \\ \vdots \\ S_{l-1,l}^1 \end{pmatrix}, \quad \eta_l = \begin{pmatrix} S_{1l}^2 \\ S_{2l}^2 \\ \vdots \\ S_{l-1,l}^2 \end{pmatrix}$$

Then

$$A_l = \left(\begin{array}{c|cc} A_{l-1} & \xi_l & \eta_l \\ \hline & I_{d_l} & 0 \\ & 0 & 0 \end{array} \right).$$

(3) We claim that

- (i) $A_l^2 = A_l$ is an idempotent matrix.
- (ii) $A_{l-1}\xi_l = 0$.
- (iii) $(I - A_{l-1})\eta_l = 0$.

In fact $A_t^2 = A^2 = A = A_t$. If $A_l^2 = A_l$, then

$$\left(\begin{array}{c|cc} A_{l-1}^2 & A_{l-1}\xi_l + \xi_l & A_{l-1}\eta_l \\ \hline & I_d & 0 \\ & 0 & 0 \end{array} \right) = \left(\begin{array}{c|cc} A_{l-1} & \xi_l & \eta_l \\ \hline & I_d & 0 \\ & 0 & 0 \end{array} \right).$$

Our claim follows by induction on $l = t, t-1, \dots, 1$

(4) We define a transformation matrix P_l inductively on $l = 1, 2, \dots, t$. Let $P_1 = I_{n_1}$, than $P_1^{-1}A_1P_1 = \begin{pmatrix} I_{d_1} & 0 \\ 0 & 0 \end{pmatrix}$. We assume that there exists an invertible matrix P_{l-1} , such

that $P_{l-1}^{-1}A_{l-1}P_{l-1} = \text{diag} \left(\begin{pmatrix} I_{d_1} & 0 \\ 0 & 0 \end{pmatrix}, \dots, \begin{pmatrix} I_{d_{l-1}} & 0 \\ 0 & 0 \end{pmatrix} \right)$. Let $P_l = \left(\begin{array}{c|cc} P_{l-1} & \xi_l & -\eta_l \\ \hline 0 & I_{d_l} & 0 \\ & 0 & I_{n_l-d_l} \end{array} \right)$,

then

$$\begin{aligned} P_l^{-1}A_lP_l &= \left(\begin{array}{c|cc} P_{l-1}^{-1} & -P_{l-1}^{-1}\xi_l & P_{l-1}^{-1}\eta_l \\ \hline & I_{d_l} & 0 \\ & 0 & I_{n_l-d_l} \end{array} \right) \left(\begin{array}{c|cc} A_{l-1} & \xi_l & \eta_l \\ \hline & I_{d_l} & 0 \\ & 0 & 0 \end{array} \right) \left(\begin{array}{c|cc} P_{l-1} & \xi_l & -\eta_l \\ \hline 0 & I_{d_l} & 0 \\ & 0 & I_{n_l-d_l} \end{array} \right) \\ &= \left(\begin{array}{c|cc} P_{l-1}^{-1}A_{l-1} & 0 & P_{l-1}^{-1}\eta_l \\ \hline & I_{d_l} & 0 \\ & 0 & 0 \end{array} \right) \left(\begin{array}{c|cc} P_{l-1} & \xi_l & -\eta_l \\ \hline & I_{d_l} & 0 \\ & 0 & I_{n_l-d_l} \end{array} \right) \\ &= \left(\begin{array}{c|cc} P_{l-1}^{-1}A_{l-1}P_{l-1} & 0 & 0 \\ \hline & I_{d_l} & 0 \\ & 0 & I_{n_l-d_l} \end{array} \right) \end{aligned}$$

by (ii), (iii) of (3). Thus we may take

$$P = \begin{pmatrix} \boxed{I_{n_1}} & \boxed{\xi_1} & \boxed{-\eta_1} & & & & \\ & \boxed{I_{n_2}} & \boxed{\xi_2} & \boxed{-\eta_2} & & & \\ & & \boxed{I_{n_3}} & & & \boxed{\xi_t} & \boxed{-\eta_t} \\ & & & \ddots & & & \\ & & & & \boxed{I_{n_t}} & & \end{pmatrix}.$$

(5) Denote P by χ_1 , we prove that $\chi_1 \in \mathcal{K}_{\underline{n} \times \underline{n}}$. In fact, since $A \in \mathcal{K}_{\underline{n} \times \underline{n}}$, we have

$$\sum_{\mathcal{I} \ni i < j \in \mathcal{J}} c_{ij}^l S_{ij} = 0$$

for $1 \leq l \leq q_{\mathcal{I}\mathcal{J}}$ and each $(\mathcal{I}, \mathcal{J}) \in (T/\sim) \times (T/\sim)$, where $S_{ij} = (S_{ij}^1 \mid S_{ij}^2)$. Therefore

$$\sum_{\mathcal{I} \ni i < j \in \mathcal{J}} c_{ij}^l S_{ij}^1 = 0 \text{ and } \sum_{\mathcal{I} \ni i < j \in \mathcal{J}} c_{ij}^l (-S_{ij}^2) = 0. \text{ Consequently,}$$

$$\sum_{\mathcal{I} \ni i < j \in \mathcal{J}} c_{ij}^l P_{ij} = 0,$$

where $P_{ij} = (S_{ij}^1 \mid -S_{ij}^2)$ for $i < j$. Thus $P \in \mathcal{K}_{\underline{n} \times \underline{n}}$.

Let $\chi = \chi_0 \chi_1 \in \mathcal{K}_{\underline{n} \times \underline{n}}$, then $\chi^{-1} \varphi \chi = \text{diag} \left(\begin{pmatrix} I_{d_1} & 0 \\ 0 & 0 \end{pmatrix}, \dots, \begin{pmatrix} I_{d_l} & 0 \\ 0 & 0 \end{pmatrix} \right)$ for some $0 \leq d_l \leq n_{\mathcal{L}}$ and each $\mathcal{L} \in T/\sim$. The proof is completed. $\square$

Lemma 3.5.1 suggests an alternative definition of direct sums in $\text{Mat}(\mathcal{K}, \mathcal{M})$, which is simple and concrete. We first define the direct sum of general matrices.

Definition 3.5.1 Let $(T, \sim)$ be a set of integers with a relation given in I of Definition 2.2.1, and let $C = (C_{ij})_{t \times t}$, $D = (D_{ij})_{t \times t}$ be two matrices partitioned by $(T, \sim)$. Then a matrix

$$\begin{pmatrix} C_{ij} & 0 \\ 0 & D_{ij} \end{pmatrix}_{t \times t}$$

is said to be a *direct sum* of C and D , and is denoted by $C \oplus_T D$, or $C \oplus D$ for simplicity.

Lemma 3.5.2 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem. If $\underline{m}_1, \underline{m}_2$ are size vectors of $(T, \sim)$ and $\underline{m} = \underline{m}_1 + \underline{m}_2$, then $H_{\underline{m}} = H_{\underline{m}_1} \oplus_T H_{\underline{m}_2}$.

Proof. Let $H = (h_{ij})_{t \times t}$. Given any pair (i, j) with $1 \leq i, j \leq t$, if $i \approx j$, then $h_{ij} = 0$, and $(H_{\underline{m}_l})_{ij} = 0$; if $i \sim j$ then $(H_{\underline{m}_l})_{ij} = h_{ij} I_{(m_l)_i}$ for $l = 0, 1, 2$, where $\underline{m}_0 = \underline{m}$. Thus

$$(H_{\underline{m}})_{ij} = (H_{\underline{m}_1})_{ij} \oplus (H_{\underline{m}_2})_{ij},$$

since $m_i = (m_1)_i + (m_2)_i$, $m_j = (m_1)_j + (m_2)_j$. The assertion follows block-wise. $\square$

Lemma 3.5.3 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem.

(1) If $M_1 \in \mathcal{M}_{\underline{m}_1}$, $M_2 \in \mathcal{M}_{\underline{m}_2}$ and $M = M_1 \oplus_T M_2$, then

$$M + H_{\underline{m}} = (M_1 + H_{\underline{m}_1}) \oplus_T (M_2 + H_{\underline{m}_2}).$$

And $M = M_1 \oplus M_2$ in the sense of Proposition 3.5.1.

(2) Conversely, $\forall M \in \mathcal{M}_{\underline{m}}$, if $\varphi : M \rightarrow M$ is an idempotent, then there exists some $M_0 \simeq M$, such that $M_0 = M_1 \oplus_T M_2$ in the sense of Definition 3.5.1.

Proof. (1) The direct sum over T is give by lemma 3.5.2. And the idempotent can be obtained easily.

(2) $\forall M \in \mathcal{M}_{\underline{m}}$ and any idempotent $\varphi : M \rightarrow M$, there exists an invertible matrix $\chi \in \mathcal{K}_{\underline{m} \times \underline{m}}$, such that $\overline{\varphi} = \chi^{-1} \varphi \chi$ is a diagonal idempotent given by Lemma 3.5.1. Define

$$M_0 = \chi^{-1} (M + H_{\underline{m}}) \chi - H_{\underline{m}},$$

then $M \simeq M_0$, and $\overline{\varphi} : M_0 \rightarrow M_0$ is a morphism. Thus $(M_0 + H_{\underline{m}}) \overline{\varphi} = \overline{\varphi} (M_0 + H_{\underline{m}})$, $\forall i, j \in T$, we have

$$((M_0)_{ij} + (H_{\underline{m}})_{ij}) \begin{pmatrix} I_{d_j} & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} I_{d_i} & 0 \\ 0 & 0 \end{pmatrix} ((M_0)_{ij} + (H_{\underline{m}})_{ij}),$$

i.e.

$$(M_0)_{ij} = \begin{pmatrix} M_{ij}^1 & 0 \\ 0 & M_{ij}^2 \end{pmatrix},$$

such that the size of $(M_1)_{ij}$ is $d_i \times d_j$, and that of $(M_2)_{ij}$ is $(m_i - d_i) \times (m_j - d_j)$. $\square$

$M \in \text{Mat}(\mathcal{K}, \mathcal{M})$ is said to be *indecomposable* if $M \simeq M_1 \oplus_T M_2$ implies that the size of M_1 or M_2 equals 0.

Theorem 3.5.1. Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem with a reduction sequence $(*)$ given by Theorem 3.4.1 for some $M \in \text{Mat}(\mathcal{K}, \mathcal{M})$.

(1) Define a size vector $\underline{m}(\mathcal{I})$ of $(T^\infty, \sim^\infty)$ for any $\mathcal{I} \in T^\infty / \sim^\infty$, such that $m(\mathcal{I})_{\mathcal{I}} = 1$ and $m(\mathcal{I})_{\mathcal{J}} = 0$, $\forall \mathcal{J} \neq \mathcal{I}$. Then the zero matrix $O(\mathcal{I}) \in \text{Mat}(\mathcal{K}^\infty, \mathcal{M}^\infty)$ of size $\underline{m}(\mathcal{I})$ is indecomposable. And

$$\{O(\mathcal{I}) \mid \forall \mathcal{I} \in T^\infty / \sim^\infty\}$$

is a complete set of isomorphism classes of indecomposables of $Mat(\mathcal{K}^\infty, \mathcal{M}^\infty)$. Moreover,

$$M^\infty = \oplus_{T^\infty} O(\mathcal{I})^{m_{\mathcal{I}}^\infty}$$

in $Mat(\mathcal{K}^\infty, \mathcal{M}^\infty)$.

(2) Let $\vartheta : Mat(\mathcal{K}^\infty, \mathcal{M}^\infty) \rightarrow Mat(\mathcal{K}, \mathcal{M})$ be the reduction functor. Then

$$\vartheta(O(\mathcal{I})) = H_{\underline{m}(\mathcal{I})}^\infty \text{ and denote it by } H^\infty(\mathcal{I}) \text{ for simplicity.}$$

Thus $H^\infty(\mathcal{I})$ is the canonical form of a direct summand of M for any $\mathcal{I} \in T^\infty / \sim^\infty$. And $M \simeq \oplus_{T^\infty} (H^\infty(\mathcal{I}))^{m_{\mathcal{I}}^\infty}$.

(3) If $M \simeq \oplus M_l^{e_l}$ in the sense of Proposition 3.5.1, such that M_l are pairwise non-isomorphic, then there exists some $\mathcal{I}_l \in T^\infty / \sim^\infty$, such that $M_l \simeq H^\infty(\mathcal{I}_l)$ and $e_l = m_{\mathcal{I}_l}^\infty$.

Proof. (1) Since

$$O(\mathcal{I}) \in \mathcal{M}_{\underline{m}(\mathcal{I})}^\infty, \text{ and } \mathcal{K}_{\underline{m}(\mathcal{I}) \times \underline{m}(\mathcal{I})}^\infty,$$

the endomorphism ring of $O(\mathcal{I})$, is local, $O(\mathcal{I})$ is indecomposable. The direct sum comes from $\underline{m}^\infty = \sum_{\mathcal{I}} \underline{m}(\mathcal{I})^{m_{\mathcal{I}}^\infty}$

(2) $\forall \mathcal{I} \in T^\infty / \sim^\infty$, denote $\underline{m}(\mathcal{I})$ by $\hat{n}^\infty$. Then corollary 3.3.1 gives a reduction sequence with a minimal end term. Thus $H^\infty(\mathcal{I})$ is a canonical form of the objects in an isomorphism classes. Let $\iota : H^\infty(\mathcal{I}) \rightarrow H_{\underline{m}^\infty}^\infty$ and $\pi : H_{\underline{m}^\infty}^\infty \rightarrow H^\infty(\mathcal{I})$ be the natural morphisms given by the direct summand respectively, and $M\varphi = \varphi H_{\underline{m}^\infty}^\infty$ be given by Remark (1) of 3.4 for some morphism $\varphi \in \mathcal{K}_{\underline{m} \times \underline{m}}$. Then the morphisms $\iota\varphi : H^\infty(\mathcal{I}) \rightarrow M$ and $\varphi^{-1}\pi : M \rightarrow H^\infty(\mathcal{I})$ tell us that $H^\infty(\mathcal{I})$ is isomorphic to a direct summand of M . The direct sum is given by lemma 3.5.3 (1).

(3) Since $\oplus M_l^{e_l} \simeq \oplus (H^\infty(\mathcal{I}))^{m_{\mathcal{I}}^\infty}$, the assertion follows from Krull-Schmidt theorem.

□

For example, see the last part of 2.2, we have $M \simeq M_1 \oplus_T M_2$, if we exchange the first and second rows and columns inside the i_{10} -row and column blocks simultaneously. Where

$$M_1 = \begin{array}{|c|c|c|c|c|c|} \hline & & & & & 1 \\ \hline & & & & \lambda & \\ \hline & & & & 1 & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array} \begin{array}{l} i_1 \\ i_2 \\ i_3 \\ i_5 \\ i_{10} \\ i_{12} \end{array}, \quad M_2 = \begin{array}{|c|c|c|c|c|c|} \hline & & & & & \\ \hline & & & & & \\ \hline & & & & 1 & \\ \hline & & & & 0 & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array} \begin{array}{l} i_1 \\ i_2 \\ i_3 \\ i_5 \\ i_{10} \end{array},$$

where the endomorphism rings are

$$\mathcal{K}_{\underline{m}_1^\infty \times \underline{m}_1^\infty}^\infty = \{\text{diag}(s_1, s_1, s_1, s_1, s_1, s_1) \mid \forall s_1 \in k\},$$

and

$$\mathcal{K}_{\underline{m}_2^\infty \times \underline{m}_2^\infty}^\infty = \{\text{diag}(s_2, s_2, s_2, s_2, s_2, s_2) \mid \forall s_2 \in k\}$$

respectively.

3.6 Corresponding bocses

The present subsection is devoted to describing the dual structure of a bimodule problem $(\mathcal{K}, \mathcal{M}, H)$, i.e. the corresponding boc $\mathfrak{A}$ of $(\mathcal{K}, \mathcal{M}, H)$. The notion of boc has been studied intensively in Kiev school since 70's of last century, and it has been used successfully for the proof of Drozd's theorem. We illustrate here an explicit relation between $(\mathcal{K}, \mathcal{M}, H)$ and $\mathfrak{A}$ in order to simplify the calculation in the proof of our main theorem.

Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem defined in 2.2.1 with a triangular basis (A, B) . Then the algebra $(\mathcal{K}, \cdot, 1)$ yields a coalgebra structure $(\Omega, \mu, \varepsilon)$ with the dual basis

$$\{E_{\mathcal{I}}^* \mid \forall \mathcal{I} \in T / \sim\} \cup B^*.$$

Namely if the multiplication of algebra $\mathcal{K}$ is given by

$$\zeta_{\mathcal{I}\mathcal{L}}^u \zeta_{\mathcal{L}\mathcal{J}}^v = \sum_{\omega} c_{\binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}}}^{\binom{\omega}{\mathcal{I}\mathcal{J}}} \zeta_{\mathcal{I}\mathcal{J}}^{\omega} \quad (14)$$

with the structure constants $c \in k$, and

$$\zeta_{\mathcal{I}\mathcal{J}}^w E_{\mathcal{J}} = \zeta_{\mathcal{I}\mathcal{J}}^w, \quad E_{\mathcal{I}} \zeta_{\mathcal{I}\mathcal{J}}^w = \zeta_{\mathcal{I}\mathcal{J}}^w,$$

then

$$\mu(\zeta_{\mathcal{I}\mathcal{J}}^w)^* = (\zeta_{\mathcal{I}\mathcal{L}}^w)^* \otimes (E_{\mathcal{J}})^* + (E_{\mathcal{I}})^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^w)^* + \delta_2(\zeta_{\mathcal{I}\mathcal{J}}^w)^*, \quad (15)$$

where

$$\delta_2(\zeta_{\mathcal{I}\mathcal{J}}^w)^* = \sum_{\mathcal{L}, u, v} c_{\binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}}}^{\binom{\omega}{\mathcal{I}\mathcal{J}}} (\zeta_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^v)^*, \quad (16)$$

and

$$\mu(E_{\mathcal{I}})^* = (E_{\mathcal{I}})^* \otimes (E_{\mathcal{I}})^*. \quad (17)$$

Therefor we obtain a linear map $\mu : \Omega \rightarrow \Omega \otimes \Omega$. On the other hand, let $\Gamma' = k \times k \times \cdots \times k$ with s copies of k be a trivial algebra, where the equivalent classes $\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_s$ may

be regarded as the vertices in the ordinary quiver of Γ' . Then we define a linear map $\varepsilon : \Omega \rightarrow \Gamma'$ given by $\varepsilon(E_I)^* = 1_I$ and $\varepsilon(\zeta_{IJ}^w)^* = 0$. Thus we have the counit ε and comultiplication μ of Ω , which satisfy the laws of coalgebra $(\mu \otimes id)\mu = (id \times \mu)\mu$, and $(\varepsilon \otimes id)\mu = id$, $(id \otimes \varepsilon)\mu = id$, where id stands for the identity morphism on Ω .

On the other hand $\mathcal{K}$ - $\mathcal{K}$ bimodule $\mathcal{M}$ yields a left and right comodule structure Γ_m , which has the dual basis $A^* = \{(\rho_{IJ}^w)^*\}$, and the comodule actions are given according to Proposition 2.4.1:

$$\begin{aligned}\gamma_R(\rho_{IJ}^w)^* &= \sum_{\mathcal{L}, u, v} \iota^c \binom{w}{\mathcal{I}\mathcal{J}} \binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}} (\zeta_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\rho_{\mathcal{L}\mathcal{J}}^v)^*, \\ \gamma_L(\rho_{IJ}^w)^* &= \sum_{\mathcal{L}, u, v} r^c \binom{w}{\mathcal{I}\mathcal{J}} \binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}} (\rho_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^v)^*,\end{aligned}$$

then $\gamma_L : \Gamma_m \rightarrow \Omega \otimes \Gamma_m$ satisfies the left comodule law $(\mu \otimes id_{\Gamma_m})\gamma = (id_{\Omega} \otimes \gamma)\gamma$ and $\gamma_R : \Gamma_m \rightarrow \Gamma_m \otimes \Omega$ satisfies the right comodule law $(id_{\Gamma} \otimes \mu)\gamma = (\gamma \otimes id_{\Omega})\gamma$.

Next we reconstruct Γ_m to an algebra Γ which is freely generated by $A^* = \{(\rho_{IJ}^w)^*\}$ over Γ' . Then Ω can be viewed as a Γ - Γ -bimodule as follows. Write $\Omega = \Omega_0 \oplus \bar{\Omega}$ as a direct sum of k -vector spaces, then $\Omega_0 \cong \Gamma$, $\bar{\Omega}$ is freely generated by $B^* = \{(\zeta_{IJ}^w)^*\}$ over Γ . Furthermore suppose that

$$d(\zeta_{IJ}^u) = \zeta_{IJ}^u H - H \zeta_{IJ}^u = \sum_w c_{IJ}^{uw} \rho_{IJ}^w, \quad (18)$$

then the derivation $d : \mathcal{K} \rightarrow \mathcal{M}$ has a duality $Dd : D\mathcal{M} \rightarrow D\mathcal{K}$ given by

$$(Dd)(\rho_{IJ}^w)^* = \sum_u c_{IJ}^{uw} (\zeta_{IJ}^u)^*. \quad (19)$$

which is in fact from $D\mathcal{M}$ to $D(\text{rad}\mathcal{K})$, because of $d(E_I) = 0$ by Corollary 2.2.1. According to formula (18), and the left and right module actions given in Proposition 2.4.1, we define a map δ_1 on A^* given by

$$\delta_1(\rho_{IJ}^w)^* = (Dd)(\rho_{IJ}^w)^* + \gamma_L(\rho_{IJ}^w)^* - \gamma_R(\rho_{IJ}^w)^*,$$

more precisely

$$\delta_1(\rho_{IJ}^w)^* = \sum_u c_{IJ}^{uw} (\zeta_{IJ}^u)^* + \sum_{\mathcal{L}, u, v} \left(\iota^c \binom{w}{\mathcal{I}\mathcal{J}} \binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}} (\zeta_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\rho_{\mathcal{L}\mathcal{J}}^v)^* - r^c \binom{w}{\mathcal{I}\mathcal{J}} \binom{u}{\mathcal{I}\mathcal{L}} \binom{v}{\mathcal{L}\mathcal{J}} (\rho_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^v)^* \right) \quad (20)$$

Then Ω satisfies the relations:

$$(\rho_{IJ}^w)^* \otimes (E_J)^* - (E_I)^* \otimes (\rho_{IJ}^w)^* = \delta_1(\rho_{IJ}^w)^*. \quad (21)$$

The reason will be given in Lemma 3.6.1. We denote the structure defined above by $\mathfrak{A}$, and write $\mathfrak{A} = (\Gamma, \Omega)$, which is called a *bocs*, see [D, CB1]. Note that both μ and ε can be viewed as Γ - Γ -bimodule morphisms.

Now we show that the maps $\delta_1 : \Gamma \rightarrow \Omega$ and $\delta_2 : \bar{\Omega} \rightarrow \bar{\Omega} \otimes \bar{\Omega}$ extended linearly are determined just by multiplications of matrices.

If we write x_{ij} , z_{ij} for the (i, j) -entries of R_0 , N_0 which satisfy 2.2.1.II and III respectively, then

$$R_0 = \sum_i E_{ii}^* E_{ii} + \sum_{i,j} \left(\sum_w (\zeta_{\mathcal{I}\mathcal{J}}^w)^* b_{\mathcal{I}\mathcal{J}}^w(i, j) \right) E_{ij} = \sum_{i,j} x_{ij} E_{ij}, \quad (22)$$

$$N_0 = \sum_{i,j} \left(\sum_w (\rho_{\mathcal{I}\mathcal{J}}^w)^* a_{\mathcal{I}\mathcal{J}}^w(i, j) \right) E_{ij} = \sum_{i,j} z_{ij} E_{ij}. \quad (23)$$

Recall Formulae (1) and (2) of 2.3, the entries $z_{p_{\mathcal{I}\mathcal{J}}^w q_{\mathcal{I}\mathcal{J}}^w} = (\rho_{\mathcal{I}\mathcal{J}}^w)^*$ of N_0 are said to be *free*, and $z_{ij} = \sum_w a_{\mathcal{I}\mathcal{J}}^w(i, j) (\rho_{\mathcal{I}\mathcal{J}}^w)^*$ for $(i, j) \neq (p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w)$, $\forall w, \forall (\mathcal{I}, \mathcal{J}) \in T / \sim \times T / \sim$, are said to be *dependent*. Similarly, the entries $x_{\bar{p}_{\mathcal{I}\mathcal{J}}^w \bar{q}_{\mathcal{I}\mathcal{J}}^w} = (\zeta_{\mathcal{I}\mathcal{J}}^w)^*$ of R_0 are said to be *free* and $x_{ij} = \sum_w b_{\mathcal{I}\mathcal{J}}^w(i, j) (\zeta_{\mathcal{I}\mathcal{J}}^w)^*$ for $(i, j) \neq (\bar{p}_{\mathcal{I}\mathcal{J}}^w, \bar{q}_{\mathcal{I}\mathcal{J}}^w)$, $\forall w, \forall (\mathcal{I}, \mathcal{J}) \in T / \sim \times T / \sim$, are said to be *dependent*. Now consider the matrix equation $(N_0 + H)R_0 = R_0(N_0 + H)$, the $(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w)$ -entry, which we denote by p, q for simplicity, gives an equation:

$$z_{pq} x_{qq} - x_{pp} z_{pq} = \left(\sum_{p < l} x_{pl} h_{lq} - \sum_{l < q} h_{pl} x_{lq} \right) + \left(\sum_{p < l} x_{pl} z_{lq} - \sum_{l < q} z_{pl} x_{lq} \right) \quad (24)$$

By Formulae (1) and (2) of 2.3 the left and right module actions given in Proposition 2.4.1 are equivalent respectively to the formulae

$$\sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} \left(\sum_{l \in \mathcal{L}} b_{\mathcal{I}\mathcal{L}}^u(i, l) a_{\mathcal{L}\mathcal{J}}^v(l, j) \right) E_{ij} = \sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} \left(\sum_w {}^l c_{\left(\begin{smallmatrix} \mathcal{I} \\ \mathcal{I}\mathcal{L} \end{smallmatrix}\right)}^{\left(\begin{smallmatrix} w \\ \mathcal{I}\mathcal{J} \end{smallmatrix}\right)} a_{\mathcal{I}\mathcal{J}}^w(i, j) \right) E_{ij}, \quad (25)$$

and

$$\sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} \left(\sum_{l \in \mathcal{L}} a_{\mathcal{I}\mathcal{L}}^u(i, l) b_{\mathcal{L}\mathcal{J}}^v(l, j) \right) E_{ij} = \sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} \left(\sum_w {}^r c_{\left(\begin{smallmatrix} \mathcal{I} \\ \mathcal{I}\mathcal{L} \end{smallmatrix}\right)}^{\left(\begin{smallmatrix} w \\ \mathcal{I}\mathcal{J} \end{smallmatrix}\right)} a_{\mathcal{I}\mathcal{J}}^w(i, j) \right) E_{ij}, \quad (26)$$

Furthermore Formulae (18),(2) and (3) yield

$$\sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} \left(\sum_{l \in \mathcal{L}} (b_{\mathcal{I}\mathcal{L}}^u(i, l) h_{lj} - h_{il} b_{\mathcal{L}\mathcal{J}}^u(l, j)) \right) E_{ij} = \sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} \left(\sum_w c_{\mathcal{I}\mathcal{J}}^{uw} a_{\mathcal{I}\mathcal{J}}^w(i, j) \right) E_{ij}. \quad (27)$$

Denote $(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w)$ still by (p, q) . Since $a_{\mathcal{I}\mathcal{J}}^w(p, q) = 1$, $a_{\mathcal{I}\mathcal{J}}^{w'}(p, q) = 0$, $\forall w' \neq w$, we have

$$\begin{aligned}
Dd((\rho_{\mathcal{I}\mathcal{J}}^w)^*) &\stackrel{(19),(27)}{=} \sum_u \left(\sum_{l \in \mathcal{L}} (b_{\mathcal{I}\mathcal{L}}^u(p, l)h_{lq} - h_{pl}b_{\mathcal{L}\mathcal{J}}^u(l, q)) \right) (\zeta_{\mathcal{I}\mathcal{J}}^u)^* \\
&= \sum_{l \in \mathcal{L}} \left(\sum_u b_{\mathcal{I}\mathcal{L}}^u(p, l)(\zeta_{\mathcal{I}\mathcal{J}}^u)^* \right) h_{lq} - \sum_{l \in \mathcal{I}} h_{pl} \left(\sum_u b_{\mathcal{L}\mathcal{J}}^u(l, q)(\zeta_{\mathcal{I}\mathcal{J}}^u)^* \right) \\
&\stackrel{(22)}{=} \sum_l (s_{pl}h_{lq} - h_{pl}s_{lq}).
\end{aligned}$$

Moreover,

$$\begin{aligned}
&\sum_{\mathcal{L}, u, v} \left({}^l c_{\left(\begin{smallmatrix} \mathcal{I}\mathcal{J} \\ \mathcal{I}\mathcal{L} \end{smallmatrix} \right) \left(\mathcal{L}\mathcal{J} \right)}^{\left(\begin{smallmatrix} w \\ \mathcal{I}\mathcal{J} \end{smallmatrix} \right)} (\zeta_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\rho_{\mathcal{L}\mathcal{J}}^v)^* - {}^r c_{\left(\begin{smallmatrix} \mathcal{I}\mathcal{J} \\ \mathcal{I}\mathcal{L} \end{smallmatrix} \right) \left(\mathcal{L}\mathcal{J} \right)}^{\left(\begin{smallmatrix} w \\ \mathcal{I}\mathcal{J} \end{smallmatrix} \right)} (\rho_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^v)^* \right) \\
&\stackrel{(25),(26)}{=} \sum_{\mathcal{L}, u, v} \sum_{l \in \mathcal{L}} (b_{\mathcal{I}\mathcal{L}}^u(p, l)a_{\mathcal{L}\mathcal{J}}^v(l, q)(\zeta_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\rho_{\mathcal{L}\mathcal{J}}^v)^* - a_{\mathcal{I}\mathcal{L}}^u(p, l)b_{\mathcal{L}\mathcal{J}}^v(l, q)(\rho_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^v)^*) \\
&\stackrel{(22),(23)}{=} \sum_l (x_{pl}z_{lq} - z_{pl}x_{lq}).
\end{aligned}$$

Lemma 3.6.1 The Formula (24) is equivalent to

$$(\rho_{\mathcal{I}\mathcal{J}}^w)^* \otimes E_{\mathcal{J}}^* - E_{\mathcal{I}}^* \otimes (\rho_{\mathcal{I}\mathcal{J}}^w)^* = \delta_1(\rho_{\mathcal{I}\mathcal{J}}^w)^*$$

□

On the other hand, The Formulae (14) and (2) show that

$$\sum_{\mathcal{I} \ni i < j \in \mathcal{J}} \left(\sum_l b_{\mathcal{I}\mathcal{L}}^u(i, l)b_{\mathcal{L}\mathcal{J}}^v(l, j) \right) E_{ij} = \sum_{\mathcal{I} \ni i < j \in \mathcal{J}} \left(\sum_w c_{\left(\begin{smallmatrix} \mathcal{I}\mathcal{J} \\ \mathcal{I}\mathcal{L} \end{smallmatrix} \right) \left(\mathcal{L}\mathcal{J} \right)}^{\left(\begin{smallmatrix} w \\ \mathcal{I}\mathcal{J} \end{smallmatrix} \right)} b_{\mathcal{I}\mathcal{J}}^w(i, j) \right) E_{ij}. \quad (28)$$

If the basis element $(\zeta_{\mathcal{I}\mathcal{J}}^w)^*$ is given by Formula (2) of 2.3, then the $(p_{\mathcal{I}\mathcal{J}}^w, q_{\mathcal{I}\mathcal{J}}^w)$ -entry of $R_0 \cdot R_0$ equals

$$\begin{aligned}
\sum_{\bar{p} \leq l \leq \bar{q}} x_{\bar{p}l}x_{l\bar{q}} &\stackrel{(22)}{=} (\zeta_{\mathcal{I}\mathcal{J}}^w)^* \otimes E_{\mathcal{J}}^* + E_{\mathcal{I}}^* \otimes (\zeta_{\mathcal{I}\mathcal{J}}^w)^* + \sum_{\mathcal{L}, u, v} \sum_{l \in \mathcal{L}} b_{\mathcal{I}\mathcal{L}}^u(p, l)b_{\mathcal{L}\mathcal{J}}^v(l, q)(\zeta_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^v)^* \\
&\stackrel{(28)}{=} (\zeta_{\mathcal{I}\mathcal{J}}^w)^* \otimes E_{\mathcal{J}}^* + E_{\mathcal{I}}^* \otimes (\zeta_{\mathcal{I}\mathcal{J}}^w)^* + \sum_{\mathcal{L}, u, v} c_{\left(\begin{smallmatrix} \mathcal{I}\mathcal{J} \\ \mathcal{I}\mathcal{L} \end{smallmatrix} \right) \left(\mathcal{L}\mathcal{J} \right)}^{\left(\begin{smallmatrix} w \\ \mathcal{I}\mathcal{J} \end{smallmatrix} \right)} (\zeta_{\mathcal{I}\mathcal{L}}^u)^* \otimes (\zeta_{\mathcal{L}\mathcal{J}}^v)^*
\end{aligned}$$

Lemma 3.6.2 $\mu(\zeta_{\mathcal{I}\mathcal{J}}^w)^* = \sum_{\bar{p} \leq l \leq \bar{q}} x_{\bar{p}l}x_{l\bar{q}}.$

□

Theorem 3.6.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem with a triangular basis (A, B) . Then

$$\delta_1(\rho_{\mathcal{I}\mathcal{J}}^w)^* = \left(\sum_{p < l} x_{pl} h_{lq} - \sum_{l < q} h_{pl} x_{lq} \right) + \left(\sum_{p < l} x_{pl} z_{lq} - \sum_{l < q} z_{pl} x_{lq} \right)$$

where $p = p_{\mathcal{I}\mathcal{J}}^w$, $q = q_{\mathcal{I}\mathcal{J}}^w$. In particular, for the first basis element $\rho \in A$,

$$\delta(\rho^*) = \sum_{p < l} x_{pl} h_{lq} - \sum_{l < q} h_{pl} x_{lq}$$

On the other hand,

$$\delta_2(\zeta_{\mathcal{I}\mathcal{J}}^w)^* = \sum_{\bar{p} < l < \bar{q}} x_{\bar{p}l} x_{l\bar{q}}$$

where $\bar{p} = \bar{p}_{\mathcal{I}\mathcal{J}}^w$, $\bar{q} = \bar{q}_{\mathcal{I}\mathcal{J}}^w$. □

Corollary 3.6.1 The boc $\mathfrak{A}$ is normal and triangular, see [Ro, BK, CB1].

Proof. The normality is given by Formula (17) in the sense of [Ro], which follows from the algebra $\mathcal{K}$ being upper triangular. The triangularity follows from the triangularity of the basis (A, B) . In fact, δ_1 on A^* is triangular, i.e. $(\rho_{\mathcal{I}\mathcal{J}}^w)^* \succ (\rho_{\mathcal{I}\mathcal{J}}^u)^*$ and $(\rho_{\mathcal{I}\mathcal{J}}^v)^*$, by Proposition 2.4.1; δ_2 on B^* is also triangular i.e. $(\zeta_{\mathcal{I}\mathcal{J}}^w)^* \succ (\zeta_{\mathcal{I}\mathcal{J}}^u)^*$ and $(\zeta_{\mathcal{I}\mathcal{J}}^v)^*$, since $\zeta_{\mathcal{I}\mathcal{J}}^w \succ \zeta_{\mathcal{I}\mathcal{J}}^u$ and $\zeta_{\mathcal{I}\mathcal{J}}^v$ in Formula (14), see [CB1, BK]. □

4 Freely parameterized bimodule problems

4.1 Parameters

Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem, and $(*)$ be a reduction sequence. It has been shown in 2.5, that every Weyr matrix in the third sequence of $(*)$ possesses some fixed eigenvalues. From now on we allow some of the eigenvalues of Weyr matrices to be *parameters*, and $(\mathcal{K}^r, \mathcal{M}^r, H^r)$ is said to be a *parameterized bimodule problem*. Suppose that

$$P^r = \{\lambda_1, \lambda_2, \dots, \lambda_{i_r}\}$$

is the set of parameters appearing in $H_{\underline{n}}^r$, then $P^1 \subseteq P^2 \subseteq \dots \subseteq P^s$. Denote the domain of P^r by D^r , then $D^r \subseteq k^{i_r}$, $r = 1, 2, \dots, s$.

Proposition 4.1.1 [S]. D^s is determined by some polynomial equations and inequalities in i_s indeterminates. Therefore it is a locally closed subset of k^{i_s} .

Proof. We use induction on r . The case of $r = 0$ is trivial since $H_{\underline{n}} = 0$ has no parameter. Suppose the assertion is true for r . Now we perform the $(r + 1)$ -reduction. Denote by C the coefficient matrix of equation system II of Definition 2.2.1 at $(\mathcal{P}_r, \mathcal{Q}_r)$; by $\bar{C}$ that of equation system II and equation (8) of 2.6 at $(\mathcal{P}_r, \mathcal{Q}_r)$.

Case 1. Equation system II of Definition 2.2.1 implies the equation 2.6(8), if and only if $\text{rank} C = \text{rank} \bar{C}$, which is determined by some polynomial equations in $x_1, x_2, \dots, x_{i_r}$.

Case 2. Equation system II of Definition 2.2.1 does not imply the equation 2.6(8), if and only if $\text{rank} C + 1 = \text{rank} \bar{C}$, which is determined by a polynomial inequality in $x_1, x_2, \dots, x_{i_r}$. $\square$

Remarks. (1) The parameters in $P^r \setminus P^{r-1}$ are free at the r -th reduction unless otherwise assumption, since the restriction equations or inequalities for these parameters only appear in the reductions after the r -th step.

(2) If $N^s = (0)$ in $(*)$, then the parameterized matrix $H_{\underline{n}}^\infty(\lambda_1, \lambda_2, \dots, \lambda_{i_s})$ can be regarded as a *representation variety* over $(\mathcal{K}, \mathcal{M})$, which is locally closed.

(3) Note that it is difficult to define a representation category over a parameterized bimodule problem in general, (see examples of 4.6). Thus we will focus on a very special situation of the parameterized bimodule problems defined in the next subsection.

4.2 Freely parameterized bimodule Problems

Definition 4.2.1 A bimodule problem with i parameters is said to be *freely parameterized*, provided

- (1) for any $\mathcal{I} \in T/\sim$, there exists at most one parameter λ attached to $\mathcal{I}$;
- (2) all parameters are algebraically independent, and the domain

$$D = k \setminus \{\text{the roots of } g(\lambda_1, \dots, \lambda_i) = g_1(\lambda_1) \cdots g_i(\lambda_i)\}$$

□

An equivalent class $\mathcal{I} \in T/\sim$ is said to be *non-trivial* if there exists a parameter $\lambda_{\mathcal{I}}$ attached to $\mathcal{I}$; otherwise, $\mathcal{I}$ is said to be *trivial*. Suppose that $\{\mathcal{I}_1, \dots, \mathcal{I}_i\}$ is a complete set of non-trivial equivalent classes of $(T, \sim)$, and $(\mathcal{J}_1, \dots, \mathcal{J}_j)$ is a complete set of trivial ones respectively.

Now we construct a bocs based on the freely parameterized bimodule problem $(\mathcal{K}, \mathcal{M}, H)$. Let Γ' be a category with indecomposable objects $\mathcal{I}_1, \dots, \mathcal{I}_i; \mathcal{J}_1, \dots, \mathcal{J}_j$ and morphisms

$$\Gamma'(\mathcal{I}_l, \mathcal{I}_l) = k[\lambda_l, g_l(\lambda_l)^{-1}], \quad l = 1, \dots, i; \quad \Gamma'(\mathcal{J}_l, \mathcal{J}_l) = k, \quad l = 1, \dots, j.$$

Then Γ' is equivalent to the following algebra

$$k[\lambda_1, g_1(\lambda_1)^{-1}] \times \cdots \times k[\lambda_i, g_i(\lambda_i)^{-1}] \times k_1 \times \cdots \times k_j,$$

where $k_1 = \cdots = k_j = k$. Γ' is called a *minimal category*, and can be shown in the following diagram [CB1]:

$$\begin{array}{ccccccc} \lambda_1 & \circlearrowleft & \lambda_2 & \circlearrowleft & \cdots & \lambda_i & \circlearrowleft \\ & \mathcal{I}_1 & & \mathcal{I}_2 & & \mathcal{I}_i & \end{array} \quad \begin{array}{ccccccc} \bullet & & \bullet & & \bullet & & \bullet \\ \mathcal{J}_1 & & \mathcal{J}_2 & & \cdots & & \mathcal{J}_j \end{array}$$

Denote A^* by $\{a_1, \dots, a_n\}$, B^* by $\{v_1, \dots, v_m\}$ according to the order of Definition 2.4.1, in order to simplify the notations. Then $\mathfrak{A}$ is said to be a *layered boc*s with a layer


$$L = (\Gamma'; \omega; a_1, a_2, \dots, a_n; v_1, v_2, \dots, v_m),$$

where $\omega : \Gamma' \rightarrow \Omega$ is a Γ' - Γ' -bimodule map given by $\omega(E_{\mathcal{I}}) = (E_{\mathcal{I}})^*$ for any $\mathcal{I} \in T/\sim$. Note that Γ' is of infinite dimension if $i > 0$.

A layered boc can be illustrated as a *differential biquiver* as follows. The set of vertices of the biquiver is $T/\sim$, the set of solid arrows is $\{\lambda_1, \dots, \lambda_i; a_1, \dots, a_n\}$ and the set of dotted arrows is $\{v_1, \dots, v_m\}$ [CB2]. The examples given in 2.1 correspond to the following differential biquivers, if the k -basis of $\text{rad}(\Lambda)$, a, b, c, d, f still stand for the k -basis of $\text{rad}(\tilde{\Lambda})$, as well as $\begin{pmatrix} 0 & \text{rad} \tilde{\Lambda} \\ 0 & 0 \end{pmatrix}$ with $a \prec b \prec c \prec d \prec f$, and a^*, b^*, c^*, d^*, f^* are


dual basis.

Example 1.




$$\begin{aligned}\delta(a^*) &= 0, \\ \delta(b^*) &= 0, \\ \delta(c^*) &= 0.\end{aligned}$$

Example 2.




$$\begin{aligned}\delta(a^*) &= 0, \\ \delta(b^*) &= 0, \\ \delta(c^*) &= 0.\end{aligned}$$

Example 3.



$$\begin{aligned}\delta(a^*) &= 0, \\ \delta(b^*) &= 0, \\ \delta(c^*) &= v'b^* - b^*v, \\ \delta(d^*) &= \alpha v'a^* - a^*v + u'b^* - \alpha b^*u.\end{aligned}$$

Example 4.



$$\begin{aligned}\delta(a^*) &= 0, \\ \delta(b^*) &= 0, \\ \delta(c^*) &= 0, \\ \delta(d^*) &= 0, \\ \delta(f^*) &= 0.\end{aligned}$$

A freely parameterized bimodule problem is said to be *minimal*, if $\mathcal{M} = \{(0)\}$. Correspondingly a triple $(N, R, \underline{n})$ is said to be *minimal*, if $N = (0)$. Moreover, a layered boc $\mathfrak{A} = (\Gamma, \Omega)$ is called a *minimal boc*, if $\Gamma = \Gamma'$, then the layer $L = (\Gamma'; \omega; v_1, \dots, v_m)$ without any solid arrows. The corresponding differential byquiver consists of isolated vertices, vertices with single solid loops, and dotted arrows.

From now on we will not worry about any differences between a freely parameterized

bimodule problem and its layered boc. We will use freely the two concepts depending on convenience of concerned problems.

Remark Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem, and $(*)$ be a reduction sequence. We usually obtain a freely parameterized bimodule problem as the following way: every parameter appears in a loop reduction, i.e. $\overline{N}_{p_{r-1}q_{r-1}} = W \oplus (\lambda)$ with W being a Weyr matrix of fixed eigenvalues and the domain $D^r|_{\{\lambda\}} = k \setminus \{\text{the eigenvalues of } W\}$, for some step $1 \leq r < s$. Then the end term $(N^s, R^s, \underline{n}^s)$ of $(*)$ is said to be a *freely parameterized triple*. In this case, $(\mathcal{K}^s, \mathcal{M}^s, H^s)$, the end term of the corresponding sequence $(**)$, is a freely parameterized bimodule problem.

4.3 The differential of the first arrow

We will illustrate all the possibilities of the differential of the first arrow of a layered boc in this subsection, which are not mentioned before in any references.

Let $\mathfrak{A} = (\Gamma, \Omega)$ be a boc with a layer $L = (\Gamma'; \omega; a_1, \dots, a_n; v_1, \dots, v_m)$. Then the differential $\delta(a_1)$ of the first arrow $a_1 : \mathcal{P} \rightarrow \mathcal{Q}$ has the following possibilities:

A1. $\lambda \begin{array}{c} \mathcal{P} \\ \curvearrowright \end{array} \bullet \begin{array}{c} \mathcal{P} \\ \curvearrowright \end{array} a_1, \mathcal{P} = \mathcal{Q}, \Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda, g_{\mathcal{P}}(\lambda)^{-1}]$ or

$\lambda \begin{array}{c} \mathcal{P} \\ \curvearrowright \end{array} \bullet \xrightarrow{a_1} \begin{array}{c} \mathcal{Q} \\ \curvearrowright \end{array} \mu, \mathcal{P} \neq \mathcal{Q}, \Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda, g_{\mathcal{P}}(\lambda)^{-1}]$ and $\Gamma'(\mathcal{Q}, \mathcal{Q}) = k[\mu, g_{\mathcal{Q}}(\mu)^{-1}]$. Then

$$\delta(a_1) = \sum_{j=1}^m f_j(\lambda, \mu) v_j \quad (29)$$

where we may assume that $f_j(\lambda, \mu) \in k[\lambda, \mu]$ by dividing v_j by some power of $g(\lambda)$; and λv_j stands for the left multiplication of v_j by λ , and μv_j for the right multiplication by λ in the first case or by μ in the second case.

Recall from [CB1], let $f(\lambda, \mu)$ be the highest common factor of the $f_j(\lambda, \mu)$ and let $q_j(\lambda, \mu) = f_j(\lambda, \mu)/f(\lambda, \mu)$. Since $q_j(\lambda, \mu)$ are coprime, there are polynomials $s_j(\lambda, \mu)$ and a non-zero polynomial $c(\lambda)$, such that

$$c(\lambda) = \sum_{j=1}^m s_j(\lambda, \mu) q_j(\lambda, \mu) \quad (30)$$

in ring $k[\lambda, \mu]$. Thus

$$1 = \sum_{j=1}^m c(\lambda)^{-1} s_j(\lambda, \mu) q_j(\lambda, \mu)$$

in a Hermite ring $S = k[\lambda, \mu, g_{\mathcal{P}}(\lambda)^{-1} c(\lambda)^{-1}, g_{\mathcal{Q}}(\mu)^{-1}]$. So there is an invertible matrix Q in $M_m(S)$ with the first row $(q_j(\lambda, \mu))_{j=1, \dots, m}$, and we can make a change of basis of the

form

$$(w_1, w_2, \dots, w_m)^T = Q(v_1, v_2, \dots, v_m)^T$$

so that

$$\delta(a_1) = f(\lambda, \mu)w_1. \quad (31)$$

A2. $\lambda \begin{array}{c} \curvearrowright \\ \bullet \end{array} \xrightarrow{a_1} \mathcal{Q}$, $\mathcal{P} \neq \mathcal{Q}$, $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda, g_{\mathcal{P}}(\lambda)^{-1}]$, $\Gamma'(\mathcal{Q}, \mathcal{Q}) = k$.
(Or dually, $\mathcal{P} \xrightarrow{a_1} \mathcal{Q} \begin{array}{c} \curvearrowright \\ \bullet \end{array} \mu$, $\Gamma'(\mathcal{P}, \mathcal{P}) = k$, $\Gamma'(\mathcal{Q}, \mathcal{Q}) = k[\mu, g_{\mathcal{Q}}(\mu)^{-1}]$). Then

$$\delta(a_1) = \sum_{j=1}^m f_j(\lambda)v_j, \quad (32)$$

where $f_j(\lambda) \in k[\lambda]$. If $f(\lambda)$ is the highest common factor of $f_j(\lambda)$, then $q_j(\lambda) = f_j(\lambda)/f(\lambda)$, $j = 1, \dots, m$, are coprime, and there are polynomials $s_j(\lambda)$ such that $1 = \sum_{j=1}^m s_j(\lambda)q_j(\lambda)$. After a basis transformation as in Case 1 given by an invertible matrix $Q \in M_m(k[\lambda, g_{\mathcal{P}}(\lambda)^{-1}])$, we have

$$\delta(a_1) = f(\lambda)w_1. \quad (33)$$

A3. $a_1 \begin{array}{c} \curvearrowright \\ \bullet \end{array} \mathcal{P}$, $\mathcal{P} = \mathcal{Q}$, $\Gamma'(\mathcal{P}, \mathcal{P}) = k$; or
 $\mathcal{P} \xrightarrow{a_1} \mathcal{Q}$, $\mathcal{P} \neq \mathcal{Q}$, $\Gamma'(\mathcal{P}, \mathcal{P}) = k$, $\Gamma'(\mathcal{Q}, \mathcal{Q}) = k$. Then

$$\delta(a_1) = \sum_{j=1}^m f_j v_j, \quad (34)$$

where $f_j \in k$. After a basis transformation given by an invertible matrix $Q \in M_m(k)$, we have $\delta(a_1) = w_1$, or $\delta(a_1) = 0$.

Proposition 4.3.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem, and $(\mathcal{K}', \mathcal{M}', H')$ be induced from $(\mathcal{K}, \mathcal{M}, H)$ by one of the 3 reductions of 3.1. Let $a_1 : \mathcal{P} \rightarrow \mathcal{Q}$ be the first arrow. Then $(\mathcal{K}', \mathcal{M}', H')$ is still freely parameterized and preserves all the free parameters if and only if $\delta(a_1)$ and the reduction for a_1 has the following forms according to A1, A2, A3 respectively.

A1 (1) $\delta(a_1) = 0$, i.e. $f_j(\lambda, \mu) = 0$ for all j in Formula (29).

When $\mathcal{P} = \mathcal{Q}$, we set $\overline{N}_{pq} = (\lambda^0)$ for a fixed eigenvalue λ^0 . Then $g'_{\mathcal{P}}(\lambda) = g_{\mathcal{P}}(\lambda)$.

When $\mathcal{P} \neq \mathcal{Q}$, we set $\overline{N}_{pq} = (0)$, Then $g'_{\mathcal{P}}(\lambda) = g_{\mathcal{P}}(\lambda)$, $g'_{\mathcal{Q}}(\mu) = g_{\mathcal{Q}}(\mu)$.

(2) $\delta(a_1) = f(\lambda, \mu)w_1 \neq 0$ in Formula (30), where $f(\lambda, \mu)$ must be equal to

$f_{\mathcal{P}}(\lambda)f_{\mathcal{Q}}(\mu)$, and $c(\lambda)$ is given by Formula (30).

When $\mathcal{P} = \mathcal{Q}$, we set $\overline{N}_{pq} = \emptyset_{1 \times 1}$. Then $g'_{\mathcal{P}}(\lambda) = g_{\mathcal{P}}(\lambda)c(\lambda)f_{\mathcal{P}}(\lambda)$.

When $\mathcal{P} \neq \mathcal{Q}$, we set $\overline{N}_{pq} = \emptyset_{1 \times 1}$. Then $g'_{\mathcal{P}}(\lambda) = g_{\mathcal{P}}(\lambda)c(\lambda)f_{\mathcal{P}}(\lambda)$,
 $g'_{\mathcal{Q}}(\mu) = g_{\mathcal{Q}}(\mu)f_{\mathcal{Q}}(\mu)$.

A2. (1) $\delta(a_1) = 0$, i.e. $f_j(\lambda) = 0$ for all j in Formula (32), we set

$$\overline{N}_{pq} = (0 \ \cdots \ 0 \ 0)_{1 \times n_q}, \quad \text{then } g'_{\mathcal{P}}(\lambda) = g_{\mathcal{P}}(\lambda); \quad \text{or}$$

$$\overline{N}_{pq} = (0 \ \cdots \ 0 \ 1)_{1 \times n_q}, \quad g'_{(\mathcal{P} \cup \mathcal{Q})'}(\lambda) = g_{\mathcal{P}}(\lambda).$$

(2) $\delta(a_1) = f(\lambda)w_1 \neq 0$ in Formula (33), we set

$$\overline{N}_{pq} = \emptyset_{1 \times n_q} \quad \text{then } g'_{\mathcal{P}}(\lambda) = g_{\mathcal{P}}(\lambda)f(\lambda).$$

Or dually,

(1') $\delta(a_1) = 0$, we set

$$\overline{N}_{pq} = (0 \ \cdots \ 0 \ 0)_{1 \times n_p}^T, \quad \text{then } g'_{\mathcal{Q}}(\mu) = g_{\mathcal{Q}}(\mu); \quad \text{or}$$

$$\overline{N}_{pq} = (1 \ \cdots \ 0 \ 0)_{1 \times n_p}^T, \quad g'_{(\mathcal{P} \cup \mathcal{Q})'}(\mu) = g_{\mathcal{Q}}(\mu).$$

(2') $\delta(a_1) = f(\mu)w_1 \neq 0$, we set $\overline{N}_{pq} = \emptyset_{n_p \times 1}$, then $g'_{\mathcal{Q}}(\mu) = g_{\mathcal{Q}}(\mu)f(\mu)$.

A3. (1) $\delta(a_1) = 0$, we set

$$\overline{N}_{pq} = \begin{cases} W, & \text{when } \mathcal{P} = \mathcal{Q}, \\ \begin{pmatrix} 0 & I \\ 0 & 0 \end{pmatrix}, & \text{when } \mathcal{P} \neq \mathcal{Q}, \end{cases}$$

where W is a Weyr matrix with some fixed eigenvalues.

(2) $\delta(a_1) = w$, we set $\overline{N}_{pq} = \emptyset$.

(3) $\delta(a_1) = 0$ and $\mathcal{P} = \mathcal{Q}$, then $\overline{N}_{pq}$ may be equal to $W \oplus (\lambda)$. Thus a new parameter λ appears at this step, and the domain of λ is

$$k \setminus \{\text{the eigenvalues of } W\}$$

Proof. In case A1, we must have $n_{\mathcal{P}} = 1$, $n_{\mathcal{Q}} = 1$, since there are free parameters λ attached to $\mathcal{P}$, and μ to $\mathcal{Q}$ respectively.

(1) If $\mathcal{P} = \mathcal{Q}$, $\overline{N}_{pq}$ can not be taken as a parameter, and if $\mathcal{P} \neq \mathcal{Q}$, $\overline{N}_{pq}$ can not be taken as 1×1 identity matrix because of condition (2) of Definition 4.2.1.

(2) If $f(\lambda, \mu) = 0$, λ, μ would be algebraically dependent, which is a contradiction to condition (3) of Definition 4.2.1. Thus $f(\lambda, \mu) \neq 0$. And the condition (3) of 4.2.1 forces that $f(\lambda, \mu) = f_{\mathcal{P}}(\lambda)f_{\mathcal{Q}}(\mu)$. Thus the proof of A1 is completed.

Cases A2 and A3 are easy. $\square$

Remark. The reduction sequences of freely parameterized bimodule problems is very special in the set of general reduction sequences of parameterized bimodule problems. In fact, in Formula (31) of case A1, if $f(\lambda, \mu) = f_{\mathcal{P}}(\lambda)f_{\mathcal{Q}}(\mu)f_0(\lambda, \mu)$, such that $f_0(\lambda, \mu) \in k[\lambda, \mu]$ is not a constant, and does not contain any non-constant factor in $k[\lambda]$ or $k[\mu]$, then we are not able to continue the reduction, such that the induced triple is still freely parameterized and preserves the parameters λ and μ . We will see in 5.2 that such a situation can not occur in tame case, but really occurs in wild case see Examples in 7.4 and 7.6.

Corollary 4.3.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem having a reduction sequence $(*)$ of freely parameterized triples given in the Remark of 4.2, and $\hat{n}^s$ be a size vector of $(T^s, \sim^s)$ such that $\hat{n}_{\mathcal{I}}^s = 1$ or 0 whenever $\mathcal{I} \in T^s / \sim^s$ is non-trivial. Then $(\hat{*})$ of Corollary 3.3.1 is still a reduction sequence of freely parameterized triples.

Proof. The only problem is, whether the domain at each step of $(*)$ can be regarded as a domain at the same step of $(\hat{*})$. Suppose it is the case up to $(\hat{N}^r, \hat{R}^r, \hat{n}^r)$. Now consider the $(r+1)$ -th reduction. Let

$$C^r = \{\mathcal{I} \in T^r / \sim^r \mid \hat{n}_{\mathcal{I}}^r = 0\}.$$

(1) If $\mathcal{P}_r \in C^r$ or $\mathcal{Q}_r \in C^r$, we have an identity functor from $(\hat{N}^r, \hat{R}^r, \hat{n}^r)$ to $(\hat{N}^{r+1}, \hat{R}^{r+1}, \hat{n}^{r+1})$ by Proposition 3.2.1. Thus $\hat{D}^r = \hat{D}^{r+1}$. However there may be some restriction from D^r to D^{r+1} or there may be a new parameter appearing in D^{r+1} .

(2) If $\mathcal{P}_r \notin C^r$, $\mathcal{Q}_r \notin C^r$, the equation system II of Definition 2.2.1 and equation (8) of 2.6 at $(\mathcal{P}_r, \mathcal{Q}_r)$ are the same both for $\mathcal{K}^r$ and $\hat{\mathcal{K}}^r$ by Proposition 3.2.1 once again. Thus items (1) and (2) of A1, A2, A3 of Proposition 4.3.1 give the same restriction from $\hat{D}^r$ to $\hat{D}^{r+1}$, as that from D^r to D^{r+1} .

In item (3) of A3, the restriction on the new parameter λ in $\hat{D}^{r+1}$ is equal to or less than that in D^{r+1} , since the eigenvalues of $\hat{W}$ may be equal to or less than those of W by Proposition 3.2.1.

By induction the sequence of the domains of $(*)$ suits that of $(\hat{*})$ well. Therefore $(\hat{*})$ is also a reduction sequence of freely parameterized triples. $\square$

If 3.3 $(**)$ is a sequence of freely parameterized bimodule problems, there exists a

sequence of corresponding layered bocses:

$$(***) \quad \mathfrak{A}, \mathfrak{A}^1, \dots, \mathfrak{A}^r, \dots, \mathfrak{A}^s.$$

4.4 Valuation matrices

Let $(\mathcal{K}, \mathcal{M}, H)$ be a parameterized bimodule problem with a set of parameters $\{\lambda_1, \dots, \lambda_i\}$ of domain D . It is natural to ask whether the parameters can be regarded as the solid arrows. If the answer is yes, then $\lambda_{\mathcal{I}_1}, \lambda_{\mathcal{I}_2}, \dots, \lambda_{\mathcal{I}_i}$ and $a_1, a_2, \dots, a_n$ are free generators of Γ over some trivial category $k \times \dots \times k$, and we are able to define a wider representation category over $(\mathcal{K}, \mathcal{M}, H)$ such that $\lambda_{\mathcal{I}_i}$ can be taken as Weyr matrices of fixed eigenvalues. This idea will be realized in the rest of the section without difficulty when $(\mathcal{K}, \mathcal{M}, H)$ is a freely parameterized bimodule problem. And a more detailed observation shows that the 2 conditions of Definition 4.2.1 are necessary for our purpose: Condition (1) gives the minimality of Γ' ; Condition (2) is the most important condition in order to define the morphisms in a wilder category (see Examples in 4.6).

Definition 4.4.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem with i free parameters.

(1) By a *valuation of a parameter* $\lambda_{\mathcal{I}}$ we mean the assigning to $\lambda_{\mathcal{I}}$ a Weyr matrix

$$W(\lambda_{\mathcal{I}}^0) = W_{\lambda_{\mathcal{I}}^1} \oplus \dots \oplus W_{\lambda_{\mathcal{I}}^{\alpha_{\mathcal{I}}}}$$

(see 2.5), where $\lambda_{\mathcal{I}}^1, \dots, \lambda_{\mathcal{I}}^{\alpha_{\mathcal{I}}}$ are pairwise different eigenvalues taken from the domain $D|_{\{\lambda_{\mathcal{I}}\}}$. When $\mathcal{I}$ runs over the non-trivial equivalent classes of $T/\sim$, we then obtain a set of valuations:

$$W(\lambda_{\mathcal{I}_1}^0), W(\lambda_{\mathcal{I}_2}^0), \dots, W(\lambda_{\mathcal{I}_i}^0).$$

(2) Define a size vector $\underline{m}$ of $(T, \sim)$, such that $m_{\mathcal{I}}$ equals the size of $W(\lambda_{\mathcal{I}}^0)$ whenever the parameter $\lambda_{\mathcal{I}}$ is attached to $\mathcal{I}$, and m_j can be taken as any non-negative integer for any trivial equivalent class $\mathcal{J} \in T/\sim$. Then we have a matrix $H_{\underline{m}}$, which is not freely parameterized if there exists some non-trivial vertex $\mathcal{I}$, such that $m_{\mathcal{I}} > 1$. Write $\hat{\underline{m}} = \underline{m}$, then $(\hat{N}, \hat{R}, \hat{\underline{m}})$ is a triple of $(\mathcal{K}, \mathcal{M}, H)$.

(3) Let $\underline{e}_{\mathcal{I}}$ be the size vector determined by $W(\lambda_{\mathcal{I}}^0)$ (see 2.5). We define a vector $\tilde{\underline{m}}$ such that $\tilde{m}_{\mathcal{J}} = m_{\mathcal{J}}$ for any trivial $\mathcal{J} \in T/\sim$, and $\tilde{m}_{\mathcal{I}} = (\underline{e}_{\mathcal{I}})$ for any non-trivial $\mathcal{I} \in T/\sim$. A matrix $\tilde{H}_{\tilde{\underline{m}}}(\lambda^0)$ is said to be a *valuated matrix* of H , provided that $H_{\tilde{\underline{m}}}(\lambda^0)$ is obtained from $H_{\underline{m}}$ by substituting each $W(\lambda_{\mathcal{I}}^0)$ for $\lambda_{\mathcal{I}} I_{m_{\mathcal{I}}}$.

(4) A triple $(\tilde{N}, \tilde{R}, \tilde{\underline{m}})$ is constructed based on $(\hat{N}, \hat{R}, \hat{\underline{m}})$, such that $\tilde{N} = \hat{N}$, but $\tilde{N}$ has a size vector $\tilde{\underline{m}}$; and $\tilde{R}$ satisfies the matrix equation

$$\tilde{R} \tilde{H}_{\tilde{\underline{m}}}(\lambda^0) \equiv_{\prec(p,q)} \tilde{H}_{\tilde{\underline{m}}}(\lambda^0) \tilde{R}.$$

It will be proved in the next subsection that the triple $(\tilde{N}, \tilde{R}, \tilde{m})$ is induced by a sequence of reductions.

4.5 The reduction sequence towards a valuation matrix

Lemma 4.5.1 Let $f(\lambda, \mu) = \sum_{i,j \geq 0} a_{ij} \lambda^i \mu^j \in k[\lambda, \mu]$. Let $X = (x(\alpha, \beta))_{m \times n} \in M_{m \times n}(k)$, and L, R be upper triangular nilpotent matrices. Then the (α, β) -entry in the matrix

$$\sum_{i,j \geq 0} a_{ij} (\lambda I_m + L)^i X (\mu I_n + R)^j$$

equals

$$\sum_{i,j \geq 0} \left(a_{ij} \lambda^i \mu^j x(\alpha, \beta) + \sum_{(\alpha', \beta') \prec (\alpha, \beta)} d(\lambda, \mu, \alpha', \beta') x(\alpha', \beta') \right),$$

where $d(\lambda, \mu, \alpha', \beta')$ are polynomials in λ, μ as coefficients of $x(\alpha', \beta')$, and $(\alpha', \beta') \prec (\alpha, \beta)$ are given under the matrix order of Definition 2.3.1. $\square$

Let $\sum_{i,j} c_{ij}(\lambda, \mu) x_{ij} = 0$ be a linear equation in the indeterminates x_{ij} , and the coefficients $c_{ij}(\lambda, \mu) \in k[\lambda, \mu]$. Let $W(\underline{\lambda}^0) = W_{\lambda^1} \oplus \cdots \oplus W_{\lambda^g}$, $W(\underline{\mu}^0) = W_{\mu^1} \oplus \cdots \oplus W_{\mu^h}$ be two Weyr matrices with sizes m and n respectively. Let matrix $X_{ij} = (X_{ij}^{\gamma\eta})_{g \times h}$ be partitioned according to $W(\underline{\lambda}^0)$ and $W(\underline{\mu}^0)$, i.e. $X_{ij}^{\gamma\eta}$ are $m_\gamma \times n_\eta$ matrices whenever the size of W_{λ^γ} is m_γ , and that of W_{μ^η} is n_η , thus $X_{ij}^{\gamma\eta} = (x_{ij}^{\gamma\eta}(\alpha, \beta))_{m_\gamma \times n_\eta}$. For example,

W_{λ^1}		X_{ij}^{11}	X_{ij}^{12}	X_{ij}^{13}	W_{μ^1}		
	W_{λ^2}	X_{ij}^{21}	X_{ij}^{22}	X_{ij}^{23}		W_{μ^2}	
							W_{μ^3}

Now suppose that we have a bimodule problem $(\mathcal{K}, \mathcal{M}, H = 0)$ and a reduction sequence $(*)$ of freely parameterized triples given in the Remark of 4.2. Consider the $(r+1)$ -th reduction of $(*)$. Let $a_1 : \mathcal{P}_r \rightarrow \mathcal{Q}_r$ be the first arrow of $(\mathcal{K}^r, \mathcal{M}^r, H^r)$, where $\mathcal{P}_r$ (resp. $\mathcal{Q}_r$) is either non-trivial with a free parameter λ (resp. μ) or trivial. Assume that the equation system II of Definition 2.2.1 at $(\mathcal{P}_r, \mathcal{Q}_r)$ for $\mathcal{K}^r$ is given by

$$(II) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} c_{ij}^l(\lambda, \mu) x_{ij} = 0,$$

and the equation (8) of 2.6 is given by

$$(8) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} c_{ij}(\lambda, \mu) x_{ij} = 0.$$

We denote the coefficient matrix of (II) by $C(\lambda, \mu)$; that of (II) and (8) by $C'(\lambda, \mu)$. Furthermore assume that we also have the matrix equations:

$$(\tilde{\text{II}}) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} c_{ij}^l(W(\underline{\lambda}^0), W(\underline{\mu}^0)) X_{ij} = 0,$$

$$(\tilde{8}) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} c_{ij}(W(\underline{\lambda}^0), W(\underline{\mu}^0)) X_{ij} = 0,$$

or equivalently,

$$(\tilde{\text{II}}_{\gamma\eta}) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} c_{ij}^l(W_{\lambda^\gamma}, W_{\mu^\eta}) X_{ij}^{\gamma\eta} = 0,$$

$$(\tilde{8}_{\gamma\eta}) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} c_{ij}(W_{\lambda^\gamma}, W_{\mu^\eta}) X_{ij}^{\gamma\eta} = 0$$

for $1 \leq \gamma \leq g$, $1 \leq \eta \leq h$. Which are equivalent to a system of linear equations:

$$(\tilde{\text{II}}_{\gamma\eta}(\alpha, \beta)) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} \left(c_{ij}^l(\lambda^\gamma, \mu^\eta) x_{ij}^{\gamma\eta}(\alpha, \beta) + \sum_{(\alpha', \beta') \prec (\alpha, \beta)} d_{ij}^l(\lambda^\gamma, \mu^\eta, \alpha', \beta') x_{ij}^{\gamma\eta}(\alpha', \beta') \right) = 0,$$

$$(\tilde{8}_{\gamma\eta}(\alpha, \beta)) : \sum_{\mathcal{P}_r \ni i < j \in \mathcal{Q}_r} \left(c_{ij}(\lambda^\gamma, \mu^\eta) x_{ij}^{\gamma\eta}(\alpha, \beta) + \sum_{(\alpha', \beta') \prec (\alpha, \beta)} d_{ij}(\lambda^\gamma, \mu^\eta, \alpha', \beta') x_{ij}^{\gamma\eta}(\alpha', \beta') \right) = 0,$$

for $1 \leq \alpha \leq m_\gamma$, $1 \leq \beta \leq n_\eta$, and each pair (γ, η) . Next we fix a pair (γ, η) , and for each index pair (α, β) , we list the equations at the (α, β) -entry. If we count the index pairs (α, β) according to the matrix order of Definition 2.3.1, then we obtain an equation system $\tilde{\Pi}_{\gamma\eta}^0 = \bigcup_{(\alpha, \beta)} \tilde{\Pi}_{\gamma\eta}(\alpha, \beta)$ whose coefficient matrix is

$$\tilde{C}(W_{\lambda^\gamma}, W_{\mu^\eta}) = \begin{pmatrix} C(\lambda^\gamma, \mu^\eta) & 0 & \cdots & 0 \\ * & C(\lambda^\gamma, \mu^\eta) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & C(\lambda^\gamma, \mu^\eta) \end{pmatrix}_{(m_\gamma n_\eta) \times (m_\gamma n_\eta)}.$$

And the coefficient matrix of $\tilde{\Pi}_{\gamma\eta}^1 = \bigcup_{(\alpha, \beta)} \tilde{\Pi}_{\gamma\eta}(\alpha, \beta) \cup (\tilde{8}_{\gamma\eta}(\alpha, \beta))$ is

$$\tilde{C}'(W_{\lambda^\gamma}, W_{\mu^\eta}) = \begin{pmatrix} C'(\lambda^\gamma, \mu^\eta) & 0 & \cdots & 0 \\ * & C'(\lambda^\gamma, \mu^\eta) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \cdots & C'(\lambda^\gamma, \mu^\eta) \end{pmatrix}_{(m_\gamma n_\eta) \times (m_\gamma n_\eta)}.$$

We denote the coefficient matrix of $\bigcup_{(\gamma,\eta)}(\tilde{\Pi}_{\gamma\eta}^0)$ by $\tilde{C}(W(\underline{\lambda}^0), W(\underline{\mu}^0))$, and that of $\bigcup_{(\gamma,\eta)}(\tilde{\Pi}_{\gamma\eta}^1)$ by $\tilde{C}'(W(\underline{\lambda}^0), W(\underline{\mu}^0))$. Then

$$\begin{aligned}\tilde{C}(W(\underline{\lambda}^0), W(\underline{\mu}^0)) &= \text{diag}(\tilde{C}(W_{\lambda^\gamma}, W_{\mu^\eta}))_{(\gamma,\eta)}, \\ \tilde{C}'(W(\underline{\lambda}^0), W(\underline{\mu}^0)) &= \text{diag}(\tilde{C}'(W_{\lambda^\gamma}, W_{\mu^\eta}))_{(\gamma,\eta)},\end{aligned}$$

Proposition 4.5.1 (1) If $\text{rank}C'(\lambda, \mu) = \text{rank}C(\lambda, \mu)$ in the domain D^{r+1} , then

$$\text{rank}\tilde{C}'(W_{\lambda^\gamma}, W_{\mu^\eta}) = \text{rank}\tilde{C}(W_{\lambda^\gamma}, W_{\mu^\eta}),$$

when $(\lambda^\gamma, \mu^\eta) \in D^{r+1}|_{(\lambda, \mu)}$. Moreover

$$\text{rank}\tilde{C}'(W(\underline{\lambda}^0), W(\underline{\mu}^0)) = \text{rank}\tilde{C}(W(\underline{\lambda}^0), W(\underline{\mu}^0))$$

when $(\lambda^\gamma, \mu^\eta) \in D^{r+1}|_{(\lambda, \mu)}$ for $1 \leq \gamma \leq g$, $1 \leq \eta \leq h$.

(2) If $\text{rank}C'(\lambda, \mu) = \text{rank}C(\lambda, \mu) + 1$ in the domain D^{r+1} , then

$$\text{rank}\tilde{C}'(W_{\lambda^\gamma}, W_{\mu^\eta}) = \text{rank}\tilde{C}(W_{\lambda^\gamma}, W_{\mu^\eta}) + m_\gamma \times n_\eta,$$

when $(\lambda^\gamma, \mu^\eta) \in D^{r+1}|_{(\lambda, \mu)}$. Moreover

$$\text{rank}\tilde{C}'(W(\underline{\lambda}^0), W(\underline{\mu}^0)) = \text{rank}\tilde{C}(W(\underline{\lambda}^0), W(\underline{\mu}^0)) + m_{\mathcal{P}_r} m_{\mathcal{Q}_r},$$

when $(\lambda^\gamma, \mu^\eta) \in D^{r+1}|_{(\lambda, \mu)}$ for $1 \leq \gamma \leq g$, $1 \leq \eta \leq h$.

Proof. (1) It is not difficult to see that

$$\text{rank}\tilde{C}(W_{\lambda^\gamma}, W_{\mu^\eta}) = m_\gamma n_\eta \text{rank}(C(\lambda, \mu)), \text{ and } \text{rank}\tilde{C}'(W_{\lambda^\gamma}, W_{\mu^\eta}) = m_\gamma n_\eta \text{rank}(C'(\lambda, \mu)).$$

In fact, the the left hand sides in two equalities have the nice expressions above, which imply that “ $\geq$ ” hold in both formulae. On the other hand, if some row of $C(\lambda, \mu)$ is a linear combination of other rows, then the corresponding block row of $\tilde{C}(W_{\lambda^\gamma}, W_{\mu^\eta})$ is the same linear combination of the same block rows. The similar assertion is true for C' and $\tilde{C}'$. Therefor “ $\leq$ ” hold in both formulae.

(2) The reason is: $\sum_{(\gamma,\eta)} m_\gamma n_\eta = m_{\mathcal{P}_r} n_{\mathcal{Q}_r}$. □

Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem with a reduction sequence $(*)$ of freely parameterized triples given in the Remark of 4.2. Suppose that H^s in the end term of $(**)$ possesses a set of parameters $\lambda_{\mathcal{I}_1}, \lambda_{\mathcal{I}_2}, \dots, \lambda_{\mathcal{I}_i}$. Let a set of valuation matrices

$$(\text{Vm}) : \quad \{W(\underline{\lambda}_{\mathcal{I}_1}^0), W(\underline{\lambda}_{\mathcal{I}_2}^0), \dots, W(\underline{\lambda}_{\mathcal{I}_i}^0)\}$$

be given by item (1) of Definition 4.4.1, and let $H_{\underline{m}^s}^s, \tilde{H}_{\underline{m}^s}^s(\underline{\lambda}^0)$ be given in the items (2), (3) of Definition 4.4.1 respectively. Assume that $(\hat{\underline{m}}, \hat{\underline{m}}^1, \dots, \hat{\underline{m}}^s)$ is a sequence of size vectors

calculated by Formula (11) of 3.2 inductively from $\hat{m}^s = \underline{m}^s$, and $(H_{\hat{m}}, H_{\hat{m}^1}^1, \dots, H_{\hat{m}^s}^s)$ is a sequence of matrices given by $(H, H^1, \dots, H^s)$ and $(\hat{m}, \hat{m}^1, \dots, \hat{m}^s)$. Then we are able to construct $\tilde{m}^r$ and $\tilde{H}_{\tilde{m}^r}^r(\Delta^0(r))$ according to the valuation matrices (Vm) for $r = s, s-1, \dots, 1, 0$, by Definition 4.4.1.

Thus we have a sequence of triples $(\hat{*})$ given in Corollary 4.3.1 (but the restriction of $m_{\mathcal{I}} = 1$ or 0 for any non-trivial $\mathcal{I}$ is relaxed). And we have a sequence of valuation matrices

$$0 = \tilde{H}_{\tilde{m}}(\Delta^0(0)), \tilde{H}_{\tilde{m}^1}^1(\Delta^0(1)), \dots, \tilde{H}_{\tilde{m}^r}^r(\Delta^0(r)), \dots, \tilde{H}_{\tilde{m}^{s-1}}^{s-1}(\Delta^0(s-1)), \tilde{H}_{\tilde{m}^s}^s(\Delta^0(s)),$$

and a sequence of triples

$$(\tilde{*}) \quad (\tilde{N}, \tilde{R}, \tilde{m}), (\tilde{N}^1, \tilde{R}^1, \tilde{m}^1), \dots, (\tilde{N}^r, \tilde{R}^r, \tilde{m}^r), \dots, (\tilde{N}^{s-1}, \tilde{R}^{s-1}, \tilde{m}^{s-1}), (\tilde{N}^s, \tilde{R}^s, \tilde{m}^s),$$

such that each triple $(\tilde{N}^r, \tilde{R}^r, \tilde{m}^r)$ is obtained from $(\hat{N}^r, \hat{R}^r, \hat{m}^r)$ by Definition 4.4.1.

Theorem 4.5.1 The sequence $(\tilde{*})$ is a reduction sequence. And either the type of reductions from $(\tilde{N}^r, \tilde{R}^r, \tilde{m}^r)$ to $(\tilde{N}^{r+1}, \tilde{R}^{r+1}, \tilde{m}^{r+1})$ is the same as that from (N^r, R^r, m^r) to $(N^{r+1}, R^{r+1}, m^{r+1})$, or the triple $(\tilde{N}^{r+1}, \tilde{R}^{r+1}, \tilde{m}^{r+1})$ is obtained from $(\tilde{N}^r, \tilde{R}^r, \tilde{m}^r)$ by identity, $0 \leq r \leq s-1$.

(We stress that the reductions in $(\tilde{*})$ are ordered according to the matrix order in $(\hat{*})$, which differs from the usual order of Definition 2.3.1.)

Proof. For $r = 0$, we set $\tilde{R} = \hat{R}$, $\tilde{N} = \hat{N}$, $\tilde{m} = \hat{m}$. Suppose we have already a sequence of reductions up to $(\tilde{N}^r, \tilde{R}^r, \tilde{m}^r)$, now consider the $(r+1)$ -th reduction towards $(\tilde{N}^{r+1}, \tilde{R}^{r+1}, \tilde{m}^{r+1})$. If $(\hat{N}^{r+1}, \hat{R}^{r+1}, \hat{m}^{r+1})$ is obtained from $(\hat{N}^r, \hat{R}^r, \hat{m}^r)$ by identity, so is $(\tilde{N}^{r+1}, \tilde{R}^{r+1}, \tilde{m}^{r+1})$ from $(\tilde{N}^r, \tilde{R}^r, \tilde{m}^r)$, we have done. Otherwise, let us go back to Proposition 4.3.1, and let $a_1 : \mathcal{P}_r \rightarrow \mathcal{Q}_r$ be the first arrow of $(\mathcal{K}^r, \mathcal{M}^r, H^r)$.

In item (1) of A1,A2,A3 (as well as (1') of A2) of Proposition 4.3.1, $\delta(a_1) = 0$, which means that the equation system (II) implies (8). Then item (1) of Proposition 4.5.1 tells that $(\tilde{\text{II}})$ implies $(\tilde{8})$. Thus we are able to set

$$\overline{\tilde{N}}_{p_r q_r}^r = \overline{\tilde{N}}_{p_r q_r}^r$$

by an edge reduction, or a loop reduction with fixed eigenvalues.

In item (2) of A1,A2,A3 (as well as (2') of A2) of Proposition 4.3.1, $\delta(a_1) \neq 0$, which means that the equation system (II) does not imply (8). Then item (2) of Proposition 4.5.1 tells that

$$\text{rank} \tilde{C}'(W_{\Delta^0}, W_{\mu^0}) = \text{rank} \tilde{C}(W(\Delta^0), W(\mu^0)) + m_{\mathcal{P}_r} n_{\mathcal{Q}_r}.$$

Thus we are able to set

$$\overline{\widetilde{N}}_{p_r q_r}^r = \emptyset = \overline{\widetilde{N}}_{p_r q_r}^r$$

by regularization.

In item (3) of A3 of Proposition 4.3.1, $\delta(a_1) = 0$. We are able to set

$$\overline{\widetilde{N}}_{p_r q_r}^r = \hat{W} \oplus W(\underline{\lambda}_{\mathcal{I}}^0),$$

whenever $\overline{\widetilde{N}}_{p_r q_r}^r = \hat{W} \oplus \lambda I_{m_{\mathcal{I}}}$ by a loop reduction, where $\mathcal{I} \in T^{r+1} / \sim^{r+1}$.

Our theorem follows by induction. $\square$

4.6 Valuation representation categories

Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem given by Definition 4.2.1, and $\mathfrak{A} = (\Gamma, \Omega)$ be the layered bocs corresponding to $(\mathcal{K}, \mathcal{M}, H)$. The present subsection is devoted to showing a valuation representation category $\widetilde{Mat}(\mathcal{K}, \mathcal{M})$, and equivalently $\widetilde{R}(\mathfrak{A})$.

Given any size vector $\underline{m}$ and any valuation matrix $\widetilde{H}_{\underline{m}}(\underline{\lambda}^0)$ defined in 4.4.1, an object of size vector $\underline{m}$ in $\widetilde{Mat}(\mathcal{K}, \mathcal{M})$ is defined as a matrix $M \in \mathcal{M}_{\underline{m}}$ with an *adjoint valuation matrix* $\widetilde{H}_{\underline{m}}(\underline{\lambda}^0)$. If L is also a matrix of size $\underline{l}$ with an adjoint valuation matrix $\widetilde{H}_{\underline{l}}(\underline{\mu}^0)$, then a morphism $\varphi : M \rightarrow L$ in $\widetilde{Mat}(\mathcal{K}, \mathcal{M})$ is a matrix $\varphi \in \mathcal{K}_{\underline{m} \times \underline{l}}$, such that

$$(M + \widetilde{H}_{\underline{m}}(\underline{\lambda}^0))\varphi = \varphi(L + \widetilde{H}_{\underline{l}}(\underline{\mu}^0)).$$

Thus we obtain a category of representations, which is called a *valuation representation category* of $(\mathcal{K}, \mathcal{M}, H)$, and we denote it by $\widetilde{Mat}(\mathcal{K}, \mathcal{M}, H)$.

Representation category $\widetilde{R}(\mathfrak{A})$ has been well-defined in [D] and [CB1]. The objects of $\widetilde{R}(\mathfrak{A})$ are Γ -modules. In fact, if $M \in \widetilde{Mat}(\mathcal{K}, \mathcal{M})$, an object $\overline{M} \in \widetilde{R}(\mathfrak{A})$ corresponding to M is defined by a set of vector spaces: $\overline{M}_{\mathcal{I}} = k^{m_{\mathcal{I}}}$, $\forall \mathcal{I} \in T / \sim$; a set of linear maps: $\overline{M}(\lambda_{\mathcal{I}}) = W(\underline{\lambda}_{\mathcal{I}}^0)$ for $\lambda_{\mathcal{I}}$ attached to $\mathcal{I}$, and $\overline{M}(\rho_{\mathcal{I}\mathcal{J}}^w)^* = M_{p_{\mathcal{I}\mathcal{J}}^w q_{\mathcal{I}\mathcal{J}}^w}$ for any basis element $\rho_{\mathcal{I}\mathcal{J}}^w \in A$ (see Formula (1) in 2.3). Thus $\overline{M}$ is a Γ -module. If $\overline{L} \in \widetilde{R}(\mathfrak{A})$ corresponds to $L \in \widetilde{Mat}(\mathcal{K}, \mathcal{M})$, and $\varphi (= S) : M \rightarrow L$ is a morphism in $\widetilde{Mat}(\mathcal{K}, \mathcal{M})$, a morphism $\overline{\varphi} : \overline{M} \rightarrow \overline{L}$ corresponding to φ in $\widetilde{R}(\mathfrak{A})$ is defined by a set of linear maps: $(\varphi_{\mathcal{I}}, \varphi(\zeta_{\mathcal{I}\mathcal{J}}^w)^* \mid \forall \mathcal{I} \in T / \sim, \forall \zeta_{\mathcal{I}\mathcal{J}}^w \in B)$, where $\varphi(\zeta_{\mathcal{I}\mathcal{J}}^w)^* = S_{\overline{p}_{\mathcal{I}\mathcal{J}}^w \overline{q}_{\mathcal{I}\mathcal{J}}^w}$. It is clear that $\varphi_{\mathcal{I}} : \overline{M}_{\mathcal{I}} \rightarrow \overline{L}_{\mathcal{I}}$ satisfy

$$\overline{M}(\rho_{\mathcal{I}\mathcal{J}}^w)^* \varphi_{\mathcal{J}} - \varphi_{\mathcal{I}} \overline{L}(\rho_{\mathcal{I}\mathcal{J}}^w)^* = \varphi(\delta(\rho_{\mathcal{I}\mathcal{J}}^w)^*)$$

(see the last part of 3.6). If $\overline{\psi} : \overline{L} \rightarrow \overline{E}$ is also a morphism, then the composition $\overline{\eta} = \overline{\varphi} \overline{\psi} : \overline{L} \rightarrow \overline{E}$ is given by $\overline{\eta} = (\eta_{\mathcal{I}}, \eta(\zeta_{\mathcal{I}\mathcal{J}}^w)^* \mid \forall \mathcal{I} \in T / \sim, \forall \zeta_{\mathcal{I}\mathcal{J}}^w \in B)$, such that

$\eta_{\mathcal{I}} = \varphi_{\mathcal{I}}\psi_{\mathcal{I}}$, and

$$\eta(\zeta_{\mathcal{I}\mathcal{J}}^w)^* = \varphi(\zeta_{\mathcal{I}\mathcal{J}}^w)^*\psi_{\mathcal{J}} + \varphi_{\mathcal{I}}\psi(\zeta_{\mathcal{I}\mathcal{J}}^w)^* + \sum_{\mathcal{L}, u, v} c_{\binom{\mathcal{I}\mathcal{J}}{\mathcal{I}\mathcal{L}}}^{\binom{w}{u}} \varphi(\zeta_{\mathcal{I}\mathcal{L}}^u)^*\psi(\zeta_{\mathcal{I}\mathcal{L}}^v)^*$$

(see also the last part of 3.6).

Now suppose the freely parameterized bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ (respectively $\mathfrak{A}$) is minimal. Given any non-trivial $\mathcal{I} \in T/\sim$ with an attached parameter $\lambda_{\mathcal{I}}$ of domain $k \setminus \{\text{the roots of } g_{\mathcal{I}}(\lambda_{\mathcal{I}})\}$, let $\underline{m}(\mathcal{I}, d)$ be a size vector of $(T, \sim)$, such that $m(\mathcal{I}, d)_{\mathcal{I}} = d$ for some $d \in \mathbb{N}$ and $m(\mathcal{I}, d)_{\mathcal{L}} = 0, \forall \mathcal{L} \in T/\sim \setminus \{\mathcal{I}\}$. For any $\lambda_{\mathcal{I}}^0 \in k$ with $g_{\mathcal{I}}(\lambda_{\mathcal{I}}^0) \neq 0$, we define a zero matrix $O(\mathcal{I}, d, \lambda_{\mathcal{I}}^0) \in \widetilde{\mathcal{M}}_{\underline{m}(\mathcal{I}, d)}$ with an adjoint valuation matrix $\widetilde{H}_{\underline{m}(\mathcal{I}, d)}(J_d(\lambda_{\mathcal{I}}^0))$, and we denote it by $\widetilde{H}(J_d(\lambda_{\mathcal{I}}^0))$ for simplicity.

Corollary 4.6.1 (1) $O(\mathcal{I}, d, \lambda_{\mathcal{I}}^0)$ is indecomposable.

(2) $\{O(\mathcal{J}) \mid \forall \text{ trivial } \mathcal{J} \in T/\sim\}$ and

$$\{O(\mathcal{I}, d, \lambda_{\mathcal{I}}^0) \mid \forall \text{ non-trivial } \mathcal{I} \in T/\sim, \forall d \in \mathbb{N}, \forall \lambda_{\mathcal{I}}^0 \in k \text{ with } g_{\mathcal{I}}(\lambda_{\mathcal{I}}^0) \neq 0\}$$

form a complete set of indecomposables of $\widetilde{Mat}(\mathcal{K}, \mathcal{M})$.

Proof. (1) Let $O(\mathcal{I}, d, \lambda_{\mathcal{I}}^0)$ be an object of $\widetilde{Mat}(\mathcal{K}, \mathcal{M}, H)$ with an adjoint matrix $\widetilde{H}(J_d(\lambda_{\mathcal{I}}^0))$. We denote $O(\mathcal{I}, d, \lambda_{\mathcal{I}}^0)$ by O for simplicity. Suppose $\overline{O} \in \widetilde{R}(\mathfrak{A})$ corresponds to O . Thus $\overline{O}_{\mathcal{I}} = k^d$, $\overline{O}_{\mathcal{J}} = 0$, and $\overline{O}(\lambda_{\mathcal{I}}) = J_d(\lambda_{\mathcal{I}}^0)$. Given any morphism $\overline{\varphi} : \overline{O} \rightarrow \overline{O}$, then

$$\overline{\varphi}_{\mathcal{I}} = \begin{pmatrix} x_1 & x_2 & \cdots & x_d \\ & x_1 & \cdots & x_{d-1} \\ & & \ddots & \vdots \\ & & & x_1 \end{pmatrix} \quad (35)$$

is given by $J_d(\lambda_{\mathcal{I}}^0)\overline{\varphi}_{\mathcal{I}} = \overline{\varphi}_{\mathcal{I}}J_d(\lambda_{\mathcal{I}}^0)$, and $\overline{\varphi}_{\mathcal{J}} = 0$. Therefore $End_{\widetilde{R}(\mathfrak{A}^{\infty})}(\overline{O})$ is local. Consequently, $\overline{O}$ is indecomposable, so is O .

(2) See [CB1, 6.2]. □

From now on we will not worry about any difference between $\widetilde{Mat}(\mathcal{K}, \mathcal{M})$ and $\widetilde{R}(\mathfrak{A})$. And we denote them still by $Mat(\mathcal{K}, \mathcal{M})$ and $R(\mathfrak{A})$ respectively for simplicity.

If $(**)$ is a reduction sequence of freely parameterized bimodule problems given in 3.3, the *reduction functors*

$$\vartheta_{r-1, r} : Mat(\mathcal{K}^r, \mathcal{M}^r) \rightarrow Mat(\mathcal{K}^{r-1}, \mathcal{M}^{r-1})$$

are defined as follows. Given any matrix $M^r \in \mathcal{M}_{\underline{m}}^r$ with an adjoint valuation matrix $\widetilde{H}_{\underline{m}}^r(\underline{\Delta}^0(r))$, then

$$\vartheta_{r-1, r}(M^r) = M^r + \widetilde{H}_{\underline{m}}^r(\underline{\Delta}^0(r))_{p_{r-1}q_{r-1}} \otimes \rho^{r-1}$$

with an adjoint valuation matrix $\tilde{H}_{\underline{m}}^r(\underline{\lambda}^0(r))$, where ρ^{r-1} is the first basis element of $\mathcal{M}^{r-1}$. Moreover, the composition

$$\vartheta_{0r} = \vartheta_{01} \cdots \vartheta_{r-1,r} : \text{Mat}(\mathcal{K}^r, \mathcal{M}^r) \rightarrow \text{Mat}(\mathcal{K}, \mathcal{M})$$


is obtained by sending M^r to $M^r + \tilde{H}_{\underline{m}}^r(\underline{\lambda}^0(r))$. And the action of the reduction functors on morphisms is an identity as the same as that in 3.3. In particular if the end term of $(**)$ is minimal, then

$$\vartheta_{0,\infty}(O(\mathcal{I}, d, \lambda_{\mathcal{I}}^0)) = \tilde{H}^\infty(J_d(\lambda_{\mathcal{I}}^0)),$$

which is indecomposable by Corollary 4.6.1.

Finally we claim why the conditions (2) of Definition 4.2.1 must hold for a freely parameterized bimodule problem by some examples to end the subsection.

Example 1. Let $\mathfrak{A} = (\Gamma, \Omega)$ be a local bocs with a layer $L = (\Gamma'; \omega; a; v)$



where $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda]$, $\delta(a) = v\lambda - \lambda v$, i.e. $f(\lambda, \mu) = -\lambda + \mu$. If we require $f(\lambda, \mu) \neq 0$, and set $M_{\mathcal{P}} = k$, $M(\lambda) = (\lambda^0)$, $\forall \lambda^0 \in k$, then there is not any morphism $\varphi : M \rightarrow M$. In fact, if φ were a morphism, then $M(a)\varphi_{\mathcal{P}} - \varphi_{\mathcal{P}}M(a) = v\lambda^0 - \lambda^0v = 0$, which would lead to a contradiction to the requirement of $f(\lambda^0, \lambda^0) \neq 0$.

Example 2. We have the same picture as Example 1, but $\delta(a) = 2v\lambda - \lambda v$, i.e. $f(\lambda, \mu) = -\lambda + 2\mu$. If we require $f(\lambda, \mu) = 0$, and set $M_{\mathcal{P}} = k$, $M(\lambda) = (\lambda^0)$, $\forall \lambda^0 \in k$, and $L_{\mathcal{P}} = k$, $L(\lambda) = (\mu^0)$, then there is not any morphism $\varphi : M \rightarrow L$ when $\lambda^0 \neq 2\mu^0$. In fact, if φ were a morphism, then $M(a)\varphi_{\mathcal{P}} - \varphi_{\mathcal{P}}L(a) = 2v\lambda^0 - \mu^0v \neq 0$, which would lead to a contradiction to the requirement of $f(\lambda^0, \mu^0) = 0$.

5 Minimally wild bimodule problems

This section is devoted to classifying the minimally wild bimodule problems.

5.1 The wild theorem

We will give an alternative proof of the well-known Drozd's wild theorem in this subsection. Our proof may not be an essential improvement, since the idea of the original proof is perfect. But the new proof lowers the dimension from the original 43 to 20. This makes the matrices simpler.

Definition 5.1.1([CB1]) We say a layered boc $\mathfrak{A}$ is *wild*, if there is a functor

$$F : k\langle x, y \rangle\text{-mod} \rightarrow R(\mathfrak{A})$$

which preserves iso-classes and indecomposability.

Theorem 5.1.1([D], [CB1]) Let $\mathfrak{A} = (\Gamma, \Omega)$ be a boc with layer $L = (\Gamma'; \omega; a_1, \dots, a_n; v_1, \dots, v_m)$. Suppose that $a_1 : \mathcal{P} \rightarrow \mathcal{Q}$. Then $\mathfrak{A}$ must be wild in the following two cases:

Case 1. $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda, g_{\mathcal{P}}(\lambda)^{-1}]$ and $\Gamma'(\mathcal{Q}, \mathcal{Q}) = k[\mu, g_{\mathcal{Q}}(\mu)^{-1}]$;


$$\delta(a_1) = f(\lambda, \mu)v_1$$

where $f(\lambda, \mu) \in k[\lambda, \mu, g_{\mathcal{P}}(\lambda)^{-1}, g_{\mathcal{Q}}(\mu)^{-1}]$ is non-invertible.

Case 2. $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda, g_{\mathcal{P}}(\lambda)^{-1}]$ and $\Gamma'(\mathcal{Q}, \mathcal{Q}) = k$;

$$\delta(a_1) = 0, \quad \text{or dual.}$$

Proof. Case 1. Without loss of generality we may assume that $\mathfrak{A}$ is local when $\mathcal{P} = \mathcal{Q}$; or $\mathfrak{A}$ has two vertices when $\mathcal{P} \neq \mathcal{Q}$:



We may also assume that $f(\lambda, \mu) \in k[\lambda, \mu]$ after dividing v_1 by some power of $g_{\mathcal{P}}(\lambda)g_{\mathcal{Q}}(\mu)$, write

$$f(\lambda, \mu) = \alpha(\lambda - \lambda^0) + \beta(\mu - \mu^0) + \gamma_1(\lambda - \lambda^0)^2 + \gamma_2(\lambda - \lambda^0)(\mu - \mu^0) + \gamma_3(\mu - \mu^0)^2 + \dots \quad (36)$$

where $f(\lambda^0, \mu^0) = 0$, $g_{\mathcal{P}}(\lambda^0) \neq 0$, $g_{\mathcal{Q}}(\mu^0) \neq 0$. If $\mathcal{P} \neq \mathcal{Q}$, and $\alpha\beta \neq 0$, we set $m_{\mathcal{P}} = 9$, $m_{\mathcal{Q}} = 11$, and define an object $M \in R(\mathfrak{A})$, such that $M(\lambda) \simeq J_1(\lambda^0) \oplus J_3(\lambda^0) \oplus J_5(\lambda^0)$, $M(\mu) \simeq$

$J_2(\mu^0) \oplus J_4(\mu^0) \oplus J_5(\mu^0)$. Then

$$M(\lambda) = \begin{pmatrix} \boxed{\begin{matrix} \lambda^0 & 0 & 0 & 1 & 0 \\ & \lambda^0 & 0 & 0 & 1 \\ & & \lambda^0 & 0 & 0 \end{matrix}} & & & & \\ & \boxed{\begin{matrix} \lambda^0 & 0 & 1 & 0 \\ & \lambda^0 & 0 & 1 \end{matrix}} & & & \\ & & \boxed{\begin{matrix} \lambda^0 & 0 & 1 \\ & \lambda^0 & 0 \end{matrix}} & & \\ & & & \boxed{\begin{matrix} \lambda^0 & 1 \\ & \lambda^0 \end{matrix}} \end{pmatrix}$$

and

$$S_{\mathcal{P}} = \begin{pmatrix} \boxed{\begin{matrix} s_1 & s_{13}^1 & s_{15}^1 & s_{11}^2 & s_{13}^2 & s_{11}^3 & s_{13}^3 & s_{11}^4 & s_{11}^5 \\ & s_3 & s_{35}^1 & 0 & s_{33}^2 & s_{31}^3 & s_{33}^3 & s_{31}^4 & s_{31}^5 \\ & & s_5 & 0 & 0 & 0 & s_{53}^3 & 0 & s_{51}^5 \end{matrix}} & & & & \\ & \boxed{\begin{matrix} s_1 & s_{13}^1 & s_{11}^2 & s_{13}^2 & s_{11}^3 & s_{11}^4 \\ & s_3 & 0 & s_{33}^2 & s_{31}^3 & s_{31}^4 \end{matrix}} & & & \\ & & \boxed{\begin{matrix} s_1 & s_{13}^1 & s_{11}^2 & s_{11}^3 \\ & s_3 & 0 & s_{31}^3 \end{matrix}} & & \\ & & & \boxed{\begin{matrix} s_1 & s_{11}^2 \\ & s_{11}^3 \end{matrix}} \end{pmatrix}$$

is given by $M(\lambda)S_{\mathcal{P}} - S_{\mathcal{P}}M(\lambda) = 0$. On the other hand

$$M(\mu) = \begin{pmatrix} \boxed{\begin{matrix} \mu^0 & 0 & 0 & 1 & 0 & 0 \\ & \mu^0 & 0 & 0 & 1 & 0 \\ & & \mu^0 & 0 & 0 & 1 \end{matrix}} & & & & \\ & \boxed{\begin{matrix} \mu^0 & 0 & 0 & 1 & 0 \\ & \mu^0 & 0 & 0 & 1 \\ & & \mu^0 & 0 & 0 \end{matrix}} & & & \\ & & \boxed{\begin{matrix} \mu^0 & 0 & 1 & 0 \\ & \mu^0 & 0 & 1 \end{matrix}} & & \\ & & & \boxed{\begin{matrix} \mu^0 & 0 & 1 \\ & \mu^0 & 0 \end{matrix}} \end{pmatrix},$$

and

$$S_Q = \begin{pmatrix} \boxed{\begin{matrix} t_1 & t_{12}^1 & t_{14}^1 \\ & t_2 & t_{24}^1 \\ & & t_4 \end{matrix}} & \boxed{\begin{matrix} t_{11}^2 & t_{12}^2 & t_{14}^2 \\ t_{21}^2 & t_{22}^2 & t_{24}^2 \\ 0 & 0 & t_{44}^2 \end{matrix}} & * \\ & \boxed{\begin{matrix} t_1 & t_{12}^1 & t_{14}^1 \\ & t_2 & t_{24}^1 \\ & & t_4 \end{matrix}} & \\ & & * \end{pmatrix}.$$

is given by $M(\mu)S_Q - S_QM(\mu) = 0$.

The 9×11 matrix $S(\delta(a_1))$ equals

$$\begin{pmatrix} * & * & * & & & & & & & & \\ * & * & * & & & & & & & & \\ 0 & 0 & 0 & & & & & & * & & \\ * & * & * & & & & & & & & \\ \alpha v_{71} & \alpha v_{72} & * & & & & & & & & \\ \alpha v_{81} + \gamma_1 v_{91} & \alpha v_{82} + \gamma_1 v_{92} & * & & & & & & & & \\ 0 & 0 & 0 & \beta v_{71} & \beta v_{72} & * & & & & & \\ \alpha v_{91} & \alpha v_{92} & \alpha v_{93} & \alpha v_{94} + \beta v_{81} + \gamma_2 v_{91} & \alpha v_{95} + \beta v_{82} + \gamma_2 v_{92} & * & & & & & \\ 0 & 0 & 0 & \beta v_{91} & \beta v_{92} & \beta v_{93} & \beta v_{94} + \gamma_3 v_{91} & \beta v_{95} + \gamma_3 v_{92} & * & \dots & \end{pmatrix}$$

Let

$$M(a_1) = \begin{pmatrix} \emptyset & \emptyset & \emptyset & & & & & & & & \\ \emptyset & \emptyset & \emptyset & & & & & & & & \\ 1 & \xi & \emptyset & & & & & & & & \\ \emptyset & \emptyset & \emptyset & & & & & & & & \\ \eta & 1 & \emptyset & & & & & & & & \\ 1 & \emptyset & \emptyset & & & & & & & & \\ 0 & 0 & 1 & & & & & & & & \\ 0 & 0 & 1 & & & & & & & & \\ 0 & 0 & 0 & & & & & & & & \end{pmatrix}_{9 \times 11}$$

where η and ξ are two algebraically independent parameters with a domain $k \times k$. Thus $\forall S \in \text{End}_{\mathfrak{A}}(M)$, we have $S_{\mathcal{P}} = sI_9 + L_1$ and $S_Q = sI_{11} + L_2$ for some upper triangular nilpotent matrices L_1, L_2 , and $s \in k$.

If $\alpha\beta = 0$, we can take some lower sizes to construct a canonical matrix with two independent parameters η, ξ , [ZX].

If $\mathcal{P} = \mathcal{Q}$, and there exists a pair (λ^0, μ^0) with $\lambda^0 = \mu^0$, $f(\lambda^0, \mu^0) = 0$, (i.e. λ^0 is a root of $f(\lambda, \lambda)$), $g_{\mathcal{P}}(\lambda^0) \neq 0$, we may set $m_{\mathcal{P}} = 4$, $M(\lambda) \simeq J_3(\lambda^0) \oplus J_1(\lambda^0)$. Thus

$$M(\lambda) = \begin{pmatrix} \boxed{\begin{matrix} \lambda^0 & 0 & 1 \\ & \lambda^0 & 0 \\ & & \lambda^0 & 1 \\ & & & \lambda^0 \end{matrix}} \end{pmatrix}, \quad \text{and} \quad S_{\mathcal{P}} = \begin{pmatrix} \boxed{\begin{matrix} s_1 & s_{13}^1 & s_{11}^2 & s_{11}^3 \\ & s_3 & 0 & s_{31}^3 \\ & & s_1 & s_{11}^2 \\ & & & s_1 \end{matrix}} \end{pmatrix}$$

is given by $M(\lambda)S_{\mathcal{P}} = S_{\mathcal{P}}M(\lambda)$. When $\alpha\beta \neq 0$,

$$S(\delta(a_1)) = \begin{pmatrix} * & * & * & * \\ 0 & * & * & * \\ \alpha v_{41} & * & * & * \\ 0 & 0 & \beta v_{41} & * \end{pmatrix}, \quad M(a_1) = \begin{pmatrix} \emptyset & \emptyset & \emptyset & \emptyset \\ \xi & \emptyset & \emptyset & \emptyset \\ \eta & \emptyset & \emptyset & \emptyset \\ 0 & 1 & \emptyset & \emptyset \end{pmatrix},$$

where η and ξ are independent parameters with a domain $k \times k$. Thus given any $S \in \text{End}_{\mathfrak{A}}(M)$, $S_{\mathcal{P}} = sI_4 + L$ for some upper triangular nilpotent matrix L and $s \in k$. When $\alpha\beta = 0$ or $\alpha \neq 0, \beta = 0$, the method is similar [ZX].

If $\mathcal{P} = \mathcal{Q}$, and all the pairs (λ^0, μ^0) with $f(\lambda^0, \mu^0) = 0$, $g_{\mathcal{P}}(\lambda^0)g_{\mathcal{Q}}(\mu^0) \neq 0$ have the property that $\lambda^0 \neq \mu^0$, we may use a rolled up version $\overline{M}$ of the construction given in the case $\mathcal{P} \neq \mathcal{Q}$, i.e. let $m_{\mathcal{P}} = 20$, $\overline{M}(\lambda) = M(\lambda^0) \oplus M(\mu^0)$ and $\overline{M}(a_1) = \begin{pmatrix} 0 & M(a_1) \\ 0 & 0 \end{pmatrix}$.

Case 2. $\lambda \overset{\mathcal{P}}{\curvearrowright} \overset{a_1}{\bullet} \mathcal{Q}$, (or we have a dual diagram). Let $m_{\mathcal{P}} = 5$, $m_{\mathcal{Q}} = 2$, $M(\lambda) = J_5(\lambda^0)$, where $g_{\mathcal{P}}(\lambda^0) \neq 0$, then

$$S_{\mathcal{P}} = \begin{pmatrix} s_1 & s_2 & s_3 & s_4 & s_5 \\ & s_1 & s_2 & s_3 & s_4 \\ & & s_1 & s_2 & s_3 \\ & & & s_1 & s_2 \\ & & & & s_1 \end{pmatrix}, \quad S_{\mathcal{Q}} = \begin{pmatrix} t_{11} & t_{12} \\ t_{21} & t_{22} \end{pmatrix}; \quad M(a_1) = \begin{pmatrix} \xi & \emptyset \\ \eta & \emptyset \\ \emptyset & \emptyset \\ 1 & \emptyset \\ 0 & 1 \end{pmatrix},$$

where η and ξ are independent parameters with a domain $k \times k$. Thus $\forall S \in \text{End}_{\mathfrak{A}}(M)$, $S_{\mathcal{P}} = sI_5$, $S_{\mathcal{Q}} = sI_2$, for any $s \in k$. The proof for the dual case is similar.

It is clear from the construction of M , that we are able to define a local layered boc $\mathfrak{B}$ induced from $\mathfrak{A}$ just before the appearing of ξ in all the cases (see also [ZX]), i.e. $\mathfrak{B}$ has a layer $L_{\mathfrak{B}} = (\Gamma'_{\mathfrak{B}}; \omega_{\mathfrak{B}}; b_0, b_1, \dots, b_n; u_1, \dots, u_m)$ with a unique non-trivial vertex $\mathcal{I}$, and $\Gamma'_{\mathfrak{B}}(\mathcal{I}, \mathcal{I}) = k[\eta]$. Moreover, $\delta(b_0) = 0$ still by the construction, and we are able to set $b_0 = \xi$. We denote the reduction functor from $R(\mathfrak{B})$ to $R(\mathfrak{A})$ by G . And define a functor

$$F : k\langle \eta, \xi \rangle\text{-mod} \rightarrow R(\mathfrak{B})$$

by sending any $k\langle\eta, \xi\rangle$ -module

$$L = \begin{array}{ccc} L(\eta) & \xrightarrow{k^m} & L(\xi) \\ \curvearrowright & \bullet & \curvearrowleft \end{array}$$

for any dimension m and any linear maps $L(\eta), L(\xi)$ to $\overline{L} = F(L)$, such that

$$\overline{L}_{\mathcal{P}} = k^m, \overline{L}(\eta) = L(\eta), \overline{L}(b_0) = L(\xi), \overline{L}(b_i) = 0 \text{ for } i = 1, 2, \dots, n.$$

If $L' \in k\langle\eta, \xi\rangle\text{-mod}$ and $\overline{L}' = F(L')$, then a morphism $\varphi : L \rightarrow L'$ is sent to a morphism $\overline{\varphi} : \overline{L} \rightarrow \overline{L}'$, such that $\overline{\varphi}_{\mathcal{P}} = \varphi$, and $\overline{\varphi}(u_j) = 0$ for $j = 1, 2, \dots, m$. Thus, if L is indecomposable, so is $\overline{L}$; and if $\overline{L} \simeq \overline{L}'$, then $L \simeq L'$ both by Corollary 2.2.2. Finally, the functor $GF : k\langle\eta, \xi\rangle\text{-mod} \rightarrow R(\mathfrak{A})$ preserves indecomposability and isomorphism classes, so that $\mathfrak{A}$ is of wild type. $\square$

5.2 The tame theorem

In this subsection we will give a slightly different statement of the well-known tame theorem [D, CB1] by constructing a finite set of minimal local bimodule problems whose valued representation categories cover all the canonical forms of the indecomposable matrices of dimension at most n for each non-negative integer n .

Lemma 5.2.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a non-wild bimodule problem having a reduction sequence $(*)$. Suppose that $\overline{N}_{pr_0, qr_0}^{r_0} = W \oplus (\lambda)$ with λ attached to $\mathcal{I} \in T^{r_0+1} / \sim^{r_0+1}$. Given any $r > r_0$, if the first arrow a_1 in $(\mathcal{K}^r, \mathcal{M}^r, H^r)$ starts or ends at $\mathcal{I}$, then the reduction for a_1 must be a regularization. Consequently, the equivalent class $\mathcal{I}$ remains until the end of the sequence $(*)$, i.e. $\mathcal{I} \in T^\infty / \sim^\infty$. In particular, if R^∞ is local, then $\overline{N}_{pr_0, qr_0}^{r_0} = (\lambda)$.

Proof. Item (1) of A1 and A2 in Proposition 4.3.1 can not occur because of non-wildness, and item (2) of A1, A2 always yields a regularization. $\square$

Lemma 5.2.2 Let $(\mathcal{K}, \mathcal{M}, H)$ be a non-wild bimodule problem. Then any reduction sequence $(*)$ of freely parameterized triples, which preserves all the free parameters, can reach a minimal triple.

Proof. By the same reason as in the proof of Lemma 5.2.1. $\square$

Lemma 5.2.3 Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a non-wild bimodule problem having a reduction sequence $(*)$ and $N^s = (0)$. Then $H^\infty(\mathcal{J})$ for any trivial $\mathcal{J} \in T^\infty / \sim^\infty$ defined in 3.5, and $\tilde{H}^\infty(J_d(\lambda_{\mathcal{I}}^0))$ for any non-trivial $\mathcal{I} \in T^\infty / \sim^\infty$ defined in 4.6 are both canonical forms under the fixed orders of matrix and the base field.

Proof. Let $\hat{n}^\infty = \underline{m}(\mathcal{J})$ when $\mathcal{J}$ is trivial. We obtain a reduction sequence $(\hat{*})$ in Corollary 3.3.1 under the orders mentioned in the lemma. Thus $H^\infty(\mathcal{J})$ is a canonical form.

Let $\hat{n}^\infty = \underline{m}(\mathcal{I}, 1)$ when $\mathcal{I}$ is non-trivial, which has dimension 1 at $\mathcal{I}$ and 0 at any $\mathcal{J} \neq \mathcal{I}$. We obtain a reduction sequence $(\hat{*})$ in Corollary 4.3.1, which contains a unique parameter $\lambda_{\mathcal{I}}$ and $\hat{R}^\infty$ is local. Thus $\overline{N}_{p_r q_r}^r = (\lambda_{\mathcal{I}})$ for some $1 \leq r \leq s$ by Lemma 5.2.1. It is obvious that the order of the base field is not destroyed in $\tilde{H}^{r+1}(J_d(\lambda_{\mathcal{I}}^0))$. On the other hand, all the reductions after the $(r+1)$ -th step are regularization in $(\hat{*})$ given by $J_d(\lambda_{\mathcal{I}}^0)$ by Lemma 5.2.1 and Theorem 4.5.1, which are made according to the partitions of $(\hat{*})$. But they can also be made according to the order of Definition 2.3.1, whenever we write the matrix equations (8) of 2.6 element-wise. Therefore $\tilde{H}^\infty(J_d(\lambda_{\mathcal{I}}^0))$ is a canonical form under the fixed orders of both matrix and base field. $\square$

Theorem 5.2.1 Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a non-wild bimodule problem. Suppose we fix the orders both on matrix and the base field. Then for any fixed $n \in \mathbb{N}$, there exists a finite set of sequences of freely parameterized triples $\{*_1, *_2, \dots, *_h\}$ such that

(1) the end terms of corresponding $(**)$'s:

$$(\mathcal{K}_1^\infty, \mathcal{M}_1^\infty, H_1^\infty), (\mathcal{K}_2^\infty, \mathcal{M}_2^\infty, H_2^\infty), \dots, (\mathcal{K}_h^\infty, \mathcal{M}_h^\infty, H_h^\infty)$$

are minimal and local;

(2) for any indecomposable $M \in \text{Mat}(\mathcal{K}, \mathcal{M})$ of size at most n , there exists a unique sequence $(*_l)$ in the set and a matrix

$$M_l^\infty \in \text{Mat}(\mathcal{K}_l^\infty, \mathcal{M}_l^\infty) \quad \text{with} \quad M \simeq \vartheta_{0,\infty}^l(M_l^\infty).$$

(3) $\vartheta_{0,\infty}^l(M_l^\infty)$ is the canonical form of M .

Proof. We use the induction firstly on size n . $n = 0$ is trivial. Suppose that the statement is true for a non-negative integer $(n-1)$. We will start from a triple $(N, R, \underline{n})$ of size n . We use the induction for the second time on the reduction steps r .

When $r = 0$, there are only finitely many choices of size vector $\underline{n}$, since the equation

$$n = n_1 + n_2 + \dots + n_t$$

has only finitely many solutions of non-negative integers. Suppose that for each fixed triple $(N, R, \underline{n})$, there are only finitely many possibilities to construct the reduction sequences up to the r -th step. Consider one of such sequences. We will show that the possibilities to construct $\overline{N}_{p_r q_r}^r$ are also finite. Thus our conclusion holds, since a reduction sequence towards a minimal bimodule problem can be taken at most n^2 steps.

We treat the $(r+1)$ -th reduction still according to Proposition 4.3.1. Item (1) of A1 and A2 can not occur because of non-wildness. Item (2) of A1, A2, A3 gives $\overline{N}_{p_r q_r}^r = \emptyset$, so we have only one choice.

In item (1) of A3 and $\mathcal{P} \neq \mathcal{Q}$,

$$\overline{N}_{p_r q_r}^r = \begin{pmatrix} 0 & I_d \\ 0 & 0 \end{pmatrix}_{n_{p_r}^r \times n_{q_r}^r},$$

then we have finitely many choices for $d = 0, 1, \dots, \min\{n_{p_r}^r, n_{q_r}^r\}$.

In item (1) of A3 and $\mathcal{P} = \mathcal{Q}$ or item (3) of A3. If R^r is not local, then either $T^r / \sim^r \setminus \{\mathcal{P}_r\} \neq \emptyset$, or $T^r / \sim^r = \{\mathcal{P}_r\}$, and $n_{p_r} > 1$. Let $(\hat{*})$ be a reduction sequence of Corollary 4.3.1 given by $\hat{n}^r$, such that $\hat{n}_{p_r}^r = 1$, $\hat{n}_{\mathcal{J}}^r = 0$, $\forall \mathcal{J} \in T^r / \sim^r$, $\mathcal{J} \neq \mathcal{P}_r$, then $\hat{n}^r < n^r$. The induction hypothesis and Lemma 5.2.1 tell us that the sequence $(\hat{*})$ with $\hat{N}_{p_r q_r}^r = (\lambda)$ already exists in our set and reaches a minimal triple with domain $k \setminus \{\text{the roots of } \hat{g}(\lambda)\}$. Thus Lemma 5.3.3 ensures that $\tilde{H}^\infty(J_d(\lambda^0))$, for some $d \in \mathbb{N}$ and $\hat{g}(\lambda^0) \neq 0$, have covered the canonical forms of indecomposables of size at most n such that $\hat{N}_{p_r q_r}^r$ has eigenvalues $\lambda^0 \in (k \setminus \{\text{the roots of } \hat{g}(\lambda)\})$. Therefore

$$\overline{N}_{p_r q_r}^r \simeq \oplus_{l,d} J_d(\lambda^l)^{e_{l,d}}$$

with $\hat{g}(\lambda^l) = 0$ such that

$$n_{p_r} = \sum_{l,d} d e_{l,d}.$$

But such a equation in variables $e_{l,d}$ has only finitely many solutions of non-negative integers, i.e. $\overline{N}_{p_r q_r}^r$ has only finitely many choices.

If R^r is local, then $T^r / \sim^r = \{\mathcal{P}_r\}$ and $n_{p_r} = 1$. Item (3) of A3 gives $\overline{N}_{p_r q_r}^r = (\lambda)$ with a domain $k \setminus \{\text{the roots of } g(\lambda)\}$, or $\overline{N}_{p_r q_r}^r = (\lambda^l)$ with $g(\lambda^l) = 0$, thus $\overline{N}_{p_r q_r}^r$ has only finitely many choices.

Therefore we obtain finitely many reduction sequences $(*)$ of size n . Finally all the sequences having local end terms are retained, and others are excluded, since their local direct summands have already been included in the case of $(n-1)$. The proposition follows by double induction on n and r . $\square$

5.3 Local wild bimodule problems

This subsection is devoted to constructing a reduction sequence consisting of local parameterized bimodule problems (not necessarily freely!). In the subsection we will not distinguish the multiplication of the parameters whether they act from left or right. And

we denote such kind of differentials by δ^c , where c in δ^c means the commutativity of the multiplication from left and right.

Lemma 5.3.1. Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem with $T/\sim = \{\mathcal{P}\}$, and H have a free parameter λ_0 with the domain $k \setminus \{\text{the roots of } f(\lambda_0)\}$. Suppose that $\underline{n}$ is a size vector with $n_{\mathcal{P}} = 1$ and $(N, R, \underline{n})$ is given by Formula (6) of 2.6. Then we are able to construct a reduction sequence $(*)$ of 3.3 satisfying the following properties:

- (1) $\underline{n} = \underline{n}^1 = \cdots = \underline{n}^r = \cdots = \underline{n}^s$, $N^s = (0)$. H^s contains parameters $\lambda_0, \lambda_1, \dots, \lambda_\gamma$ with $\gamma \geq 0$, which are algebraically independent.
- (2) When $\delta^c(\rho^{r_l-1})^* = 0$ in $(\mathcal{K}^{r_l-1}, \mathcal{M}^{r_l-1})$, we set $(H^{r_l})_{p_{r_l-1}, q_{r_l-1}} = (\lambda_l)$, where $l = 1, \dots, \gamma$.
- (3) When $\delta^c(\rho^{r-1})^* = w^{r-1} \neq 0$ in $(\mathcal{K}^{r-1}, \mathcal{M}^{r-1})$, we set $(H^r)_{p_{r-1}, q_{r-1}} = \emptyset$ and $w^{r-1} = 0$ after a localization given by $c^r(\lambda_0, \lambda_1, \dots, \lambda_l)$, where $r_l < r < r_{l+1}$, $l = 0, 1, \dots, \gamma$, $r_0 = 0$, $r_{\gamma+1} = s + 1$.
- (4) The domain of the parameters in H^r , for $r_l < r < r_{l+1}$, is an open subset of k , determined by $f^r(\lambda_0, \lambda_1, \dots, \lambda_l) \neq 0$, where

$$f^r(\lambda_0, \lambda_1, \dots, \lambda_l) = f(\lambda_0) \left(\prod_{l'=0}^{l-1} \prod_{r_l' < r' < r_{l'+1}} c^{r'}(\lambda_0, \lambda_1, \dots, \lambda_{l'}) \right) \prod_{r_l < r' \leq r} c^{r'}(\lambda_0, \lambda_1, \dots, \lambda_l), \quad (37)$$

and $c^{r'}(\lambda_0, \lambda_1, \dots, \lambda_{l'})$ are polynomials in $\lambda, \lambda_1, \dots, \lambda_{l'}$ appearing at r' -th reductions.

- (5) $f^{r_l}(\lambda_0, \lambda_1, \dots, \lambda_{l-1}) = f^{r_l-1}(\lambda_0, \lambda_1, \dots, \lambda_{l-1})$, for $l = 1, \dots, \gamma$.

Proof. (1) is obvious. We will prove (2), (3), (4) and (5) by induction on r . $r = 0$ is clear. Suppose that we have already had a sequence of parameterized bimodule problems

$$(\mathcal{K}, \mathcal{M}, H), (\mathcal{K}^1, \mathcal{M}^1, H^1), \dots, (\mathcal{K}^r, \mathcal{M}^r, H^r)$$

and parameters $\lambda_0, \lambda_1, \dots, \lambda_l$ in H^r with a polynomial

$$f^r(\lambda_0, \lambda_1, \dots, \lambda_l) \neq 0$$

for some $r \geq r_l$. Let $(\text{rad } \mathcal{K}^r)^*$ have a k -basis $\{v_1, v_2, \dots, v_m\}$. If $a_1 = (\rho^r)^*$ is the first arrow of $(\mathcal{K}^r, \mathcal{M}^r, H^r)$, then

$$\delta^c(a_1) = \sum_{j=1}^m f_j(\lambda_0, \lambda_1, \dots, \lambda_l) v_j \quad (38)$$

If $\delta^c(a_1) \neq 0$, we make a basis change in the field of rational functions $k(\lambda_0, \lambda_1, \dots, \lambda_l)$, say $(w_1, \dots, w_m)^T = Q(v_1, \dots, v_m)^T$ with the first row of Q being $(f_1, f_2, \dots, f_m)$. Then

$\delta(a_1) = w^{r+1}$ and thus $(H^{r+1})_{p_r q_r} = \emptyset$. Denote by $c^{r+1}(\lambda_0, \lambda_1, \dots, \lambda_l)$ the product of the denominators of the entries of Q and Q^{-1} , let

$$f^{r+1}(\lambda_0, \lambda_1, \dots, \lambda_l) = f^r(\lambda_0, \lambda_1, \dots, \lambda_l) c^{r+1}(\lambda_0, \lambda_1, \dots, \lambda_l).$$

In case of $\delta^c(a_1) = 0$, we set $r_{l+1} = r + 1$, $(H^{r+1})_{p_r q_r} = (\lambda_{l+1})$, and

$$f^{r+1}(\lambda_0, \lambda_1, \dots, \lambda_l) = f^r(\lambda_0, \lambda_1, \dots, \lambda_l).$$

Thus we complete the proof from $(\mathcal{K}^r, \mathcal{M}^r, H^r)$ to $(\mathcal{K}^{r+1}, \mathcal{M}^{r+1}, H^{r+1})$. Our assertion follows by induction. $\square$

5.4 Triangular formulae

Based on Lemma 5.3.1, we will in this subsection construct two kinds of freely parameterized bimodule problems which are established by Bautista. First we fix some notations.

Let $\mathfrak{A} = (\Gamma, \Omega)$ be a local boc, with a layer $L = (\Gamma'; \omega; a_1, a_2, \dots, a_n; v_1, v_2, \dots, v_m)$. Suppose that the indecomposable object of Γ' is $\mathcal{P}$, and $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda, f(\lambda)^{-1}]$. Denote by Δ the Γ' - Γ' -bimodule freely generated by $a_1, a_2, \dots, a_n$,

$$\Delta^{\otimes p} = \underbrace{\Delta \otimes_{\Gamma'} \Delta \otimes \dots \otimes_{\Gamma'} \Delta}_p$$

and $\Delta^{\otimes 0} = \Gamma'$. Then

$$\begin{cases} \Gamma = \bigoplus_{p \geq 0} \Delta^{\otimes p}, \\ \overline{\Omega} = \bigoplus_{p, q \geq 0} \bigoplus_{j=1}^m \Delta^{\otimes p} v_j \Delta^{\otimes q}. \end{cases} \quad (39)$$

For any $v \in \overline{\Omega}$, denote by v^0 the projection of v into $\bigoplus_{j=1}^m \Gamma' v_j \Gamma'$.

Definition 5.4.1 Given any constants $\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-1}^0 \in k$ with

$$f^{r\gamma}(\lambda_0^0, \dots, \lambda_{\gamma-1}^0) \neq 0$$

given in Lemma 5.3.1, and denote λ_γ by ν , we construct a local boc $\mathfrak{A}_{(\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-1}^0)}$ with a layer

$$L = (\Gamma'; \omega; b_1, \dots, b_j; u_1, \dots, u_m),$$

such that if $\gamma = 0$,

$$\mathfrak{A}_{(\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-1}^0)} = \mathfrak{A},$$

thus $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\nu, f(\nu)^{-1}]$; If $\gamma > 0$,

$$\mathfrak{A}_{(\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-1}^0)} = \mathfrak{A}^{r\gamma},$$

thus $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\nu]$, and ν appears for the first time in $\mathfrak{A}^{r_\gamma}$.

Lemma 5.4.1 There exists a sequence of localizations given by

$$\sigma_1(\nu), \sigma_2(\nu), \dots, \sigma_j(\nu),$$

and a triangular formula:

$$\begin{cases} \delta(b_1)^0 &= g_{11}(\nu, \kappa)u_1 \\ \delta(b_2)^0 &= g_{21}(\nu, \kappa)u_1 + g_{22}(\nu, \kappa)u_2 \\ \dots &\dots \\ \delta(b_j)^0 &= g_{j1}(\nu, \kappa)u_1 + g_{j2}(\nu, \kappa)u_2 + \dots + g_{j,j}(\nu, \kappa)u_j \end{cases} \quad (40)$$

where ν stands for the left multiplication by ν , and κ for the right multiplication by ν ;

$$g_{ll'}(\nu, \kappa) \in k[\nu, \kappa, \sigma_l(\nu)^{-1}\sigma_l(\kappa)^{-1}], \quad g_{ll'}(\nu, \nu) \neq 0, \quad l' \leq l, \quad l = 1, 2, \dots, j.$$

Proof. $j = 0$ is trivial. If $j > 0$, then $\delta^c(b_1) \neq 0$, since $\nu = \lambda_\gamma$ is the last parameter. Therefore $\delta(b) \neq 0$. Without loss of generality, we may divide the dotted arrows by some power of $f(\nu)f(\kappa)$ when $\gamma = 0$, such that the coefficients in $\delta(b_1)$ are all polynomials. After a localization given by $c_1(\nu)$ in Formula (30) of 4.3, $\delta(b_1) = g_{11}(\nu, \kappa)u_1$ with $g_{11}(\nu, \nu) \neq 0$. Let $\sigma_1(\nu) = f(\nu)c_1(\nu)$ if $\gamma = 0$, and $\sigma_1(\nu) = c_1(\nu)$ if $\gamma > 0$. In case of $j = 1$, the proof is completed. Otherwise,

$$\delta(b_2)^0 = g_{21}(\nu, \kappa)u_1 + \sum_{l=2}^m g'_{2l}(\nu, \kappa)u_{2l},$$

and $g'_{2l}(\nu, \kappa) \in k[\nu, \kappa, \sigma_1(\nu)^{-1}\sigma_1(\kappa)^{-1}]$. Without loss of generality we may divide u_{2l} by some power of $\sigma_1(\nu)\sigma_1(\kappa)$, such that $g'_{2l}(\nu, \kappa) \in k[\nu, \kappa]$. Suppose the highest common factor of $g'_{2l}(\nu, \kappa)$, $l = 2, \dots, m$, is $g_{22}(\nu, \kappa)$, and $c_2(\nu)$ is obtained by (30) to make a basis change from $u_{22}, \dots, u_{2m}$ to $u_2, u_{33}, \dots, u_{3m}$ in a Hermit ring $k[\nu, \kappa, \sigma_2(\nu)^{-1}\sigma_2(\kappa)^{-1}]$ with $\sigma_2(\mu) = \sigma_1(\mu)c_2(\mu)$. Then $g_{22}(\nu, \nu) \neq 0$. By induction we finally reach to the integer j and Formula (40). Which depends on a sequence of localizations given by

$$\sigma_l(\nu) = \begin{cases} f(\nu) \prod_{p=1}^l c_p(\nu), & \text{when } \gamma = 0; \\ \prod_{p=1}^l c_p(\nu), & \text{when } \gamma > 0, \end{cases} \quad \text{write } \sigma(\nu) = \sigma_j(\nu). \quad (41)$$

The proof is completed. $\square$

Definition 5.4.2 If for any $\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-1}^0$, $g_{ll}(\nu, \kappa) \in k[\nu, \kappa, \sigma_l(\nu)^{-1}\sigma_l(\kappa)^{-1}]$ are invertible, we may fix some constants $\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-2}^0$ with

$$f^{r_{\gamma-1}}(\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-2}^0) \neq 0.$$

Denote $\lambda_{\gamma-1}$ by λ , we then construct a local boc $\mathfrak{A}_{(\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-2}^0)} = (\Gamma, \Omega)$ with a layer

$$L = (\Gamma'; \omega'; a_1, \dots, a_i, b_1, \dots, b_j; v_1, \dots, v_m),$$

such that if $\gamma = 1$,

$$\mathfrak{A}_{(\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-2}^0)} = \mathfrak{A},$$

thus $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda, f(\lambda)^{-1}]$; if $\gamma > 1$,

$$\mathfrak{A}_{(\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-2}^0)} = \mathfrak{A}^{r_{\gamma-1}},$$

thus $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda]$, and λ appears for the first time in $\mathfrak{A}^{r_{\gamma-1}}$.

Lemma 5.4.2 There exists a localization given by $\tau(\lambda)$ and a triangular formula:

$$\left\{ \begin{array}{lcl} \delta(a_1)^0 & = & h_{11}(\lambda, \mu)w_1 \\ \delta(a_2)^0 & = & h_{21}(\lambda, \mu)w_1 + h_{22}(\lambda, \mu)w_2 \\ \dots & & \dots \\ \delta(a_{i-1})^0 & = & h_{i-1,1}(\lambda, \mu)w_1 + h_{i-1,2}(\lambda, \mu)w_2 + \dots + h_{i-1,i-1}(\lambda, \mu)w_{i-1} \\ \delta(a_i)^0 & = & h_{i1}(\lambda, \mu)w_1 + h_{i2}(\lambda, \mu)w_2 + \dots + h_{i,i-1}(\lambda, \mu)w_{i-1} + \tilde{h}(\lambda, \mu)\tilde{w} \end{array} \right. \quad (42)$$

where

$$h_{ll'}(\lambda, \mu) \in k[\lambda, \mu, \tau(\lambda)^{-1}\tau(\mu)^{-1}], \quad h_{ll}(\lambda, \lambda) \neq 0 \text{ for } l' \leq l \text{ and } l = 1, 2, \dots, i;$$

$w_1, w_2, \dots, w_{i-1}$ are free generators of $\overline{\Omega}$, $\tilde{w}$ is a linear form of $w_1, \dots, w_{i-1}$, or $\tilde{w}$ is linearly independent of $\{w_1, w_2, \dots, w_{i-1}\}$, and $\tilde{h}(\lambda, \mu) \neq 0$ but $\tilde{h}(\lambda, \lambda) = 0$.

Proof. If $\delta(a_1) = 0$ in Formula (29) of case A1 of 4.3, then $i = 1$, $\tilde{w} = 0$. Otherwise, after a localization given by $c_1(\lambda)$ in Formula (30) of 4.3, $\delta(a_1)^0 = h_{11}(\lambda, \mu)w_1$. If $h_{11}(\lambda, \lambda) = 0$, then $i = 1$, $\tilde{w} \neq 0$. If $h_{11}(\lambda, \lambda) \neq 0$, we use the similar procedure given in the proof of Lemma 5.4.1. Then we reach to Formula (42), since $a_i = \lambda_\gamma$ must appear at some stage i .

We use the following notations

$$\tau(\lambda) = \begin{cases} f(\lambda) \prod_{l=1}^i c_l(\lambda), & \text{when } \gamma = 1; \\ \prod_{l=1}^i c_l(\lambda), & \text{when } \gamma > 1, \end{cases} \quad \text{and } h(\lambda) = \tau(\lambda) \prod_{l=1}^{i-1} h_{ll}(\lambda, \lambda). \quad (43)$$

The proof is completed. $\square$

5.5 The minimal size assumption

Lemma 5.5.1. Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem, and $(*)$ be a reduction sequence consisting of freely parameterized triples, such that the end term $\mathfrak{A}^s$ of $(***)$ has two vertices, and is in one of the following configurations of Theorem 5.1.1:

$$\text{Case (1) } \lambda \begin{array}{c} \curvearrowright \\ \cdot \end{array} \xrightarrow[\substack{\mathcal{P} \\ a_1 \\ w_1}]{\mathcal{Q}} \mu, \quad \text{Case (2) } \lambda \begin{array}{c} \curvearrowright \\ \cdot \end{array} \xrightarrow[\mathcal{P}]{a_1} \cdot \mathcal{Q} \text{ (or dual).}$$


If we assume in addition that the size of the triples of $(*)$ is minimal such that $(*)$ meets any configuration of 5.1.1. Then the local boc $\mathfrak{A}_{\mathcal{P}}^s$ at $\mathcal{P}$ satisfies Formula (40) of 5.4 with all the

$$g_l(\lambda, \lambda') \in k[\lambda, \lambda', \sigma_l(\lambda)^{-1} \sigma_l(\lambda')^{-1}]$$

being invertible, where λ' stands for the right multiplication of λ . And so does $\mathfrak{A}_{\mathcal{Q}}^s$ at $\mathcal{Q}$.

Proof. Suppose the contrary, i.e. either we have Formula (42) of 5.4, or there exists some $1 \leq e \leq j$ with $g_{ee}(\lambda, \lambda') \in k[\lambda, \lambda', \sigma_e(\lambda)^{-1} \sigma_e(\lambda')^{-1}]$ being non-invertible in Formula (40). Then we would meet case (1) of 5.1.1 in sequence $(\hat{*})$, which is obtained from $(*)$ by deletion of $(T \setminus \{\mathcal{P}\})$ according to Corollary 4.3.1. But the size of the triples of $(\hat{*})$ is less than those of $(*)$, which is a contradiction to the minimal size assumption. $\square$

Lemma 5.5.2. Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem, and $(*)$ be a reduction sequence consisting of freely parameterized triples, such that the end term $\mathfrak{A}^s$ of $(***)$ is local and in the configuration of case (1) of Theorem 5.1.1:



Suppose that λ appears for the first time in $\mathfrak{A}^r$ attached to a vertex $\mathcal{P}$ for some $1 \leq r < s$. If we assume in addition that the size of the triples of $(*)$ is minimal, such that $(*)$ meets any configuration of 5.1.1. Then either

(1) $\mathfrak{A}^r$ is local, such that after performing the procedure of Lemma 5.3.1 starting from $\mathfrak{A}^r$, we have Formula (42) of 5.4, or Formula (40) with

$$g_{ee}(\lambda, \lambda') \in k[\lambda, \lambda', \sigma_e(\lambda)^{-1} \sigma_e(\lambda')^{-1}]$$

being non-invertible for some $1 \leq e \leq j$; or

(2) there exists some $r \leq l < s$, such that $\mathfrak{A}^l$ is in case (2) of Lemma 5.5.1.

Proof. (1) If $\gamma > 0$ in Lemma 5.3.1, we have done. If $\gamma = 0$, such index e must exist, since $\delta(a_1)$ is not invertible.

(2) If $\mathfrak{A}^r$ is not local, then the induced local boc $\mathfrak{A}_{\mathcal{P}}^r$ at $\mathcal{P}$ satisfies Formula (40) of 5.4 with all the $g_l(\lambda, \lambda') \in k[\lambda, \lambda', \sigma_l(\lambda)^{-1} \sigma_l(\lambda')^{-1}]$ being invertible by the minimal size assumption. There must exist some edges connecting $\mathcal{P}$ and other vertices. Otherwise $\mathcal{P}$ would be an isolated vertex, which contradicts to $\mathfrak{A}^s$ being local. Continue the reduction


from $\mathfrak{A}^r$, there must exist some first edge a_1 connecting $\mathcal{P}$ and other vertex in some induced boc $\mathfrak{A}^l$, such that $\delta(a_1) = 0$. Otherwise all such a_1 were deleted by regularizations, then $\mathcal{P}$ would be again an isolated vertex. Thus $\mathfrak{A}^l$ is in case (2) of Lemma 5.5.1 by the minimal size assumption. $\square$


5.6 The classification of minimally wild bimodule problems


Theorem 5.6.1. Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, and $(*)$ be a reduction sequence consisting of freely parameterized bimodule problems. Suppose that

- (1) the size of the triples of $(*)$ is minimal;
- (2) there exists some minimal integer r such that $\mathfrak{A}^r$ is in one of the configurations of Theorem 5.1.1.

Then there must exist a minimal integer s , such that the end term $\mathfrak{A}^s$ of $(***)$ is in one of the following five cases:

MW1.  Bocs $\mathfrak{B}_1$ has two vertices $\mathcal{P}, \mathcal{Q}$, and a layer $L_1 = (\Gamma'_1; \omega_1; a_1, \dots, a_n; v_1, \dots, v_m)$, where $\Gamma'_1(\mathcal{P}, \mathcal{P}) = k[\nu, g_{\mathcal{P}}(\nu)^{-1}]$, $\Gamma'_1(\mathcal{Q}, \mathcal{Q}) = k[\kappa, g_{\mathcal{Q}}(\kappa)^{-1}]$, $\delta(a_1) = f(\nu, \kappa)w_1$, such that $f(\nu, \kappa) \in k[\nu, \kappa, g_{\mathcal{P}}(\nu)^{-1}g_{\mathcal{Q}}(\kappa)^{-1}]$ is non-invertible. Moreover the local boc $(\mathfrak{B}_1)_{\mathcal{P}}$ at $\mathcal{P}$ satisfies Formula (40) of 5.4, with all the $g_{ll}(\nu, \nu') \in k[\nu, \nu', \sigma_l(\nu)^{-1}\sigma_l(\nu')^{-1}]$ given by Formula (41) being invertible, and so does $(\mathfrak{B}_1)_{\mathcal{Q}}$ at $\mathcal{Q}$.

MW2.  (or dual) Bocs $\mathfrak{B}_2$ has two vertices $\mathcal{P}, \mathcal{Q}$, and a layer $L_2 = (\Gamma'_2; \omega_2; a_1, \dots, a_n; v_1, \dots, v_m)$, where $\Gamma'_2(\mathcal{P}, \mathcal{P}) = k[\nu, g_{\mathcal{P}}(\nu)^{-1}]$, $\Gamma'_2(\mathcal{Q}, \mathcal{Q}) = k$, $\delta(a_1) = 0$. Moreover the induced local boc $(\mathfrak{B}_2)_{\mathcal{P}}$ at $\mathcal{P}$ satisfies Formula (40) of 5.4, with all the $g_{ll}(\nu, \nu') \in k[\nu, \nu', \sigma_l(\nu)^{-1}\sigma_l(\nu')^{-1}]$ given by Formula (41) being invertible.

MW3.  Bocs $\mathfrak{B}_3$ is local, and has a layer $L_3 = (\Gamma'_3; \omega_3; b_1, \dots, b_j; v_1, \dots, v_m)$, where $\Gamma'_3(\mathcal{P}, \mathcal{P}) = k[\nu]$, and the differentials $b_1, b_2, \dots, b_j$ are given by Formula (40) of 5.4, such that $g_{ll}(\nu, \kappa) \in k[\nu, \kappa, \sigma_l(\nu)^{-1}, \sigma_l(\kappa)^{-1}]$ for $1 \leq l \leq e-1$ are invertible, but $g_{e,e}(\nu, \kappa) \in k[\nu, \kappa, \sigma_e(\nu)^{-1}, \sigma_e(\kappa)^{-1}]$ is not invertible.

MW4.  Bocs $\mathfrak{B}_4$ is local, and has a layer $L_4 = (\Gamma'_4; \omega_4; a_1, \dots, a_i; b_1, \dots, b_j; v_1, \dots, v_m)$, where $\Gamma'_4(\mathcal{P}, \mathcal{P}) = k[\lambda]$, the differentials $a_1, a_2, \dots, a_i$ are given by Formula (42) of 5.4, such that

- (i) for any fixed $\lambda^0 \in k$, $h(\lambda^0) \neq 0$ in Formula (43), set $a_i = \nu$, then the differentials of $b_1, \dots, b_j$ given by Formulae(40) have the property that $g_l(\nu, \kappa) \in k[\nu, \kappa, \sigma_l(\nu)^{-1} \sigma_l(\kappa)^{-1}]$ are all invertible.
- (ii) $\tilde{w}$ is a linear combination of $w_1, \dots, w_{i-1}$ or $\tilde{w}$ is linearly independent of $w_1, \dots, w_{i-1}$, but $(\lambda - \mu)^2 \mid \tilde{h}(\lambda, \mu)$.

MW5.  Bocs $\mathfrak{B}_5$ is the same as $\mathfrak{B}_4$, except the hypothesis (ii) is changed to

(iii) $\tilde{w}$ is linearly independent of $w_1, \dots, w_{i-1}$, and $(\lambda - \mu)^2 \nmid \tilde{h}(\lambda, \mu)$.

Proof. Suppose that $\mathfrak{A}^r$ is in one case of the configurations of Theorem 5.1.1 with two vertices, then $r = s$ and we obtain MW1 or MW2 by Lemma 5.5.1. Now suppose $\mathfrak{A}^r$ is local, then λ appears for the first time in $\mathfrak{A}^r$ by Lemma 5.5.2. Denote λ by λ_0 , it ensures that the domain of λ_0 in $\mathfrak{A}^r$ is k . We perform the procedure given by Lemma 5.4.1. If there exist some $\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-1}^0$, and $1 \leq e \leq j$ in Lemma 5.4.1, such that $g_l(\nu, \kappa) \in k[\nu, \kappa, \sigma_l(\nu)^{-1}, \sigma_l(\kappa)^{-1}]$ for $1 \leq l \leq e - 1$ are invertible, but $g_{e,e}(\nu, \kappa) \in k[\nu, \kappa, \sigma_e(\nu)^{-1}, \sigma_e(\kappa)^{-1}]$ is not invertible, we have MW3. Otherwise, γ must be greater than zero, since $\mathfrak{A}^r$ is in case (1) of Theorem 5.1.1. Thus we may fix some $\lambda_0^0, \lambda_1^0, \dots, \lambda_{\gamma-2}^0$ and obtain MW4 or MW5. $\square$

Remark We always assume that the reductions are given according to the order of Definition 2.3.1, and the localizations are given to the parameters at the left with respect to the multiplication in case of the first arrows $a_1 : \mathcal{I} \rightarrow \mathcal{J}$ with $\mathcal{I}, \mathcal{J}$ both non-trivial. In particular Theorem 5.1.1 and 5.2.1 are proved by the above fixed procedure. Consequently if $\mathfrak{A}^s$ given in Theorem 5.6.1 satisfy MW3 or MW4 or MW5, then the parameter ν in $\mathfrak{B}_3$, or the parameter λ in $\mathfrak{B}_4$ or $\mathfrak{B}_5$ appears for the first time in $\mathfrak{A}^s$, i.e. $\Gamma'_3(\mathcal{P}, \mathcal{P}) = k[\nu]$, and $\Gamma'_4(\mathcal{P}, \mathcal{P}) = k[\lambda]$, $\Gamma'_5(\mathcal{P}, \mathcal{P}) = k[\lambda]$.

6 Exact categories and almost split conflations

6.1 Some basic concepts

Let $\mathcal{A}$ be an additive category with Krull-Schmidt property. We recall from [GR] and [DRSS] the following notions. A pair (ι, π) of composable morphisms $M \xrightarrow{\iota} E \xrightarrow{\pi} L$ in $\mathcal{A}$ is called *exact* if ι is a kernel of π and π is a cokernel of ι . Recall that ι is a *kernel* of π , provided that (1) ι is monic; (2) $\iota\pi = 0$; (3) $\forall \varphi : N \rightarrow E$ with $\varphi\pi = 0$, there exists a $\varphi' : N \rightarrow M$ such that $\varphi'\iota = \varphi$. Dually we have the definition of π being a *cokernel* of ι .

Let $\mathcal{E}$ be a class of exact pairs $M \xrightarrow{\iota} E \xrightarrow{\pi} L$ which is closed under isomorphisms. The morphisms ι and π appearing in a pair (ι, π) of $\mathcal{E}$ are called an *inflation* and a *deflation* of $\mathcal{E}$ respectively, and the pair itself is called a *conflation*. The class $\mathcal{E}$ is said to be an *exact structure* on $\mathcal{A}$, and $(\mathcal{A}, \mathcal{E})$ an *exact category* if the following axioms are satisfied:

E1. The composition of two deflations is a deflation.

E2. For each φ in $\mathcal{A}(L', L)$ and each deflation π in $\mathcal{A}(E, L)$, there are some E' in $\mathcal{A}$, an φ' in $\mathcal{A}(E', E)$ and a deflation $\pi' : E' \rightarrow L'$ such that $\pi'\varphi = \varphi'\pi$.

E3. Identities are deflations. If $\varphi\pi$ is a deflation, then so is π .

E3^{op}. Identities are inflations. If $\iota\varphi$ is an inflation, then so is ι .

An object L in $\mathcal{A}$ is said to be $\mathcal{E}$ -*projective* (or *projective* for short) if any conflation ending at L is split. Dually an object M in $\mathcal{A}$ is said to be $\mathcal{E}$ -*injective* (or *injective* for short) if any conflation starting at M is split.

Let $\mathcal{A}$ be a Krull-Schmidt category. A morphism $\pi : E \rightarrow L$ in $\mathcal{A}$ is called *right almost split* if it is not a retraction and for any non-retraction $\varphi : N \rightarrow L$, there exists a morphism $\psi : N \rightarrow E$ such that $\varphi = \psi\pi$. We say that $\mathcal{A}$ has *right almost split morphisms* if for all indecomposable L there exist right almost split morphisms ending at L . Dually we define *left almost split morphisms*. We say that $\mathcal{A}$ has *almost split morphisms* if $\mathcal{A}$ has right and left almost split morphisms.

A morphism $\pi : E \rightarrow L$ is called *right minimal* if every endomorphism $\eta : E \rightarrow E$ with the property that $\pi = \eta\pi$ is an isomorphism. A *left minimal morphism* $\iota : M \rightarrow E$ is defined dually.

Suppose that the Krull-Schmidt category $\mathcal{A}$ carries an exact structure $\mathcal{E}$. Let

$$(e) \quad M \xrightarrow{\iota} E \xrightarrow{\pi} L$$

be a conflation. Then the following assertions are equivalent.

(i) ι is minimal left almost split;

- (ii) π is minimal right almost split;
- (iii) ι is left almost split and π is right almost split.

The conflation (e) in the above condition is said to be an *almost split conflation*. The exact category $(\mathcal{A}, \mathcal{E})$ is said to *have almost split conflations* if

- (i) $\mathcal{A}$ has almost split morphisms;
- (ii) for any indecomposable non-projective L , there exists an almost split conflation (e) ending at L ;
- (iii) for any indecomposable non-injective M , there exists an almost split conflation (e) starting at M .

An almost split conflation $M \xrightarrow{\iota} E \xrightarrow{\pi} L$ is uniquely determined by M or L . We introduce the notation $M = \tau(L)$ or $L = \tau^{-1}(M)$.

Let us fix an exact category $(\mathcal{A}, \mathcal{E})$. For given objects M and L of $\mathcal{A}$, two conflations (ι, π) and (ι', π') starting from M and ending at L respectively are said to be *equivalent*, if there exists a commutative diagram as follows:

$$\begin{array}{ccccc} M & \xrightarrow{\iota} & E & \xrightarrow{\pi} & L \\ id \downarrow & & \downarrow \eta & & \downarrow id \\ M & \xrightarrow{\iota'} & E' & \xrightarrow{\pi'} & L \end{array}$$

6.2 Exact categories $P(\Lambda)$ and $P_1(\Lambda)$

The results of this subsection are mainly quoted from [B2] and [ZZ]. Let Λ be a finite-dimensional k -algebra, and $P(\Lambda)$ be defined in 2.1. A sequence of $P(\Lambda)$,

$$(e) \quad \begin{array}{ccccccc} 0 & \longrightarrow & P_1 & \xrightarrow{f_1} & W_1 & \xrightarrow{g_1} & Q_1 \longrightarrow 0 \\ & & \alpha \downarrow & & \downarrow \gamma & & \downarrow \beta \\ 0 & \longrightarrow & P_0 & \xrightarrow{f_0} & W_0 & \xrightarrow{g_0} & Q_0 \longrightarrow 0 \end{array} \quad (44)$$

is an *exact pair* if the two rows are exact and therefore split in $\Lambda\text{-mod}$. If $\mathcal{E}$ stands for the set of such exact pairs, then (e) is a conflation.

Theorem 6.2.1 $\mathcal{E}$ is an exact structure and $(P(\Lambda), \mathcal{E})$ is an exact category.

Proof. E1, E3 and $E3^{op}$ are obvious. We only check E2. If (g_1, g_0) given in (44) is a deflation, and $(\varphi_1, \varphi_0) : (Q'_1 \xrightarrow{\beta'} Q'_0) \rightarrow (Q_1 \xrightarrow{\beta} Q_0)$ is a morphism in $P(\Lambda)$. Then the following diagram gives a deflation (g'_1, g'_0) and a morphism (φ'_1, φ'_0) , which satisfy E2.

$$\begin{array}{ccccc}
W'_1 & \xrightarrow{g'_1} & Q'_1 & \xrightarrow{\beta'} & Q'_0 \\
\downarrow \varphi'_1 & \searrow \gamma & \downarrow \varphi'_0 & \downarrow \varphi_1 & \downarrow \varphi_0 \\
W_1 & \xrightarrow{g_1} & Q_1 & \xrightarrow{\beta} & Q_0 \\
\downarrow \varphi_1 & \searrow \gamma & \downarrow \varphi_0 & \downarrow \varphi_1 & \downarrow \varphi_0 \\
W_0 & \xrightarrow{g_0} & Q_0 & \xrightarrow{\beta} & Q_0
\end{array}$$

where $W'_0 = \{(w_0, q_0) \in W_0 \oplus Q'_0 \mid g_0(w_0) = \varphi_0(q_0)\}$, $W'_1 = \{(w_1, q_1) \in W_1 \oplus Q'_1 \mid g_1(w_1) = \varphi_1(q_1)\}$, $\gamma' = \begin{pmatrix} \gamma & 0 \\ 0 & \beta' \end{pmatrix}$, $g'_0 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, $g'_1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, $\varphi'_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\varphi'_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. $\square$

It is well known that $\text{rad}P \xrightarrow{\iota} P$ is a right almost split morphism in $\Lambda\text{-mod}$ for any indecomposable $P \in \Lambda\text{-proj}$. We denote by $r(P)$ a projective cover of $\text{rad}P$, and by ρ the composition of $r(P) \xrightarrow{\pi} \text{rad}P \xrightarrow{\iota} P$, then $r(P) \xrightarrow{\rho} P$ is a minimal right almost split morphism in $\Lambda\text{-proj}$. On the other hand, we denote by $\Lambda\text{-inj}$ the full subcategory of $\Lambda\text{-mod}$ consisting of injective Λ -modules. It is also well known that $I \xrightarrow{p} I/\text{soc}I$ is a left almost split morphism in $\Lambda\text{-mod}$ for any indecomposable $I \in \Lambda\text{-inj}$. There are two nice equivalent functors $F = D\Lambda \otimes_{\Lambda} - : \Lambda\text{-proj} \rightarrow \Lambda\text{-inj}$ and $G = \text{Hom}_{\Lambda}(D(\Lambda), -) : \Lambda\text{-inj} \rightarrow \Lambda\text{-proj}$. We denote $G(I)$ by P , and if $e(I)$ is an injective envelope of $I/\text{soc}(I)$, we denote $G(e(I))$ by $l(P)$. Let λ be the image of the composition $I \xrightarrow{p} I/\text{soc}I \xrightarrow{i} e(I)$ under the functor G , then $P \xrightarrow{\lambda} l(P)$ is a left almost split morphism in $\Lambda\text{-proj}$. Now consider the following objects of $P(\Lambda)$:

$$J(P) = (P \xrightarrow{id} P), \quad T(P) = (0 \xrightarrow{0} P), \quad Z(P) = (P \xrightarrow{0} 0)$$

and

$$R(P) = (r(P) \xrightarrow{\rho} P), \quad L(P) = (P \xrightarrow{\lambda} l(P)), \quad B(P) = (r(P) \xrightarrow{\rho\lambda} l(P))$$

We refer $U(P)$ to a minimal projective presentation of $\text{rad}P$, and $V(P)$ to the image of a minimal injective copresentation of $I/\text{soc}I$ under the functor G .

Lemma 6.2.1 (1) $\{J(P), T(P) \mid \forall \text{ indecomposable } P \in \Lambda\text{-proj}\}$ is a complete set of iso-classes of indecomposable $\mathcal{E}$ -projectives. Moreover, $R(P) \xrightarrow{(\rho \ id)} J(P)$ and $U(P) \xrightarrow{(0 \ \rho)} T(P)$ are right minimal almost split morphisms in $P(\Lambda)$.

(2) $\{J(P), Z(P) \mid \forall \text{ indecomposable } P \in \Lambda\text{-proj}\}$ is a complete set of iso-classes of indecomposable $\mathcal{E}$ -injectives. Moreover, $J(P) \xrightarrow{(id \ \lambda)} L(P)$ and $Z(P) \xrightarrow{(\lambda \ 0)} V(P)$ are left minimal almost split morphisms in $P(\Lambda)$. $\square$

Proposition 6.2.1 The exact category $(P(\Lambda), \mathcal{E})$ has almost split conflations. $\square$

We will pay most attention to $P_1(\Lambda)$ whose objects have no direct summands $J(P)$. If $\mathcal{E}_1$ stands for the induced exact structure of $\mathcal{E}$ restricted to $P_1(\Lambda)$, then $(P_1(\Lambda), \mathcal{E}_1)$ is also

an exact category.

Lemma 6.2.2 (1) $\{T(P), L(P) \mid \forall \text{ indecomposable } P \in \Lambda\text{-proj}\}$ is a complete set of iso-classes of indecomposable $\mathcal{E}_1$ -projectives. Moreover, $U(P) \xrightarrow{(0 \ \rho)} T(P)$ and $B(P) \xrightarrow{(\rho \ id)} L(P)$ are right minimal almost split morphisms in $P_1(\Lambda)$.

(2) $\{Z(P), R(P) \mid \forall \text{ indecomposable } P \in \Lambda\text{-proj}\}$ is a complete set of iso-classes of indecomposable $\mathcal{E}_1$ -injectives. Moreover, $Z(P) \xrightarrow{(\lambda \ 0)} V(P)$ and $R(P) \xrightarrow{(id \ \lambda)} B(P)$ are left minimal almost split morphisms in $P_1(\Lambda)$. $\square$

Proposition 6.2.2 The exact category $(P_1(\Lambda), \mathcal{E}_1)$ has almost split conflations. $\square$

Theorem 6.2.2 If (e) given by Formula (44) is a conflation in $P_1(\Lambda)$, and

$$Cok(e) : \quad Cok(\alpha) \xrightarrow{Cok(f_1, f_0)} Cok(\gamma) \xrightarrow{Cok(g_1, g_0)} Cok(\beta) \rightarrow 0$$

is the corresponding exact sequence in $\Lambda\text{-mod}$, such that $Cok(\alpha)$ is non-zero and neither injective nor simple. Then $Coker(e)$ is an almost split sequence in $\Lambda\text{-mod}$, if and only if (e) is an almost split conflation in $P_1(\Lambda)$.

Proof. By 2.3, 2.4 and 2.6 in [ZZ], Proposition 5.6 in [B2]. $\square$

6.3 Morphisms in $R(\mathfrak{A})$

Let $\mathfrak{A}$ be a layered boc. The purpose of this subsection is to transfer some morphisms of $R(\mathfrak{A})$ to the morphisms of a module category over an algebra in order to use some nice properties of module categories.

Lemma 6.3.1[Ro]. Let $\mathfrak{A} = (\Gamma, \Omega)$ be a boc with a layer $L = (\Gamma'; \omega; a_1, \dots, a_n; v_1, \dots, v_m)$, and $M \in R(\mathfrak{A})$. Suppose that

$$\{M'_{\mathcal{I}} = M_{\mathcal{I}} \mid \forall \mathcal{I} \in \text{ind}\Gamma'\}$$

is a set of k -vector spaces,

$$\{\varphi_{\mathcal{I}} = id : M_{\mathcal{I}} \rightarrow M'_{\mathcal{I}}; \varphi(v_j) : M_{\mathcal{I}_j} \rightarrow M'_{\mathcal{J}_j} \mid \forall \mathcal{I} \in \text{ind}\Gamma', v_j : \mathcal{I}_j \rightarrow \mathcal{J}_j\}$$

is a set of linear maps. Then there exists a unique $M' \in R(\mathfrak{A})$ such that $\varphi : M \rightarrow M'$ is an isomorphism in $R(\mathfrak{A})$.

Proof. We present a brief proof here for reader's convenience. Let $M'(\lambda) = M(\lambda)$ for any free parameter λ , and we set $M'(a_1) = M(a_1) - \varphi(\delta(a_1))$. Suppose that $M'(a_1), \dots, M'(a_{i-1})$ have been fixed, then we set $M'(a_i) = M(a_i) - \varphi(\delta(a_i))$, since $\delta(a_i)$ involves only $a_1, \dots, a_{i-1}$. Thus our assertion follows by induction. $\square$

Let $\mathfrak{A} = (\Gamma, \Omega)$ be a bocs with a layer $L = (\Gamma'; \omega; a_1, \dots, a_n; v_1, \dots, v_m)$. Then the bocs $\mathfrak{A}_0 = (\Gamma, \Gamma)$ is called the *principal bocs* of $\mathfrak{A}$. It is clear that the representation category $R(\mathfrak{A}_0)$ is just the module category $\Gamma\text{-mod}$. If $\varphi : M \rightarrow N$ is a morphism in $R(\mathfrak{A})$, then $\varphi = (\varphi_{\mathcal{I}}, \varphi(v_j) \mid \forall \mathcal{I} \in \text{ind}\Gamma', j = 1, 2, \dots, m)$ (see 4.6). We denote $(\varphi_{\mathcal{I}} \mid \forall \mathcal{I} \in \text{ind}\Gamma')$ by φ_0 . From now on we assume that the bocs is triangular also on $\{v_1, \dots, v_m\}$, i.e. $\delta(v_j)$ involves only $v_1, \dots, v_{j-1}$. If $\mathfrak{A}$ corresponds a bimodule problem, then it is obviously triangular on $\{v_1, \dots, v_m\}$.

Proposition 6.3.1 [O]. (1) If $\iota : M \rightarrow E$ is a morphism of $R(\mathfrak{A})$ with ι_0 injective, then there exists an isomorphism η and a commutative diagram in $R(\mathfrak{A})$:

$$\begin{array}{ccccc} M & \xrightarrow{\iota} & E \\ id \downarrow & & \downarrow \eta \\ 0 & \longrightarrow & M & \xrightarrow{\iota'} & E' \end{array}$$

such that the second sequence is exact in $R(\mathfrak{A}_0)$.

(2) If $\pi : E \rightarrow L$ is a morphism of $R(\mathfrak{A})$ with π_0 surjective, then there exists an isomorphism η and a commutative diagram in $R(\mathfrak{A})$:

$$\begin{array}{ccccc} E & \xrightarrow{\pi} & L \\ \eta \downarrow & & \downarrow id \\ E' & \xrightarrow{\pi'} & L & \longrightarrow & 0 \end{array}$$

such that the second sequence is exact in $R(\mathfrak{A}_0)$.

Proof. We present a complete proof here, since Ovsienko has not published his paper.

(1) Our proof will be shown by induction on the dotted arrows. First we fix a set of vector spaces $\{E_{\mathcal{I}}^1 = E_{\mathcal{I}} \mid \forall \mathcal{I} \in \text{ind}\Gamma'\}$ and a set of linear maps $\{\varphi_{\mathcal{I}} = id : E_{\mathcal{I}} \rightarrow E_{\mathcal{I}}^1; \varphi(v_j) = 0 \mid \forall \mathcal{I} \in \text{ind}\Gamma', \text{ and } j \neq 1\}$. $\varphi(v_1)$ will be defined below such that $\iota^1 = \iota\varphi : M \rightarrow E^1$ with $\iota^1(v_1) = 0$. Suppose $v_1 : \mathcal{I} \rightarrow \mathcal{J}$, then $0 = \iota^1(v_1) = \iota_{\mathcal{I}}\varphi(v_1) + \iota(v_1)\varphi_{\mathcal{J}}$ (see 4.6), i.e. $\iota_{\mathcal{I}}\varphi(v_1) + \iota(v_1) = 0$, which yields a commutative diagram since $\iota_{\mathcal{I}}$ is injective.:

$$\begin{array}{ccc} M_{\mathcal{I}} & \xrightarrow{\iota_{\mathcal{I}}} & E_{\mathcal{I}} \\ -\iota(v_1) \downarrow & & \downarrow \varphi(v_1) \\ E_{\mathcal{J}} & \xrightarrow{\varphi_{\mathcal{J}}=id} & E_{\mathcal{J}}^1 \end{array}$$

Thus $E^1 \in R(\mathfrak{A})$ is obtained by Lemma 6.3.1. And a commutative diagram

$$\begin{array}{ccc}
M & \xrightarrow{\iota} & E \\
id \downarrow & & \downarrow \eta^1 = \varphi \\
M & \xrightarrow{\iota^1} & E^1
\end{array}$$

follows from the structure with $\eta^1 = \varphi$, $\iota^1(v_1) = 0$. The assertion is obtained by the triangularity of $\{v_1, \dots, v_m\}$ and induction.

(2) is obtained dually. $\square$

Proposition 6.3.2 [O]. Let $\mathfrak{A}$ be a layered boc, and $(e) : M \xrightarrow{\iota} E \xrightarrow{\pi} L$ with $\iota\pi = 0$ be a pair of composable morphisms in $R(\mathfrak{A})$. If

$$(e_0) \quad 0 \longrightarrow M \xrightarrow{\iota_0} E \xrightarrow{\pi_0} L \longrightarrow 0$$

is exact in the category of vector spaces, then there exists an isomorphism η and a commutative diagram in $R(\mathfrak{A})$:

$$\begin{array}{ccccccc}
(e) & & M & \xrightarrow{\iota} & E & \xrightarrow{\pi} & L \\
& & id \downarrow & & \downarrow \eta & & \downarrow id \\
(e') & 0 & \longrightarrow & M & \xrightarrow{\iota'} & E' & \xrightarrow{\pi'} L \longrightarrow 0
\end{array}$$

such that (e') is an exact sequence in $R(\mathfrak{A}_0)$.

Proof. First we fix a set of vector spaces $\{E_{\mathcal{I}}^1 = E_{\mathcal{I}} \mid \forall \mathcal{I} \in \text{ind}\Gamma'\}$, and a set of linear maps $\{\varphi_{\mathcal{I}} = id : E_{\mathcal{I}} \rightarrow E_{\mathcal{I}}^1; \varphi(v_j) = 0 \mid \forall \mathcal{I} \in \text{ind}\Gamma', j \neq 1\}$. $\varphi(v_1)$ will be defined for the purpose of constructing some $E^1 \in R(\mathfrak{A})$ and some $\iota^1 = \iota\varphi : M \rightarrow E^1$, $\pi^1 = \varphi^{-1}\pi : E^1 \rightarrow L$ such that $\iota^1(v_1) = 0$, $\pi^1(v_1) = 0$. Suppose $v_1 : \mathcal{I} \rightarrow \mathcal{J}$, then $0 = \iota^1(v_1) = \iota_{\mathcal{I}}\varphi(v_1) + \iota(v_1)\varphi_{\mathcal{J}}$, i.e. $\iota_{\mathcal{I}}\varphi(v_1) + \iota(v_1) = 0$; $0 = \pi^1(v_1) = \varphi_{\mathcal{I}}^{-1}\pi(v_1) + \varphi^{-1}(v_1)\pi_{\mathcal{J}}$, i.e. $\pi(v_1) + \varphi^{-1}(v_1)\pi_{\mathcal{J}} = 0$ (see 4.6). Since $\varphi\varphi^{-1} = id$, $0 = (\varphi\varphi^{-1})(v_1) = \varphi_{\mathcal{I}}\varphi^{-1}(v_1) + \varphi(v_1)\varphi_{\mathcal{J}}^{-1}$, i.e. $\varphi^{-1}(v_1) = -\varphi(v_1)$. Therefore $\pi(v_1) - \varphi(v_1)\pi_{\mathcal{J}} = 0$. The hypothesis $(\iota\pi)(v_1) = 0$ given by $\iota\pi = 0$ yields a commutative diagram

$$\begin{array}{ccccc}
0 & \longrightarrow & M_{\mathcal{I}} & \xrightarrow{\iota_{\mathcal{I}}} & E_{\mathcal{I}} \\
& & \downarrow -\iota(v_1) & & \downarrow \pi(v_1) \\
& & E_{\mathcal{J}} & \xrightarrow{\pi_{\mathcal{J}}} & L_{\mathcal{J}} \longrightarrow 0
\end{array}$$

$\varphi(v_1) \cdot$ (dotted arrow from $M_{\mathcal{I}}$ to $E_{\mathcal{J}}$)
 ξ (dashed arrow from $L_{\mathcal{J}}$ to $E_{\mathcal{J}}$)

We claim that there exists a linear map $\varphi(v_1) : E_{\mathcal{I}} \rightarrow E_{\mathcal{J}}$ such that $\iota_{\mathcal{I}}\varphi(v_1) = -\iota(v_1)$ and $\varphi(v_1)\pi_{\mathcal{J}} = \pi(v_1)$. In fact, if $E_{\mathcal{I}} = \iota_{\mathcal{I}}(M_{\mathcal{I}}) \oplus N$ is taken in the category of vector spaces, we can define $((x)\iota_{\mathcal{I}} + y)\varphi(v_1) = (x)(-\iota(v_1)) + (y)\pi(v_1)\xi$ with $\xi : L_{\mathcal{J}} \rightarrow E_{\mathcal{J}}$ being

a retraction of $\pi_{\mathcal{J}}$. Thus $E^1 \in R(\mathfrak{A})$ is obtained by Lemma 6.3.1, and a commutative diagram

$$\begin{array}{ccccc} M & \xrightarrow{\iota} & E & \xrightarrow{\pi} & L \\ id \downarrow & & \downarrow \eta^1 = \varphi & & \downarrow id \\ M & \xrightarrow{\iota^1} & E^1 & \xrightarrow{\pi^1} & L \end{array}$$

follows from the structure, where $\eta^1 = \varphi$, $\iota^1(v_1) = 0$, $\pi^1(v_1) = 0$. The assertion is obtained by the triangularity of $\{v_1, \dots, v_m\}$ and induction. $\square$

6.4 An exact structure on $R(\mathfrak{A})$

There is a natural exact structure on $R(\mathfrak{A})$, which we will illustrate in this subsection.

Lemma 6.4.1 (1) $\iota : M \rightarrow E$ is monic if $\iota_0 : M \rightarrow E$ is injective.

(2) $\pi : E \rightarrow L$ is epic if $\pi_0 : E \rightarrow L$ is surjective.

Proof. (1) If ι_0 is injective, Proposition 6.3.1 (1) gives a commutative diagram with $\iota' : M \rightarrow E'$ in $R(\mathfrak{A}_0)$. Given any morphism $\varphi : N \rightarrow M$ with $\varphi\iota = 0$, we have $\varphi\iota\eta = \varphi\iota' = 0$. Then $\varphi_0\iota'_0 = 0$ yields $\varphi_0 = 0$; and for any $v : \mathcal{I} \rightarrow \mathcal{J}$, $0 = (\varphi_0\iota')(v) = \varphi(v)\iota'_j$ yields $\varphi(v) = 0$. Thus $\varphi = 0$ and ι is monic.

(2) is obtained dually. $\square$

Lemma 6.4.2 A pair of composable morphisms $(e) : M \xrightarrow{\iota} E \xrightarrow{\pi} L$ with $\iota\pi = 0$ is exact in $R(\mathfrak{A})$, if $(e_0) : 0 \rightarrow M \xrightarrow{\iota_0} E \xrightarrow{\pi_0} L \rightarrow 0$ is exact as a sequence of vector spaces.

Proof. (1) If (e_0) is exact, Lemma 6.4.1 (1) tells us that ι is monic.

(2) $\iota\pi = 0$.

(3) Proposition 6.3.2 gives a commutative diagram. If $\varphi : N \rightarrow E$ with $\varphi\pi = 0$, then $\varphi(\eta\pi') = 0$. Let $\xi = \varphi\eta$, then $\xi\pi' = 0$ implies that $\xi_{\mathcal{I}}\pi'_{\mathcal{I}} = 0$ and $\xi(v)\pi'_{\mathcal{J}} = 0$ for any vertex $\mathcal{I}$ and any dotted arrow $v : \mathcal{I} \rightarrow \mathcal{J}$. Let $\varphi' : N \rightarrow M$ be given by $\varphi'_{\mathcal{I}}\iota'_{\mathcal{I}} = \xi_{\mathcal{I}}$, $\varphi'(v)\iota'_{\mathcal{J}} = \xi(v)$, then we obtain $\varphi'\iota' = \xi$. Thus $\varphi'\iota'\eta^{-1} = \varphi$, i.e. $\varphi'\iota = \varphi$.

Therefore ι is a kernel of π . It can be proved dually that π is a cokernel of ι . $\square$

Definition 6.4.1 Let $\mathfrak{A}$ be a layered boc, we define a set $\mathcal{E}$ consisting of exact pairs $M \xrightarrow{\iota} E \xrightarrow{\pi} L$ in $R(\mathfrak{A})$, such that $\iota\pi = 0$ and

$$0 \rightarrow M \xrightarrow{\iota_0} E \xrightarrow{\pi_0} L \rightarrow 0$$

are exact as vector spaces. $\square$

It is clear that $\mathcal{E}$ is closed under isomorphisms.

Lemma 6.4.3 (1) $\iota : M \rightarrow E$ is an inflation, if and only if ι_0 is injective.

(2) $\pi : E \rightarrow L$ is a deflation, if and only if π_0 is surjective.

Proof. (1) If ι_0 is injective, Proposition 6.3.1 (1) gives an isomorphism $\eta : E \rightarrow E'$ in $R(\mathfrak{A})$, and a morphism ι' in $R(\mathfrak{A}_0)$. Moreover $\iota_0\eta_0 = \iota'$ ensures that ι' is injective. Let $L = \text{Coker}\iota'$, and $\pi' : E' \rightarrow L$ be the natural projection. Define $\pi = \eta\pi'$, we have the following commutative diagram:

$$\begin{array}{ccccc} & & E & & \\ & \nearrow \iota & \downarrow \eta & \searrow \pi & \\ M & & & & L \\ & \searrow \iota' & \downarrow \pi' & \nearrow & \\ & & E' & & \end{array}$$

Then $\iota\pi = \iota\eta \cdot \eta^{-1}\pi = \iota'\pi' = 0$. And $\eta_0^{-1}\pi'_0 = \pi_0$ ensures that π_0 is surjective. Thus $M \xrightarrow{\iota} E \xrightarrow{\pi} L$ belongs to $\mathcal{E}$, ι is an inflation.

(2) can be proved dually. □

Theorem 6.4.1 $\mathcal{E}$ is an exact structure on $R(\mathfrak{A})$, and $(R(\mathfrak{A}), \mathcal{E})$ is an exact category.

Proof. E1, E3 and E3^{op} are obvious. It suffices to prove E2. Suppose that $\pi : E \rightarrow L$ is a deflation, and $\varphi : L' \rightarrow L$ is a morphism in $R(\mathfrak{A})$. Consider the following diagram

$$\begin{array}{ccccc} E' & \xrightarrow{\pi'} & L' & & \\ & \searrow \kappa & \nearrow \eta & & \\ & N & & & \\ & \searrow \eta & \nearrow & & \\ & E \oplus L' & & & \\ & \nearrow (1) & \searrow (\pi) & & \\ E & \xrightarrow{\pi} & L & & \end{array}$$

$\varphi' \quad \downarrow \quad \varphi$

where $(\frac{\pi}{-\varphi})_0$ is surjective, since π_0 is already surjective. Thus there exists some object N and an isomorphism $\eta : N \rightarrow E \oplus L'$ in $R(\mathfrak{A})$ such that $\eta(\frac{\pi}{-\varphi}) : N \rightarrow L$ is surjective in $R(\mathfrak{A}_0)$ by item (2) of Proposition 6.3.1. Let $E' = \text{Ker}(\eta(\frac{\pi}{-\varphi}))$ and $\kappa : E' \rightarrow N$ be the natural embedding. We set $\pi' = \kappa\eta(\frac{\pi}{-\varphi})$. Since for any $y \in L'$, $(y)\varphi_0 \in L$, there exists some $x \in E$ with $(x)\pi_0 = (y)\varphi_0$. Consequently $(x, y)\eta_0^{-1}(\eta(\frac{\pi}{-\varphi})) = 0$, i.e. $(x, y)\eta_0^{-1} \in \kappa(E')$. And $(x, y)\eta_0^{-1}\kappa^{-1}\pi'_0 = y$. Therefore π'_0 is surjective and π is a deflation by Lemma 6.4.3. Let $\varphi' = \kappa\eta(\frac{\pi}{-\varphi})$. Since $\kappa\eta(\frac{\pi}{-\varphi}) = 0$, we have $\pi'\varphi = \varphi'\pi$. The proof of E2 is finished. □

We learnt recently that [BBP] proved the theorem similarly.

Corollary 6.4.1 Let $\mathfrak{A} = (\Gamma, \Omega)$ be a layered boc s and $(e) : M \xrightarrow{\iota} E \xrightarrow{\pi} L$ be a conflation of $R(\mathfrak{A})$. Then (e) is equivalent to an exact sequence

$$(e') \quad 0 \longrightarrow M \xrightarrow{\iota'} E' \xrightarrow{\pi'} L \longrightarrow 0$$

in $R(\mathfrak{A}_0)$. Moreover we can choose some suitable basis such that

$$\iota'_{\mathcal{L}} = (0, I), \quad \pi'_{\mathcal{L}} = \begin{pmatrix} I \\ 0 \end{pmatrix}$$

for any $\mathcal{L} \in \text{ind}\Gamma'$.

Proof. By Proposition 6.3.2. □

We apply the notions given in 6.3 and 6.4 to a freely parameterized bimodule problem to end this subsection.

Remark. Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem.

(1) Let $\mathcal{K}_0 = \{S \in \mathcal{K} \mid S \text{ are diagonal matrices}\}$, then $(\mathcal{K}_0, \mathcal{M})$ is said to be the *principal bimodule problem* of $(\mathcal{K}, \mathcal{M}, H)$.

(2) Let $M, L \in \text{Mat}(\mathcal{K}, \mathcal{M})$ with size vectors $\underline{m}, \underline{l}$ respectively. Let $S : M \rightarrow L$ be a morphism, and S_0 the partitioned diagonal part of S (see 2.2). Then S_0 is injective (resp. surjective) if and only if $\text{rank}(S_0) = m$ (resp. l), if and only if $\text{rank}(S_{\mathcal{I}}) = m_{\mathcal{I}}$ (resp. $l_{\mathcal{I}}$) for any $\mathcal{I} \in T/\sim$.

(3) Let $\mathcal{E}$ be a class consisting of composable pairs $M \xrightarrow{S} E \xrightarrow{R} L$ in $\text{Mat}(\mathcal{K}, \mathcal{M})$ such that $SR = 0$, and $0 \rightarrow M \xrightarrow{S_0} E \xrightarrow{R_0} L \rightarrow 0$ are exact as vector spaces. Then $\mathcal{E}$ is an exact structure and $(\text{Mat}(\mathcal{K}, \mathcal{M}), \mathcal{E})$ is an exact category.

6.5 Homogeneous conflations

Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem corresponding to a layered boc s $\mathfrak{A}$. We have seen in 6.4 that $\text{Mat}(\mathcal{K}, \mathcal{M})$ has an exact structure.

Definition 6.5.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem.

(1) An indecomposable object M in $\text{Mat}(\mathcal{K}, \mathcal{M})$ is said to be *homogeneous* or τ -*invariant* if there exists an almost split conflation $M \xrightarrow{\iota} E \xrightarrow{\pi} M$.

(2) The category $\text{Mat}(\mathcal{K}, \mathcal{M})$ is said to be *homogeneous* if for any positive integer n , almost all (except finitely many) iso-classes of indecomposable objects with size at most n are homogeneous.

(3) The category $\text{Mat}(\mathcal{K}, \mathcal{M})$ is said to be *strongly homogeneous* if there exists neither projectives nor injectives, and all indecomposables are homogeneous.

Sometimes we say that a bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ (or a boc $\mathfrak{A}$) is homogeneous, instead of saying $\text{Mat}(\mathcal{K}, \mathcal{M})$ (or $R(\mathfrak{A})$) for short.

Remark. We **do NOT require** that $\text{Mat}(\mathcal{K}, \mathcal{M})$ has almost split conflations globally in the sense of [DRSS] when we talk about homogeneous property. We only concentrate on the individual almost split conflations with the homogeneous starting and end terms. Furthermore we will see in subsection 7.2 that the homogeneous property of a bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ can be inherited to any induced bimodule problem $(\mathcal{K}', \mathcal{M}', H')$. Thus we are able to use reduction sequences and restrict our attention to some minimally wild cases.

Examples. (1) Let

$$\mathcal{K} = \left\{ \begin{pmatrix} s & 0 \\ 0 & s \end{pmatrix} \mid \forall s \in k \right\}, \mathcal{M} = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \mid \forall a, b \in k \right\}, H = 0, T = \{1, 2\}, 1 \sim 2.$$

Then $\text{Mat}(\mathcal{K}, \mathcal{M})$ is equivalent to $k\langle x, y \rangle\text{-mod}$. It has been shown by Vossieck in [ZL] that $\text{Mat}(\mathcal{K}, \mathcal{M})$ has no any almost split conflations.

(2) There are several sufficient conditions on a layered boc to have almost split conflations given in [CB1, BK, BB] etc. In particular, the Drozd boc of a finite-dimensional algebra Λ , or equivalently $P_1(\Lambda)$, has almost split conflations.

(3) If an algebra Λ is of tame representation type, then $P_1(\Lambda)$ and also $\Lambda\text{-mod}$ are homogeneous [CB1].

(4) Define an index set $T = \{1, 2\}$, with $1 \sim 2$. Let

$$\mathcal{K} = \left\{ \begin{pmatrix} s_1 & s_2 \\ 0 & s_1 \end{pmatrix} \mid s_1, s_2 \in k \right\}, \mathcal{M} = \left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \mid a, b \in k \right\}.$$

and $H = 0$. Then $\text{Mat}(\mathcal{K}, \mathcal{M})$ has almost split conflations, and it is strongly homogeneous [BCLZ].

6.6 Minimal bocses

Corollary 4.6.1 has shown the complete set of iso-classes of indecomposables of a minimal boc, we will show in the subsection the projectives, injectives and almost split conflations of a minimal boc.

Proposition 6.6.1 Let $\mathfrak{A} = (\Gamma, \Omega)$ be a minimal boc with a layer $L = (\Gamma'; \omega; v_1, \dots, v_m)$.

(1) For any trivial $\mathcal{J} \in \text{ind}\Gamma'$, $O(\mathcal{J})$ is projective and injective in $R(\mathfrak{A})$.

(2) For any non-trivial $\mathcal{I} \in \text{ind}\Gamma'$ with $\Gamma'(\mathcal{I}, \mathcal{I}) = k[\lambda, g_{\mathcal{I}}(\lambda)^{-1}]$, and for any $\lambda^0 \in k$, $g_{\mathcal{I}}(\lambda^0) \neq 0$.

$$O(\mathcal{I}, 1, \lambda^0) \xrightarrow{\iota_1} O(\mathcal{I}, 2, \lambda^0) \xrightarrow{\pi_1} O(\mathcal{I}, 1, \lambda^0)$$

and

$$O(\mathcal{I}, d, \lambda^0) \xrightarrow{(\pi_{d-1}, \iota_d)} O(\mathcal{I}, d-1, \lambda^0) \oplus O(\mathcal{I}, d+1, \lambda^0) \xrightarrow{(\iota_{d-1}, \pi_d)} O(\mathcal{I}, d, \lambda^0)$$

are almost split conflations in $R(\mathfrak{A})$, where $d \in \mathbb{N}, d \geq 2$.

Proof. (1) We claim first that for any indecomposable $M, N \in R(\mathfrak{A})$ with $M(\mathcal{I}) \neq 0, N(\mathcal{J}) \neq 0$ (see 4.6), if $\mathcal{I} \neq \mathcal{J}$, then $\text{Hom}_\Gamma(M, N) = 0$, since

$$\Gamma = \Gamma' = k[\lambda_1, g_1(\lambda_1)^{-1}] \times \cdots \times k[\lambda_i, g_i(\lambda_i)^{-1}] \times k_1 \times \cdots \times k_j$$

(see 4.2). If $\pi : E \rightarrow O(\mathcal{J})$ is a deflation in $R(\mathfrak{A})$, then π_0 is surjective by the definition of $\mathcal{E}$ in 6.4. Thus item (2) of Proposition 6.3.1 shows a commutative diagram with $E' \xrightarrow{\pi'} O(\mathcal{J}) \rightarrow 0$ being exact in $\Gamma\text{-mod}$. Therefore $E' = M \oplus O(\mathcal{J})^e$ with $e \geq 1$ and $\pi' = \begin{pmatrix} 0 \\ \varphi \end{pmatrix}$, where M does not contain any direct summand $O(\mathcal{J})$, and φ is a surjective linear map between vector spaces. Consequently φ splits, so do π' and π . Hence $O(\mathcal{J})$ is projective.

The injectivity of $O(\mathcal{J})$ can be proved dually.

(2) See Lemma 6.6 of [CB1]. □

Corollary 6.6.1 Let $\mathfrak{A}$ be a minimal boc, then $R(\mathfrak{A})$ has almost split conflations in the sense of [DRSS]. And $R(\mathfrak{A})$ is homogeneous.

Proof. There is a left almost split morphism: $O(\mathcal{J}) \rightarrow 0$, as well as a right almost split morphism: $0 \rightarrow O(\mathcal{J})$ for any trivial vertex $\mathcal{J}$. Therefore $R(\mathfrak{A})$ has almost split morphisms. The conclusion follows from Proposition 6.6.1. □

Next we show by an example that the exact structure $\mathcal{E}$ on $R(\mathfrak{A})$ defined in 6.4 collects only part of exact pairs of composable morphisms.

Example. Let $\mathfrak{A} = (\Gamma, \Omega)$ be a minimal boc with three trivial vertices $\mathcal{P}, \mathcal{Q}, \mathcal{L}$, and two dotted arrows $u : \mathcal{P} \dashrightarrow \mathcal{Q}, v : \mathcal{Q} \dashrightarrow \mathcal{L}$, such that $\delta(u) = 0, \delta(v) = 0$. Define a pair of composable morphisms

$$(e)M \xrightarrow{\iota} N \xrightarrow{\pi} L$$

where $M_{\mathcal{P}} = k, N_{\mathcal{Q}} = k, L_{\mathcal{L}} = k$, and M, N, L are all indecomposable; $\iota(u) = id, \pi(v) = id$. Then ι is a kernel of π , and π is a cokernel of ι . Thus (e) is an exact pair, which does not belong to $\mathcal{E}$. Moreover it is not difficult to see that (e) is even an almost split pair in $R(\mathfrak{A})$.

7 Almost split conflations in reductions

7.1 Projective and injective objects

This subsection is devoted to showing the finiteness of projectives (resp. injectives) in a layered boc and also behavior of the projectives (resp. injectives) in reductions.

Lemma 7.1.1 [B1] Let $\mathfrak{A} = (\Gamma, \Omega)$ be a boc with a layer $L = (\Gamma'; \omega; a_1, \dots, a_n; v_1, \dots, v_m)$, and $M \in R(\mathfrak{A})$ having dimension vector $\underline{m}$. Suppose that $\mathcal{P} \in \text{ind}\Gamma'$ is non-trivial. If $m_{\mathcal{P}} \neq 0$, then M is neither projective nor injective.

Proof. Let λ be a free parameter attached to $\mathcal{P}$ and $M(\lambda) \simeq J_d(\lambda^0) \oplus W$, where $d > 0$, and W is a Weyr matrix. Let us define a dimension vector $\underline{n}$ such that $n_{\mathcal{P}} = m_{\mathcal{P}} + 1$, $n_{\mathcal{I}} = m_{\mathcal{I}}$, $\forall \mathcal{I} \in \text{ind}\Gamma', \mathcal{I} \neq \mathcal{P}$. Let

$$\varphi_{\mathcal{P}} = \left(\begin{array}{c|c} I_d & 0_{d \times 1} \\ \hline & I \end{array} \right)_{m_{\mathcal{P}} \times n_{\mathcal{P}}}, \quad \psi_{\mathcal{P}} = \left(\begin{array}{c|c} I_d & \\ \hline 0_{1 \times d} & I \end{array} \right)_{n_{\mathcal{P}} \times m_{\mathcal{P}}}.$$

We now define an object N in $R(\mathfrak{A})$, such that $N(\lambda) = J_{d+1}(\lambda^0) \oplus W$, and $N(\mu) = M(\mu)$ for any parameter $\mu \neq \lambda$,

$$N(a_i) = \begin{cases} M(a_i), & \text{when } \mathcal{I} \neq \mathcal{P}, \mathcal{J} \neq \mathcal{P}; \\ M(a_i)\varphi_{\mathcal{P}}, & \text{when } \mathcal{I} \neq \mathcal{P}, \mathcal{J} = \mathcal{P}; \\ \psi_{\mathcal{P}}M(a_i), & \text{when } \mathcal{I} = \mathcal{P}, \mathcal{J} \neq \mathcal{P}; \\ \psi_{\mathcal{P}}M(a_i)\varphi_{\mathcal{P}}, & \text{when } \mathcal{I} = \mathcal{P} = \mathcal{J} \end{cases}$$

for any solid arrow $a_i : \mathcal{I} \rightarrow \mathcal{J}$. Then we obtain a morphism $\psi : N \rightarrow M$, such that $\psi_{\mathcal{I}} = I_{m_{\mathcal{I}}}$, for any $\mathcal{I} \neq \mathcal{P}$, and $\psi(v_j) = 0$, $j = 1, \dots, m$. Then $N(\lambda)\psi_{\mathcal{P}} = \psi_{\mathcal{P}}M(\lambda)$ and $N(a_i)\psi_{\mathcal{J}} = \psi_{\mathcal{J}}M(a_i)$, i.e. ψ is a morphism in $R(\mathfrak{A}_0)$. It is obvious that ψ is a deflation but not split. Therefore M is not projective.

The proof of M being non-injective is dual. $\square$

Proposition 7.1.1 (1) Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem and $(\mathcal{K}', \mathcal{M}', H')$ be an induced bimodule problem given by one of the 3 reductions of 3.1. If $M' \in \text{Mat}(\mathcal{K}', \mathcal{M}')$ is non-projective (resp. non-injective) then $\vartheta(M')$ is non-projective (resp. non-injective) in $\text{Mat}(\mathcal{K}, \mathcal{M})$.

(2) Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem with a reduction sequences $(*)$ of freely parameterized triples. If $M^s \in \text{Mat}(\mathcal{K}^s, \mathcal{M}^s)$ is non-projective (resp. non-injective), then $\vartheta_{r,s}(M^s)$ is non-projective (resp. non-injective) in $\text{Mat}(\mathcal{K}^r, \mathcal{M}^r)$ for $r = s - 1, \dots, 1, 0$.

Proof. (1) Since $M' \in \text{Mat}(\mathcal{K}', \mathcal{M}')$ is non-projective, there must exists a non-split deflation $\pi' : N' \rightarrow M'$ in $\text{Mat}(\mathcal{K}', \mathcal{M}')$, then $\pi = \pi' : \vartheta(N') \rightarrow \vartheta(M')$ is also a non-split

deflation since the surjectivity of π'_0 implies that of π_0 . In fact, $\pi'_0 = \pi_0$ in case of deletion and regularization, and π'_0 is the diagonal part of π_0 according to the partition $(T', \sim')$ in case edge and loop reductions. Therefore $M \in \text{Mat}(\mathcal{K}, \mathcal{M})$ is not projective.

The proof for non-injective case is dual.

(2) By (1) and induction on $r = s, s-1, \dots, 1, 0$. $\square$

Corollary 7.1.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem. Then for each positive integer n , there are only finitely many iso-classes of projectives and injectives of size at most n in $\text{Mat}(\mathcal{K}, \mathcal{M})$.

Proof. It is clear that there are only finitely many choices of size vectors $\underline{m}$ over $(T, \sim)$ with $m \leq n$. Let $M \in \text{Mat}(\mathcal{K}, \mathcal{M})$ be a projective matrix of size vector $\underline{m}$. Then $m_{\mathcal{P}} = 0$, for any non-trivial $\mathcal{P} \in T/\sim$ by Lemma 7.1.1. Therefore $H_{\underline{m}}$ is a constant matrix. Now we start a sequence of reductions according to Theorem 3.4.1 for M . We conclude that there does not exist any loop reduction in the sequence. Otherwise, if the $(r+1)$ -th reduction were a loop reduction, M^r would be non-projective in $\text{Mat}(\mathcal{K}^r, \mathcal{M}^r)$ still by Lemma 7.1.1. Therefore M would be non-projective in $\text{Mat}(\mathcal{K}, \mathcal{M})$ by proposition 7.1.1, which contradicts to our assumption. It follows obviously that there are only finitely many choices of such a sequence consisting of regularizations and edge reductions with size vector $\underline{m}$. Summarizing the discussion, there are only finitely many iso-classes of projectives of size at most n in $\text{Mat}(\mathcal{K}, \mathcal{M})$.

The proof for injectives is dual. $\square$

7.2 Homogeneous property in reductions

We will give the key idea for proving the main theorem in this subsection, which was originally presented by R. Bautista.

Lemma 7.2.1[B1]. Let $(\mathcal{K}, \mathcal{M}, H)$ be a freely parameterized bimodule problem, and $(\mathcal{K}', \mathcal{M}', H')$ be an induced bimodule problem given by one of the 4 reductions in 3.1 or 3.2.

(1) Suppose that $M', L' \in \text{Mat}(\mathcal{K}', \mathcal{M}')$ have size vectors $\underline{m}', \underline{l}'$, and $\vartheta(M'), \vartheta(L') \in \text{Mat}(\mathcal{K}, \mathcal{M})$ have size vectors $\underline{m}, \underline{l}$ respectively. If $\underline{m} = \underline{l}$, and $\underline{m}' \leq \underline{l}'$, then $\underline{m}' = \underline{l}'$.

(2) If $\iota' : M' \rightarrow E'$ is a morphism in $\text{Mat}(\mathcal{K}', \mathcal{M}')$ with $\vartheta(\iota') : \vartheta(M') \rightarrow \vartheta(E')$ being a left minimal almost split morphism of $\text{Mat}(\mathcal{K}, \mathcal{M})$, then so is ι' . Dually if $\pi' : E' \rightarrow L'$ is a morphism in $\text{Mat}(\mathcal{K}', \mathcal{M}')$ with $\vartheta(\pi') : \vartheta(E') \rightarrow \vartheta(L')$ being a right minimal almost split morphism of $\text{Mat}(\mathcal{K}, \mathcal{M})$, then so is π' in $\text{Mat}(\mathcal{K}', \mathcal{M}')$.

(3) Let $(e') M' \xrightarrow{\iota'} E' \xrightarrow{\pi'} L'$ be a composable pair of $\text{Mat}(\mathcal{K}', \mathcal{M}')$ with M' non-

injective and L' non-projective. If $\vartheta(e')$ is an almost split conflation of $\text{Mat}(\mathcal{K}, \mathcal{M})$, then so is (e') in $\text{Mat}(\mathcal{K}', \mathcal{M}')$.

Proof. (1) It is obvious in case of deletion. The other cases given by reductions in 3.1 follow from Formula (11) of 3.2.

(2) The functor $\vartheta : \text{Mat}(\mathcal{K}', \mathcal{M}') \rightarrow \text{Mat}(\mathcal{K}, \mathcal{M})$ is fully faithful.

(3) Since L' is non-projective, there exist some N' and non-split morphism $\varphi' : N' \rightarrow L'$ with φ'_0 being surjective. But π' is a right almost split morphism by (2), we have a morphism $\psi' : N' \rightarrow E'$ such that $\psi'\pi' = \varphi'$. Consequently $\psi'_0\pi'_0 = \varphi'_0$ and π'_0 is surjective. Similarly ι'_0 is injective. Since $\vartheta(\iota')\vartheta(\pi') = 0$, we have $\iota'\pi' = 0$ and $\iota'_0\pi'_0 = 0$. Therefore $\underline{e}' \geq \underline{m}' + \underline{l}'$. But $\underline{e} = \underline{m} + \underline{l}$, so $\underline{e}' = \underline{m}' + \underline{l}'$ by (1). Hence $M' \xrightarrow{\iota'_0} E' \xrightarrow{\pi'_0} L'$ is exact as a sequence of vector spaces, i.e. (ι', π') is a conflation. And (e') is an almost split conflation of $\text{Mat}(\mathcal{K}', \mathcal{M}')$ by (2). $\square$

Theorem 7.2.1 Let $\mathfrak{A} = (\Gamma, \Omega)$ be a layered boc, and $\mathfrak{A}' = (\Gamma', \Omega')$ be induced by one of the 4 reductions of 3.1 or 3.2 with a reduction functor $\vartheta : R(\mathfrak{A}') \rightarrow R(\mathfrak{A})$. Suppose that an indecomposable $M' \in R(\mathfrak{A}')$ is neither projective nor injective. If

$$(e) \quad \vartheta(M') \xrightarrow{\iota} E \xrightarrow{\pi} \vartheta(M')$$

is an almost split conflation of $R(\mathfrak{A})$, then

- (1) there exists some $E' \in R(\mathfrak{A}')$ such that $\vartheta(E') \simeq E$;
- (2) there exists an almost split conflation

$$(e') \quad M' \xrightarrow{\iota'} E' \xrightarrow{\pi'} M'$$

in $R(\mathfrak{A}')$ such that $\vartheta(e')$ is equivalent to (e) .

Proof. We assume that $\mathfrak{A}$ is the corresponding boc of bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ and use the notions of both structures freely. Denote $\vartheta(M')$ by M for short, then we first set up two claims.

Claim 1. In case of deletion, both conclusions (1) and (2) hold.

In fact, since

$$(e_0) \quad 0 \longrightarrow \vartheta(M') \xrightarrow{\iota_0} E \xrightarrow{\pi_0} \vartheta(M') \longrightarrow 0$$

is exact as a sequence of vector spaces, and $M = \vartheta(M') = M'$, we conclude that $E_{\mathcal{I}} = 0$, $\forall \mathcal{I} \notin T'/\sim'$. Hence E belongs to $R(\mathfrak{A}')$. Therefore $E' = E$, $(e') = (e)$.

Claim 2. In other cases, (2) is true if (1) holds. Indeed, suppose the commutative diagram below comes from (1):

$$\begin{array}{ccccc}
& & E & & \\
& \nearrow \iota & \downarrow \eta & \searrow \pi & \\
\vartheta(M') & & & & \vartheta(M') \\
& \searrow \iota\eta & & \nearrow \eta^{-1}\pi & \\
& & \vartheta(E') & &
\end{array}$$

Since ϑ is fully faithful, there exists some $\iota' : M' \rightarrow E'$ with $\vartheta(\iota') = \iota\eta$ and $\pi' : E' \rightarrow M'$ with $\vartheta(\pi') = \eta^{-1}\pi$. We obtain a pair of morphisms (e') in $R(\mathfrak{A}')$ satisfying the hypothesis of item (3) of Lemma 7.2.1. Therefore, (e') is an almost split conflation in $R(\mathfrak{A}')$ with $\vartheta(e')$ equivalent to (e) .

Then it suffices to prove assertion (1) according to the 3 reductions in 3.1 respectively.

Regularization. (1) is valid because of ϑ being an equivalence.

Edge reduction. We consider first an edge reduction given by $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Let $\mathfrak{A}''$ be the induced bocs and $\vartheta_1 : R(\mathfrak{A}'') \rightarrow R(\mathfrak{A})$ be the equivalent functor. Thus $\mathfrak{A}'$ is induced from $\mathfrak{A}''$ by a deletion of $(T''/\sim'') \setminus (T'/\sim')$. And the following diagram commutes:

$$\begin{array}{ccc}
R(\mathfrak{A}') & \xrightarrow{\vartheta_2} & R(\mathfrak{A}'') \\
\vartheta \searrow & & \nearrow \vartheta_1 \\
& R(\mathfrak{A}) &
\end{array} \tag{45}$$

Since $M' \in R(\mathfrak{A}')$ is neither projective nor injective, $\vartheta_2(M') \in R(\mathfrak{A}'')$ is so by item (1) of Proposition 7.1.1. And since ϑ_1 is an equivalence, there exists some $E'' \in R(\mathfrak{A}'')$ with $E \simeq \vartheta_1(E'')$. Therefore we have an almost split conflation

$$(e'') : \vartheta_2(M') \xrightarrow{\iota''} E'' \xrightarrow{\pi''} \vartheta_2(M')$$

in $R(\mathfrak{A}'')$ with $\vartheta_1(e'')$ equivalent to (e) by Claim 2. Furthermore, $M' = \vartheta_2(M')$ and $E' = E''$, $(e') = (e'')$ follow from deletion given by Claim 1. Thus, $\vartheta(E') \simeq E$ and $\vartheta(e')$ is equivalent to (e) as desired.

Loop reduction. Suppose the reduction is given by $W \oplus (\lambda)$ (or just W), and the domain of λ equals $k \setminus \{\text{the roots of } g(\lambda)\}$. If $E_{pq} \simeq W_1 \oplus W_2$, where the eigenvalues of W_2 belong to the domain of λ , but those of W_1 do not. Then we make a new loop reduction given by

$$W \oplus W_1 \oplus (\lambda) \text{ (or } W \oplus W_1).$$

Consequently we obtain a new induced boc $\mathfrak{A}''$ from $\mathfrak{A}$ by loop reduction, and $\mathfrak{A}'$ is induced from $\mathfrak{A}''$ by deletion of $(T''/\sim'') \setminus (T'/\sim')$. Then there exist some $E'' \in R(\mathfrak{A}'')$ such that $E \cong \vartheta_1(E'')$ and an almost split conflation (e'') in $R(\mathfrak{A}'')$ such that $\vartheta_1(e'')$ is equivalent to (e) . The diagram (45) shows a similar proof for the case of loop reduction. $\square$

Corollary 7.2.1 (1) Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem and $(\mathcal{K}', \mathcal{M}', H')$ induced from $(\mathcal{K}, \mathcal{M}, H)$ by one of the four reductions given by 3.1. If $(\mathcal{K}, \mathcal{M}, H)$ is homogeneous, then so is $(\mathcal{K}', \mathcal{M}', H')$.

(2) For a given bimodule problem $(\mathcal{K}, \mathcal{M}, H)$, if there exists a reduction sequence $(*)$ of freely parameterized triples such that $Mat(\mathcal{K}^s, \mathcal{M}^s)$ is non-homogeneous, then $Mat(\mathcal{K}, \mathcal{M})$ must be non-homogeneous.

Proof. (1) Given any fixed positive integer n , we define a set of iso-classes of objects

$$\mathcal{S} = \{M_\alpha \in Mat(\mathcal{K}', \mathcal{M}') \mid M_\alpha \text{ is indecomposable of size at most } n, \\ \text{and is neither projective nor injective}\}.$$

Because of Corollary 7.1.1, $\mathcal{S}$ is a cofinite subset. If $\mathcal{S}$ is a finite set, we are done. If $\mathcal{S}$ is an infinite set, then $\{\vartheta(M_\alpha) \mid M_\alpha \in \mathcal{S}\}$ is an infinite set of objects in $Mat(\mathcal{K}, \mathcal{M})$, where $\vartheta(M_\alpha)$ is neither projective nor injective by Proposition 7.1.1, and the size of $\vartheta(M_\alpha)$ is bounded by n . Since $Mat(\mathcal{K}, \mathcal{M})$ is homogeneous, there exists a cofinite subset $\mathcal{S}_0$ of $\mathcal{S}$, such that $\forall M_\alpha \in \mathcal{S}_0$, there are almost split conflations of $Mat(\mathcal{K}, \mathcal{M})$:

$$(e_\alpha) \quad \theta(M_\alpha) \xrightarrow{\iota} E_\alpha \xrightarrow{\pi} \theta(M_\alpha)$$

Thus there are almost split conflations of $R(\mathfrak{A}')$

$$(e'_\alpha) \quad M_\alpha \xrightarrow{\iota'} E'_\alpha \xrightarrow{\pi'} M_\alpha$$

with $\vartheta(e'_\alpha)$ being equivalent to (e_α) by Theorem 7.2.1. Consequently $Mat(\mathcal{K}', \mathcal{M}')$ is homogeneous.

(2) If $Mat(\mathcal{K}, \mathcal{M})$ were homogeneous, then $Mat(\mathcal{K}^r, \mathcal{M}^r)$ would be homogeneous for $r = 1, 2, \dots, s$ by (1) and induction on r , which is a contradiction to the hypothesis. $\square$

7.3 Critically non-homogeneous bocses $\mathfrak{B}_1$ and $\mathfrak{B}_2$

Lemma 7.3.1 Let $f(\nu, \kappa)$ and $g(\nu, \kappa)$ be polynomials in two indeterminates ν, κ . If f and g are coprime, then there exists an infinite list of pairs (ν^0, κ^0) , such that $f(\nu^0, \kappa^0) = 0$ but $g(\nu^0, \kappa^0) \neq 0$. Thus ν^0 or κ^0 must range over a cofinite subset D of k .

Proof. By means of Bezout's theorem. $\square$

Corollary 7.3.1 Let $g_{\mathcal{P}}(\nu), \sigma_{\mathcal{P}}(\nu) \in k[\nu]$ with $g_{\mathcal{P}}(\nu) \mid \sigma_{\mathcal{P}}(\nu)$, and $g_{\mathcal{Q}}(\kappa), \sigma_{\mathcal{Q}}(\kappa) \in k[\kappa]$ with $g_{\mathcal{Q}}(\kappa) \mid \sigma_{\mathcal{Q}}(\kappa)$. If

$$f(\nu, \kappa) \in k[\nu, \kappa, g_{\mathcal{P}}(\nu)^{-1}, g_{\mathcal{Q}}(\kappa)^{-1}]$$

is not invertible, then there exists an infinite list of pairs (ν^0, κ^0) with $f(\nu^0, \kappa^0) = 0$, such that either

- (1) $\sigma_{\mathcal{P}}(\nu^0)g_{\mathcal{Q}}(\kappa^0) \neq 0$, and ν^0 ranges over a cofinite subset $D_{\mathcal{P}}$ of k ; or
- (2) $g_{\mathcal{P}}(\nu^0)\sigma_{\mathcal{Q}}(\kappa^0) \neq 0$, and κ^0 ranges over a cofinite subset $D_{\mathcal{Q}}$ of k .

Proof. If $f(\nu, \kappa) \neq 0$, let

$$f(\nu, \kappa) = \bar{f}(\nu, \kappa)\bar{g}_{\mathcal{P}}(\nu)\bar{g}_{\mathcal{Q}}(\kappa),$$

such that $\bar{f}(\nu, \kappa)$ is a polynomial and $(\bar{f}(\nu, \kappa), g_{\mathcal{P}}(\nu)g_{\mathcal{Q}}(\kappa)) = 1$; and $\bar{g}_{\mathcal{P}}(\nu)$ is a product of some positive or negative powers of the factors of $g_{\mathcal{P}}(\nu)$, $\bar{g}_{\mathcal{Q}}(\kappa)$ is that of $g_{\mathcal{Q}}(\kappa)$. Suppose that

$$\bar{f}(\nu, \kappa) = f_0(\nu, \kappa)f_{\mathcal{P}}(\nu)f_{\mathcal{Q}}(\kappa),$$

such that $f_0(\nu, \kappa)$ has no non-constant factors taken from $k[\nu]$ or $k[\kappa]$.

(i) If $f_0(\nu, \kappa)$ is not a constant, then $(f_0(\nu, \kappa), \sigma_{\mathcal{P}}(\nu)\sigma_{\mathcal{Q}}(\kappa)) = 1$, then there exists an infinite list of pairs (ν^0, κ^0) , such that $f_0(\nu^0, \kappa^0) = 0$, thus $f(\nu^0, \kappa^0) = 0$, but $\sigma_{\mathcal{P}}(\nu^0)\sigma_{\mathcal{Q}}(\kappa^0) \neq 0$. When ν^0 ranges over a cofinite subset $D_{\mathcal{P}}$ of k , we have (1); when κ^0 ranges over a cofinite subset $D_{\mathcal{Q}}$ of k , we have (2).

(ii) If $f_0(\nu, \kappa)$ is a constant, but $f_{\mathcal{Q}}(\kappa)$ is not, then $(f_{\mathcal{Q}}(\kappa), \sigma_{\mathcal{P}}(\nu)g_{\mathcal{Q}}(\kappa)) = 1$. $\forall \nu^0 \in k$ with $\sigma_{\mathcal{P}}(\nu^0) \neq 0$, there exists some κ^0 with $f_{\mathcal{Q}}(\kappa^0) = 0$ but $g_{\mathcal{Q}}(\kappa^0) \neq 0$, we obtain item (1).

(iii) If $f_0(\nu, \kappa)$ and $f_{\mathcal{Q}}(\kappa)$ are both constants, then $f_{\mathcal{P}}(\nu)$ must be not, thus $(f_{\mathcal{P}}(\nu), g_{\mathcal{P}}(\nu) \cdot \sigma_{\mathcal{Q}}(\kappa)) = 1$. $\forall \kappa^0 \in k$ with $\sigma_{\mathcal{Q}}(\kappa^0) \neq 0$, there exists some ν^0 with $f_{\mathcal{P}}(\nu^0) = 0$ but $g_{\mathcal{P}}(\nu^0) \neq 0$, we obtain item (2).

Finally, if $f(\nu, \kappa) = 0$, it is clear that we have (1) and (2). $\square$

Proposition 7.3.1 The boc $\mathfrak{B}_1$ given in MW1 of Theorem 5.6.1 is not homogeneous.

Proof. Consider the local bocses $(\mathfrak{B}_1)_{\mathcal{P}}$ at $\mathcal{P}$ and $(\mathfrak{B}_1)_{\mathcal{Q}}$ at $\mathcal{Q}$, suppose the triangular formula (40) of 5.4 gives the polynomials $\sigma_{\mathcal{P}}(\nu)$ and $\sigma_{\mathcal{Q}}(\kappa)$ in (41) respectively. Assume that we have item (1) of Corollary 7.3.1, i.e. there is an infinite list $\{(\nu^0, \kappa^0)\}$, such that $f(\nu^0, \kappa^0) = 0$, $\sigma_{\mathcal{P}}(\nu^0)g_{\mathcal{Q}}(\kappa^0) \neq 0$ and ν^0 ranges over a cofinite subset $D = D_{\mathcal{P}}$ of k . If $R(\mathfrak{B}_1)$ is homogeneous, we present an infinite list of one dimensional objects S_{ν^0} , $\forall \nu^0 \in D$,

with

$$(S_{\nu^0})_{\mathcal{P}} = k, (S_{\nu^0})_{\mathcal{Q}} = 0, S_{\nu^0}(\nu) = (\nu^0), S_{\nu^0}(a_l) = 0 \text{ or } \emptyset \text{ for } l = 1, \dots, i,$$

when we calculate S_{ν^0} according to Theorem 3.4.1. If $R(\mathfrak{B}_1)$ is homogeneous, there must be a cofinite subset $D_0 \subset D$, with S_{ν^0} being homogeneous for any $\nu^0 \in D_0$. We fix such a ν^0 , and denote S_{ν^0} by S for simplicity. Let $(e) : S \xrightarrow{\iota} E \xrightarrow{\pi} S$ be the almost split conflation starting and ending at S in $R(\mathfrak{B}_1)$, then (e) is also an almost split conflation of $R((\mathfrak{B}_1)_{\mathcal{P}})$ according to Theorem 7.2.1. Thus

$$E_{\mathcal{P}} = k^2, \quad E_{\mathcal{Q}} = 0, \quad E(\nu) = \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix}$$

by Proposition 6.6.1 and $E(a_l) = \emptyset$ for all the solid loops a_l at $\mathcal{P}$. We define an object $L \in R(\mathfrak{B}_1)$ with

$$L_{\mathcal{P}} = k, \quad L_{\mathcal{Q}} = k, \quad L(\nu) = (\nu^0), \quad L(\kappa) = (\kappa^0), \quad L(a_1) = (1), \quad L(a_l) = 0, \quad \forall l \geq 2,$$

and we also define a morphism $\varphi : L \rightarrow S$ given by

$$\varphi_{\mathcal{P}} = (1), \quad \varphi_{\mathcal{Q}} = (0), \quad \varphi(v) = 0$$

for any dotted arrow v .

We first claim that φ is not a split epimorphism. In fact if there were some morphism $\psi : S \rightarrow L$ with $\psi\varphi = id$, then we would have

$$\varphi_{\mathcal{P}} = (1), \quad \psi_{\mathcal{Q}} = (0); \quad S(a_1)\psi_{\mathcal{Q}} - \psi_{\mathcal{P}}L(a_1) = \psi(\delta(a_1)) = f(\nu^0, \kappa^0)\psi(w_1),$$

i.e. $-1 = 0$, which leads to a contradiction. Thus there exists some $\varphi' : L \rightarrow E$ such that $\varphi'\pi = \varphi$, since (e) is an almost split conflation. Then $\varphi'_{\mathcal{P}}\pi_{\mathcal{P}} = \varphi_{\mathcal{P}}$, i.e. $(a, b)\begin{pmatrix} 1 \\ 0 \end{pmatrix} = 1$, $\varphi'_{\mathcal{P}} = (1, b)$. On the other hand, $\varphi'_{\mathcal{P}}E(\nu) = L(\nu)\varphi'_{\mathcal{P}}$ implies that

$$(1 \ b) \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix} = \nu_0(1, \ b),$$

i.e. $(\nu^0, 1 + b\nu^0) = (\nu^0, \nu^0 b)$, which results in a contradiction.

The proof in the case of item (2) of Corollary 7.3.1 is dual. Therefore, as desired, $\mathfrak{B}_1$ is not homogeneous. $\square$

Proposition 7.3.2 The boc $\mathfrak{B}_2$ given in MW2 of Theorem 5.6.1 is not homogeneous.

Proof. Consider the local boc $(\mathfrak{B}_1)_{\mathcal{P}}$ at $\mathcal{P}$, suppose we have the triangular formula (40) and the polynomial $\sigma_{\mathcal{P}}(\nu)$ of (41) in 5.4. Then the proof is similar to that of Proposition 7.3.1. $\square$

Example. Consider Example 2 of 2.1 and 4.2. A sequence of reductions for $a = (1), b = (\nu)$ yields an induced boc $\mathfrak{B}_2$: $\nu \begin{array}{c} \curvearrowright \\ \curvearrowright \end{array} \begin{array}{c} \mathcal{P} \\ \bullet \end{array} \xrightarrow{c} \begin{array}{c} \mathcal{Q} \\ \bullet \end{array}$ with two vertices $\mathcal{P}, \mathcal{Q}$ and $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\nu]$, $\Gamma'(\mathcal{Q}, \mathcal{Q}) = k$, $\delta(c) = 0$.

7.4 A critically non-homogeneous local boc $\mathfrak{B}_3$

Proposition 7.4.1 The boc $\mathfrak{B}_3$ given in MW3 of Theorem 5.6.1 is not homogeneous.

Proof. Consider

$$g_{ee}(\nu, \kappa) \in k[\nu, \kappa, \sigma_e(\nu)^{-1}, \sigma_e(\kappa)^{-1}]$$

in Formula (40) of 5.4, and $\sigma_e(\nu), \sigma(\nu)$ given in Formula (41). Suppose that we have item (1) of Corollary 7.3.1, i.e. $g_{ee}(\nu^0, \kappa^0) = 0$, $\sigma(\nu^0)\sigma_e(\kappa^0) \neq 0$ and ν^0 ranges over a cofinite subset D of k . Then we are able to present an infinite list of iso-classes of one dimensional objects $S_{\nu^0}, \forall \nu^0 \in D \setminus \text{therootsof}\sigma(\nu)$, such that

$$(S_{\nu^0})_{\mathcal{P}} = k, S_{\nu^0}(\nu) = (\nu^0). \text{ thus } S_{\nu^0}(b_l) = \emptyset, l = 1, \dots, j,$$

when we calculate S_{ν^0} according to Theorem 3.4.1. If $R(\mathfrak{B}_3)$ is homogeneous, there must be a cofinite subset $D_0 \subset D$ such that S_{ν^0} is homogeneous for any $\nu^0 \in D_0$. We fix such a ν^0 , denote S_{ν^0} by S for simplicity. Let $(e) : S \xrightarrow{\iota} E \xrightarrow{\pi} S$ be an almost split conflation of $R(\mathfrak{B}_3)$, then $E_{\mathcal{P}} = k^2$, and the eigenvalue of $E(\nu)$ is ν^0 , since the sequence

$$(e_0) : 0 \longrightarrow S \xrightarrow{\iota_0} E \xrightarrow{\pi_0} S \longrightarrow 0$$

is exact over $k[\nu, \sigma(\nu)^{-1}]$. Thus $E(b_l) = \emptyset, l = 1, \dots, j$, according to Theorem 3.4.1 and Proposition 4.5.1. We conclude that $E(\nu) = \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix}$, otherwise $\begin{pmatrix} \nu^0 & 0 \\ 0 & \nu^0 \end{pmatrix}$ would lead to (e) to be split.

We define an object $L \in R(\mathfrak{B}_3)$, such that $L_{\mathcal{P}} = k^2$,

$$L(\nu) = \begin{pmatrix} \nu^0 & 0 \\ 0 & \kappa^0 \end{pmatrix}, \quad L(b_e) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad L(b_l) = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

for $l \neq e$. And we also define a morphism $\varphi : L \rightarrow S$ given by $\varphi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\varphi(v) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ for any dotted arrow v . We first claim that φ is not a split epimorphism. In fact, if we have a morphism $\psi : S \rightarrow L$ with $\psi\varphi = id$, then $\psi_{\mathcal{P}} = (1, a)$, and

$$S(b_e)\psi_{\mathcal{P}} - \psi_{\mathcal{P}}L(b_e) = \psi(\delta(b_e)) = g_{ee}(S(\nu), L(\nu))\psi(u_e),$$

(see Formula (40) of 5.4). Thus, $-(1, a) \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = (g_{ee}(\nu^0, \nu^0)u_{e1}, g_{ee}(\nu^0, \kappa^0)u_{e2})$ or $(0, -1) = (*, 0)$, which results in a contradiction. Thus there exists some $\varphi' : L \rightarrow E$ such that $\varphi'\pi = \varphi$, whenever (e) is an almost split conflation. Then $\varphi'_{\mathcal{P}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ yields $\varphi'_{\mathcal{P}} = \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix}$ and $L(\nu)\varphi'_{\mathcal{P}} = \varphi'_{\mathcal{P}}E(\nu)$ yields

$$\begin{pmatrix} \nu^0 & 0 \\ 0 & \kappa^0 \end{pmatrix} \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix} = \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix} \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix},$$

i.e.

$$\begin{pmatrix} * & \nu^0 b \\ 0 & * \end{pmatrix} = \begin{pmatrix} * & 1 + b\nu^0 \\ 0 & * \end{pmatrix}$$


which is a contradiction.

The proof of κ^0 running over a cofinite subset of k is dual. Therefore, as desired, $\mathfrak{B}_3$ is not homogeneous. $\square$

Example. Consider Example 3 of 2.1 and 4.2, let $\alpha \neq 0, 1$. A sequence of reductions given by

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} \emptyset & \emptyset \\ \nu & \emptyset \end{pmatrix}, \quad D = \begin{pmatrix} d_3 & d_4 \\ d_1 & d_2 \end{pmatrix}$$

yields an induced local boc $\mathfrak{B}_3$, with a layer $L = (\Gamma; \omega; d_1, d_2, d_3, d_4; v, \dots)$, where $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\nu]$, $\delta(d_1) = v\nu - (\alpha\nu)v$, i.e. $e = 1$, $g_{11}(\nu, \kappa) = -\alpha\nu + \kappa$, which is not invertible in $k[\nu, \kappa]$.



7.5 A critically non-homogeneous local boc $\mathfrak{B}_4$

Proposition 7.5.1 The boc $\mathfrak{B}_4$ given in MW4 of Theorem 5.6.1 is not homogeneous.

Proof. Suppose we have the contrary. We fix a $\lambda^0 \in k$ with $h(\lambda^0) \neq 0$ (see Formula (43) in 5.4), and continue the reductions such that $a_1 = \emptyset, \dots, a_{i-1} = \emptyset$. Let $a_i = (\nu)$ be a parameter, then we obtain an induced boc $\mathfrak{B}'_4$ with a layer

$$L' = ((\Gamma)'; \omega'; b_1, \dots, b_j; u_1, \dots, u_m),$$

where the differentials $\delta(b_l)^0$ are given by Formula (40) and a polynomial $\sigma(\nu)$ is given by Formula (41) in Lemma 5.4.1. It is clear that $\mathfrak{B}'_4$ is minimal if we set $(\Gamma')'(\mathcal{P}, \mathcal{P}) = k[\nu, \sigma(\nu)^{-1}]$. Denote by ϑ the reduction functor from $R(\mathfrak{B}'_4)$ to $R(\mathfrak{B}_4)$. Define an infinite list of iso-classes of objects $\{S'_{\nu^0} | \forall \nu^0 \in k, \sigma(\nu^0) \neq 0\}$ in $R(\mathfrak{B}'_4)$, such that

$$(S'_{\nu^0})_{\mathcal{P}} = k, \quad S'_{\nu^0}(\nu) = (\nu^0) \quad S'_{\nu^0}(b_l) = \emptyset \text{ for } l = 1, 2, \dots, j.$$

Then we obtain an infinite list $\{S_{\nu^0} = \vartheta(S'_{\nu^0})\}$ in $R(\mathfrak{B}_4)$. If $R(\mathfrak{B}_4)$ is homogeneous, there must exist a cofinite subset $D_0 \subset k \setminus \{\text{the roots of } \sigma(\nu)\}$ such that for any $\nu^0 \in D_0$, S_{ν^0} is homogeneous. We fix such a $\nu^0 \in D_0$, and denote S_{ν^0} by S for simplicity. If $(e) : S \xrightarrow{\iota} E \xrightarrow{\pi} S$ is an almost split conflation of $R(\mathfrak{B}_4)$, there exists an almost split conflation $(e') : S' \xrightarrow{\iota'} E' \xrightarrow{\pi'} S'$ of $R(\mathfrak{B}'_4)$, such that $\vartheta(e')$ is equivalent to (e) by Theorem 7.2.1. Thus

$$E'_{\mathcal{P}} = k^2, \quad E'(\nu) = \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix}, \quad E'(b_l) = \emptyset, \quad l = 1, \dots, j$$

by Proposition 6.6.1. Let us go back to $R(\mathfrak{B}_4)$, then we have $S(\lambda) = (\lambda^0)$, $S(a_l) = \emptyset$; $E(\lambda) = \lambda^0 I_2$, $E(a_l) = \emptyset$, $l = 1, \dots, i-1$. Construct an object L in $R(\mathfrak{B}_4)$, such that

$$L_{\mathcal{P}} = k^2, \quad L(\lambda) = \begin{pmatrix} \lambda^0 & 1 \\ 0 & \lambda^0 \end{pmatrix}, \quad L(a_l) = 0 \text{ for } l = 1, \dots, i-1,$$

$$L(a_i) = \begin{pmatrix} \nu^0 & 0 \\ 0 & \nu^0 \end{pmatrix}, \quad L(b_l) = 0 \text{ for } l = 1, \dots, j,$$

and a morphism $\varphi : L \rightarrow S$ with $\varphi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\varphi(v) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ for any dotted arrow v . We first claim that φ is not a split epimorphism. In fact if $\psi : S \rightarrow L$ with $\psi\varphi = id$, then $\psi_{\mathcal{P}} = (1, a)$. But $S_{\nu^0}(\lambda)\psi_{\mathcal{P}} = \psi_{\mathcal{P}}L(\lambda)$, i.e. $\lambda^0(1, a) = (1, a) \begin{pmatrix} \lambda^0 & 1 \\ 0 & \lambda^0 \end{pmatrix}$, or $(\lambda^0, \lambda^0 a) = (\lambda^0, 1 + a\lambda^0)$, which is a contradiction. Thus, there is some morphism $\varphi' : L \rightarrow E$ such that $\varphi'_{\mathcal{P}}\pi = \varphi_{\mathcal{P}}$, whenever (e) is an almost split conflation of $R(\mathfrak{B}_4)$. Then $\varphi'_{\mathcal{P}}\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ leads to $\varphi'_{\mathcal{P}} = \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix}$, and $L(a_i)\varphi'_{\mathcal{P}} - \varphi'_{\mathcal{P}}E(a_i) = \varphi'(\delta(a_i)) = 0$ yields

$$\begin{pmatrix} \nu^0 & 0 \\ 0 & \nu^0 \end{pmatrix} \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix} = \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix} \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix}, \text{ i.e. } \begin{pmatrix} \nu^0 & \nu^0 b \\ 0 & \nu^0 c \end{pmatrix} = \begin{pmatrix} \nu^0 & 1 + b\nu^0 \\ 0 & c\nu^0 \end{pmatrix}.$$

It is a contradiction. Therefore, as desired, $\mathfrak{B}_4$ is not homogeneous. $\square$

Remark. The assumption $(\lambda - \mu)^2 \mid \tilde{h}(\lambda, \mu)$ is necessary in the proof. For instance, if $\tilde{h}(\lambda, \mu) = \lambda - \mu$, $\varphi'(\tilde{w}) = \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{pmatrix}$, then


$$\begin{aligned} \varphi'(\delta(a_i)) &= \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{pmatrix} \begin{pmatrix} \lambda^0 & 0 \\ 0 & \lambda^0 \end{pmatrix} - \begin{pmatrix} \lambda^0 & 1 \\ 0 & \lambda^0 \end{pmatrix} \begin{pmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \end{pmatrix} \\ &= \begin{pmatrix} -w_{21} & -w_{22} \\ 0 & 0 \end{pmatrix}; \end{aligned}$$

On the other hand,

$$\begin{aligned}\varphi'(\delta(a_i)) &= L(a_i)\varphi'_{\mathcal{P}} - \varphi'_{\mathcal{P}}E(a_i) \\ &= \begin{pmatrix} \nu^0 & 0 \\ 0 & \nu^0 \end{pmatrix} \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix} - \begin{pmatrix} 1 & b \\ 0 & c \end{pmatrix} \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix} \\ &= \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}\end{aligned}$$

which does not lead to any contradiction.

Example. Consider Example 1 of 2.1 and 4.2. A sequence of reductions given by $a = 1$, $b = \lambda$, yields an induced local boc $\mathfrak{B}_4$ with a layer $L = (\Gamma; \omega; c; v, \dots)$, where $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda]$, $\delta(c) = 0$.



7.6 An example in case of MW5

The minimally wild local boc given in MW5 of Theorem 5.6.1 may be homogeneous, even strongly homogeneous (see Example 4 of 6.5). Therefore a general wild bimodule problem does not necessarily have the non-homogeneous property. We must concentrate on some special set of bimodule problems, so called bipartite bimodule problems (see section 10), in which $P_1(\Lambda)$ are included. In this subsection, we will show an algebra, whose corresponding bimodule problem possess a reduction sequence with minimal size in case of MW5 of Theorem 5.6.1. We first prove two lemmas in order to treat the example and also the general case given in section 9. Following Auslander and Reiten [AR] we recall the following lemma.

Lemma 7.6.1 Let Γ be an algebra (possibly infinite-dimensional). Denote by $\Gamma\text{-mod}$ the category of finite dimensional Γ -modules. Suppose that

$$(e) \quad 0 \longrightarrow M \xrightarrow{\iota} E \xrightarrow{\pi} L \longrightarrow 0$$

is an almost split sequence in $\Gamma\text{-mod}$.

(1) If L_1 is a proper submodule of L , $E_1 = (L_1)\pi^{-1}$, then the sequence

$$0 \longrightarrow M \xrightarrow{\iota_1} E_1 \xrightarrow{\pi_1} L_1 \longrightarrow 0$$

is split, where $\pi_1 = \pi|_{E_1}$, $\iota_1 = \iota$.

(2) If a submodule $0 \neq K \subseteq M$, $M_2 = M/K$ is a quotient module of M , and $E_2 = E/\iota(K)$, then the sequence

$$0 \longrightarrow M_2 \xrightarrow{\iota_2} E_2 \xrightarrow{\pi_2} L \longrightarrow 0$$

is split, where ι_2, π_2 are induced morphisms from ι and π respectively.

Proof. (2) (given by Pan Jun) Consider the following commutative diagram having two exact rows in $\Gamma\text{-mod}$:

$$\begin{array}{ccccccccc} (e) : & 0 & \longrightarrow & M & \xrightarrow{\iota} & E & \xrightarrow{\pi} & L & \longrightarrow & 0 \\ & & & \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow id & & \\ & & & M/K & \xleftarrow{\psi} & E & & & & \\ & & & \nearrow id & \searrow \xi & & & & & \\ (e_2) : & 0 & \longrightarrow & M/K & \xrightarrow{\iota_2} & E/\iota(K) & \xrightarrow{\pi_2} & L & \longrightarrow & 0 \end{array}$$

where φ_1, φ_2 are natural projections respectively. Since φ_1 is not a split epimorphism and ι is a left almost split morphism, there exists a morphism $\psi : E \rightarrow M/K$, such that $\iota\psi = \varphi_1 id$. On the other hand, since the left square of the diagram determines a pushout, we have a morphism $\xi : E/\iota(K) \rightarrow M/K$, such that $\iota_2\xi = id$. Thus (e_2) is split as required.

(1) can be proved dually using pull back. □

Lemma 7.6.2 Let $\lambda \begin{array}{c} \curvearrowright \\ \curvearrowright \end{array} \begin{array}{c} \mathcal{P} \\ \mathcal{Q} \end{array} \xrightarrow{c} \bullet$ be a quiver, and $\Gamma = kQ$ be the path algebra. We define an exact sequence

$$(e) \quad 0 \longrightarrow S \xrightarrow{\iota} E \xrightarrow{\pi} S \longrightarrow 0$$

of $\Gamma\text{-mod}$, such that

$$\begin{aligned} S_{\mathcal{P}} &= k, S_{\mathcal{Q}} = k, S(\lambda) = (\lambda^0) \text{ for some } \lambda^0 \in k, S(c) = (1); \\ E_{\mathcal{P}} &= k^2, E_{\mathcal{Q}} = k^2, E(\lambda) = J_2(\lambda^0), E(c) = I_2; \\ \iota &= (\iota_{\mathcal{P}}, \iota_{\mathcal{Q}}) \text{ with } \iota_{\mathcal{P}} = \begin{pmatrix} 0 & 1 \end{pmatrix}, \iota_{\mathcal{Q}} = \begin{pmatrix} 0 & 1 \end{pmatrix}; \\ \pi &= (\pi_{\mathcal{P}}, \pi_{\mathcal{Q}}) \text{ with } \pi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \pi_{\mathcal{Q}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \end{aligned}$$

Then (e) is not an almost split sequence of $\Gamma\text{-mod}$.

Proof. Suppose that we have the contrary. Let us define three Γ -modules $K, \overline{S}, \overline{E}$ respectively as follows: $K_{\mathcal{P}} = (0)$, $K_{\mathcal{Q}} = k$; $\overline{S}_{\mathcal{P}} = k$, $\overline{S}_{\mathcal{Q}} = 0$, $\overline{S}(\lambda) = (\lambda^0)$; $\overline{E}_{\mathcal{P}} = k^2$, $\overline{E}_{\mathcal{Q}} = k$,

$$\overline{E}(\lambda) = \begin{pmatrix} \lambda^0 & 1 \\ 0 & \lambda^0 \end{pmatrix}, \quad \overline{E}(a_1) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Then $\overline{E}$ is indecomposable, and the commutative diagram below shows a contradiction to item (2) of Lemma 7.6.1.

$$\begin{array}{c}
 K : \begin{array}{ccccc}
 0 & \xrightarrow{0} & 0 & & \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & k & \xrightarrow{1} & k & \\
 \downarrow J_1(\lambda^0) & \downarrow J_2(\lambda^0) & \downarrow J_1(\lambda^0) & & \\
 (e) \quad k & \xrightarrow{(0 \ 1)} & k^2 & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & k \\
 \downarrow 1 & \downarrow 1 & \downarrow I_2 & \downarrow I_2 & \downarrow 1 \\
 & k & \xrightarrow{(0 \ 1)} & k^2 & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & k \\
 \downarrow J_1(\lambda^0) & \downarrow J_2(\lambda^0) & \downarrow J_1(\lambda^0) & & \\
 (e_2) \quad k & \xrightarrow{(0 \ 1)} & k^2 & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & k \\
 \downarrow 0 & \downarrow 0 & \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \downarrow 1 \\
 & 0 & \xrightarrow{0} & k & \xrightarrow{1} & k
 \end{array}
 \end{array}$$

Hence the assumption fails, (e) is not an almost split sequence of Γ -mod. $\square$

Proposition 7.6.1 Let $\mathfrak{B} = (\Gamma, \Omega)$ be a bocs with a layer $L = (\Gamma; \omega; a_1, \dots, a_n; v_1, \dots, v_m)$, where $\Gamma(\mathcal{P}, \mathcal{P}) = k[\lambda, g_{\mathcal{P}}(\lambda)^{-1}]$, $\delta(a_1) = 0$.

$$\lambda \curvearrowright \bullet \xrightarrow{a_1} \bullet \xrightarrow{\mathcal{Q}} \bullet$$

Then an edge reduction of $a_1 = (1)$ yields an induced local bocs $\mathfrak{B}'$. If $\mathfrak{B}'$ is minimal and tame, then $R(\mathfrak{B})$ is not homogeneous.

Proof. Suppose we have the contrary. Let $D = k \setminus \{\text{the roots of } \sigma(\lambda)\}$ be the domain of the minimal boc $\mathfrak{B}'$ where $\sigma(\lambda)$ is determined by Formula (41) of 5.4, and we use λ instead of ν . Then we define an infinite set of iso-classes of objects of dimension 1:

$$\{S'_{\lambda^0} \mid S'_{\lambda^0}(\lambda) = (\lambda^0), S'_{\lambda^0}(a_l) = \emptyset, l \geq 2, \forall \lambda^0 \in D\}$$

in $R(\mathfrak{B}'_5)$. Let $\vartheta : R(\mathfrak{B}') \rightarrow R(\mathfrak{B})$ be the reduction functor. We obtain also a set $\{S_{\lambda^0} = \vartheta(S'_{\lambda^0})\}$ in $R(\mathfrak{B})$, where

$$(S_{\lambda^0})_{\mathcal{P}} = k, (S_{\lambda^0})_{\mathcal{Q}} = k, (S_{\lambda^0})(\lambda) = (\lambda^0), (S_{\lambda^0})(a_1) = (1).$$

If $R(\mathfrak{B})$ is homogeneous, there must be a cofinite subset $D_0 \subset D$ such that for any $\lambda^0 \in D_0$, S_{λ^0} is homogeneous. We fix such a λ^0 , and denote S_{λ^0} by S for simplicity. If $(e) : S \xrightarrow{\iota} E \xrightarrow{\pi} S$ is an almost split conflation of $R(\mathfrak{B})$, then there exists an almost

split conflation $(e') : S' \xrightarrow{\iota'} E' \xrightarrow{\pi'} S'$ of $R(\mathfrak{B}')$ such that $\vartheta(e')$ is equivalent to (e) by Theorem 7.2.1. Thus

$$E'(\lambda) = \begin{pmatrix} \lambda^0 & 1 \\ 0 & \lambda^0 \end{pmatrix}, \quad E'(a_l) = \emptyset, \quad l \geq 2$$

by Proposition 6.6.1. Let us go back to $R(\mathfrak{B})$, then

$$E_{\mathcal{P}} = k^2, \quad E_{\mathcal{Q}} = k^2, \quad E(a_1) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

It is clear that $\iota_{\mathcal{P}} = (0 \ 1)$, $\iota_{\mathcal{Q}} = (0 \ 1)$, $\pi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\pi_{\mathcal{Q}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$.

Let Γ_0 be the path algebra of quiver $\lambda \begin{matrix} \curvearrowright \\ \curvearrowright \end{matrix} \xrightarrow{\mathcal{P}} \bullet \xrightarrow{a_1} \bullet \xrightarrow{\mathcal{Q}}$. We claim that (e) is also an almost split sequence in $\Gamma_0\text{-mod}$. In fact

(1) (e) is a non-split exact sequence in $\Gamma_0\text{-mod}$. And S is neither projective, nor injective.

(2) For any module L and morphism $\varphi : L \rightarrow S$ in $\Gamma_0\text{-mod}$, if we set $L(a_l) = 0$ for $l \geq 2$ and $\varphi(v_l) = 0$ for $l = 1, 2, \dots, m$, then $\varphi : L \rightarrow S$ can be regarded as a morphism of $R(\mathfrak{B})$.

(3) If φ is not a split epimorphism in $\Gamma_0\text{-mod}$, then it is not a split epimorphism in $R(\mathfrak{B})$.

(4) There exists some $\varphi' : L \rightarrow E$ with $\varphi'\pi = \varphi$ in $R(\mathfrak{B})$, whenever (e) is an almost split conflation of $R(\mathfrak{B})$. Thus

$$L(\lambda)\varphi'_{\mathcal{P}} = \varphi'_{\mathcal{P}}E(\lambda) \quad \text{and} \quad L(a_1)\varphi'_{\mathcal{Q}} = \varphi'_{\mathcal{P}}E(a_1),$$

i.e. $\varphi'_0 = (\varphi'_{\mathcal{P}}, \varphi'_{\mathcal{Q}})$ is a morphism in $\Gamma_0\text{-mod}$ with $\varphi'_0\pi = \varphi$. Therefore $E \xrightarrow{\pi} S$ is a right almost split morphism in $\Gamma_0\text{-mod}$. The claim is proved.

Therefore (e) is also an almost split sequence of $\Gamma_0\text{-mod}$, which leads to a contradiction to Lemma 7.6.2. $\square$


Example. Consider Example 3 of 2.1 and 4.3. Let

$$\underline{n} = (2, 2, 2, 2, 2; 2, 2, 2, 2, 2),$$

$(N, R, \underline{n})$ be given in Formula (6) of 2.6 and we denote $\overline{a^*}, \overline{b^*}, \overline{c^*}, \overline{d^*}$ by A, B, C, D respectively for simplicity. Then a sequence of reductions given by

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} c_3 & c_4 \\ \lambda & c_2 \end{pmatrix}, \quad D = \begin{pmatrix} d_3 & d_4 \\ d_1 & d_2 \end{pmatrix}$$

yields a local layered boc. $\delta(c_2)^0, \delta(c_3)^0, \delta(c_4)^0$ and $\delta(d_1)^0$ satisfy Formula (42) of 5.4.



If we continue the reductions given by $c_2 = \emptyset$, $c_3 = \emptyset$, $c_4 = \emptyset$, then $\delta(d_1) = \tilde{w}\lambda - \lambda\tilde{w}$ with $\tilde{h}(\lambda, \mu) = -(\lambda - \mu)$ in Formula (42) and $h(\lambda) = 1$ in Formula (43). Furthermore, fix any $\lambda^0 \in k$, and denote a_1 by ν , then $\delta(d_2)^0$, $\delta(d_3)^0$, $\delta(d_4)^0$ satisfy Formula (40) with $g_l(\nu, \kappa) \in k[\nu, \kappa]$ being invertible for $l = 2, 3, 4$. Thus we obtain a sequence of parameterized bimodule problems with the end term in case of MW5.


Define a new size vector

$$\tilde{\underline{n}} = (2, 2, 2, 2, 2; 3, 3, 3, 3, 3),$$

let $(\tilde{N}, \tilde{R}, \tilde{\underline{n}})$ be given in Formula (6) of 2.6. Then a sequence of reductions given by

$$\tilde{A} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \tilde{B} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad \tilde{C} = \begin{pmatrix} \emptyset & \emptyset & \emptyset \\ 0 & \lambda & \emptyset \end{pmatrix}, \quad \tilde{D} = \begin{pmatrix} d'_0 & d_3 & d_4 \\ d_0 & d_1 & d_2 \end{pmatrix}$$

yields an induced boc $\mathfrak{B}$ with a layer $L = (\Gamma'; \omega; d_0, d_1, d_2, d'_0, d_3, d_4; v, \dots)$, where $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda]$, $\delta(d_0) = 0$.



If we set $d_0 = 1$, then the induced boc $\mathcal{B}'$ is minimal. Thus $\mathcal{B}$ and $\mathcal{B}'$ in our example satisfy the hypothesis of Proposition 7.6.1, and the structure given in Lemma 7.6.2 is as follows:

$$\begin{array}{llll} & A & B & C & D \\ S & \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} & \begin{pmatrix} \emptyset & \emptyset & \emptyset \\ 0 & \lambda^0 & \emptyset \end{pmatrix} & \begin{pmatrix} \emptyset & \emptyset & \emptyset \\ 1 & \emptyset & \emptyset \end{pmatrix} \\ E & \begin{pmatrix} 0_{2 \times 2} & I_2 & 0 \\ 0_{2 \times 2} & 0 & I_2 \end{pmatrix} & \begin{pmatrix} 0_{2 \times 2} & 0 & I_2 \\ 0_{2 \times 2} & 0 & 0 \end{pmatrix} & \begin{pmatrix} \emptyset_{2 \times 2} & \emptyset & \emptyset \\ 0_{2 \times 2} & J_2(\lambda^0) & \emptyset \end{pmatrix} & \begin{pmatrix} \emptyset & \emptyset & \emptyset \\ I_2 & \emptyset & \emptyset \end{pmatrix} \\ \bar{S} & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} \emptyset & \emptyset \\ \lambda^0 & \emptyset \end{pmatrix} & \begin{pmatrix} \emptyset & \emptyset \\ 0 & \emptyset \end{pmatrix} \\ \bar{E} & \begin{pmatrix} 0_{2 \times 1} & I_2 & 0 \\ 0_{2 \times 1} & 0 & I_2 \end{pmatrix} & \begin{pmatrix} 0_{2 \times 1} & 0 & I_2 \\ 0_{2 \times 1} & 0 & 0 \end{pmatrix} & \begin{pmatrix} \emptyset_{2 \times 1} & \emptyset & \emptyset \\ 0_{2 \times 1} & J_2(\lambda^0) & \emptyset \end{pmatrix} & \begin{pmatrix} \emptyset & \emptyset & \emptyset \\ \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \emptyset & \emptyset \end{pmatrix} \end{array}$$

where $\bar{E}$ is indecomposable.

8 One-sided differentials

8.1 Matrix problems

Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem of Definition 2.2.1 with a triangular basis (A, B) of Definition 2.4.1. Suppose that $\rho_{\mathcal{I}_1 \mathcal{J}_1}^{w_1} \in A$, $\mathcal{M}'$ is a $\mathcal{K}$ - $\mathcal{K}$ -subbimodule of $\mathcal{M}$ generated by

$$\{\rho_{\mathcal{I} \mathcal{J}}^w \in A \mid \rho_{\mathcal{I} \mathcal{J}}^w \succ \rho_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}\}$$

(see Proposition 2.4.1). Then $\mathcal{M}/\mathcal{M}'$ is a $\mathcal{K}$ - $\mathcal{K}$ -quotient bimodule of $\mathcal{M}$. And $(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$ is called a *partial bimodule problem* of $(\mathcal{K}, \mathcal{M}, H)$, where $d : \mathcal{K} \rightarrow \mathcal{M}/\mathcal{M}'$ is a derivation given by $d(S) = (SH - HS) + \mathcal{M}'$ for any $S \in \mathcal{K}$. Thus $(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$ belongs to the second set of 1.4, but may not satisfy Definition 2.2.1. Moreover, $(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$ corresponds to a boc $\mathfrak{B}$ given in [CB2]. Let $\mathfrak{A} = (\Gamma, \Omega)$ be the corresponding boc of $(\mathcal{K}, \mathcal{M}, H)$ with a layer

$$L = (\Gamma'; \omega; a_1, \dots, a_n; v_1, \dots, v_m),$$

$\overline{\Gamma}$ be freely generated by $a_1, \dots, a_{n_1}$ over Γ' , when $a_{n_1} = (\rho_{\mathcal{I}_1 \mathcal{J}_1}^{w_1})^*$. Since

$$\delta_2(v_i) = \sum_{j, l < i} c_{jl}^i v_j \otimes v_l$$

with the coefficients $c \in k$, which are independent of $a_1, \dots, a_n$ by Formula (25) of 3.6, Ω is also a $\overline{\Gamma}$ - $\overline{\Gamma}$ -coalgebra. Then $\mathfrak{B} = (\overline{\Gamma}, \Omega)$ has a layer

$$\overline{L} = (\Gamma'; \omega; a_1, \dots, a_{n_1}; v_1, \dots, v_m),$$

which is called a *partial boc* of $\mathfrak{A}$. For their representation categories, we have

$$Mat(\mathcal{K}, \mathcal{M}/\mathcal{M}') = \{M \in Mat(\mathcal{K}, \mathcal{M}) \mid M_{p_{\mathcal{I} \mathcal{J}}^w q_{\mathcal{I} \mathcal{J}}^w} = 0, \forall (p_{\mathcal{I} \mathcal{J}}^w, q_{\mathcal{I} \mathcal{J}}^w) \succ (p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}, q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1})\},$$

and

$$R(\mathfrak{B}) = \{\overline{M} \in R(\mathfrak{A}) \mid M(a_j) = 0, \forall j > n_1\}.$$

Both of them have exact structures inherited from $Mat(\mathcal{K}, \mathcal{M})$ and $R(\mathfrak{A})$ respectively. In order to treat with their reductions, we define another algebraic structure, so called matrix problem.

Definition 8.1.1 A *matrix problem* $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ consists of the following datum:

- I'. A set of integers $T = \{1, 2, \dots, t\}$ and an equivalent relation $\sim$ on T . Two subsets $T_1, T_2 \subset T$, which have t_1, t_2 elements respectively, and $\sim_1 = \sim|_{T_1}$, $\sim_2 = \sim|_{T_2}$.

II'. Two upper triangular matrix algebras

$$\mathcal{K}_1 = \{(s_{ij}^1)_{t_1 \times t_1}\} \text{ and } \mathcal{K}_2 = \{(s_{ij}^2)_{t_2 \times t_2}\},$$

such that

$$s_{ii}^1 = s_{jj}^1, s_{ii}^2 = s_{jj}^2, s_{ii}^1 = s_{jj}^2$$

when $i \sim j$, and a set of matrices $\mathcal{V} = \{(v_{ij})_{t_1 \times t_2}\}$, s_{ij}^1, s_{ij}^2 for $i < j$ and v_{ij} satisfy the following equation system:

$$\sum_{\mathcal{I} \ni i < j \in \mathcal{J}} (c_{ij}^{1l} x_{ij}^1 + c_{ij}^{2l} x_{ij}^2) + \sum_{(i', j') \in \mathcal{I} \times \mathcal{J}} c_{i'j'}^{3l} x_{i'j'}^3 = 0,$$

which are indexed by $1 \leq l \leq q_{\mathcal{I}\mathcal{J}}$ for some $q_{\mathcal{I}\mathcal{J}} \in \mathbb{N}$, and for each pair

$$(\mathcal{I}, \mathcal{J}) \in (T_1 / \sim_1) \times (T_2 / \sim_2).$$

III'. A $\mathcal{K}_1$ - $\mathcal{K}_2$ -bimodule $\mathcal{M} = \{(m_{ij})_{t_1 \times t_2}\}$, where m_{ij} satisfy the equation system:

$$\sum_{(i,j) \in \mathcal{I} \times \mathcal{J}} d_{ij}^l z_{ij} = 0$$

which are indexed by $1 \leq l \leq q'_{\mathcal{I}\mathcal{J}}$ for some $q'_{\mathcal{I}\mathcal{J}} \in \mathbb{N}$, and for each pair

$$(\mathcal{I}, \mathcal{J}) \in (T_1 / \sim_1) \times (T_2 / \sim_2).$$

□

Remark We stress that the matrix problem given in Definition 8.1.1 is not necessarily a bimodule problem, since the derivation is not defined. But it corresponds to a bocs. □

We also have a basis $A'_{\mathcal{I}\mathcal{J}} = \{\chi_{\mathcal{I}\mathcal{J}}^w\}$ and A' of the solution space of equation system III' similar to Formula (1) of 2.3. A basis of the solution space of equation system II' for each pair $(\mathcal{I}, \mathcal{J})$ is denoted by $B'_{\mathcal{I}\mathcal{J}}$. And B' is given similarly to Formula (2) of 2.3. It is clear that $\text{rad}\mathcal{K}_1$, $\text{rad}\mathcal{K}_2$ and $\mathcal{V}$ are subspaces of the solution space of the equation system II'. We may choose basis of $\text{rad}\mathcal{K}_1$, $\text{rad}\mathcal{K}_2$, and $\mathcal{V}$ respectively as follows. First let

$$W_0 = \text{rad}\mathcal{K}_1 \cap \text{rad}\mathcal{K}_2 \cap \mathcal{V},$$

take a basis $\{\xi_{\mathcal{I}\mathcal{J}}^{v_{123}}\}$ of W_0 . Secondly, let W_{12} , W_{23} , W_{31} be the complement spaces of W_0 in

$$\text{rad}\mathcal{K}_1 \cap \text{rad}\mathcal{K}_2, \text{rad}\mathcal{K}_2 \cap \mathcal{V}, \mathcal{V} \cap \text{rad}\mathcal{K}_1,$$

and take basis $\{\xi_{\mathcal{IJ}}^{v_{12}}\}, \{\xi_{\mathcal{IJ}}^{v_{23}}\}, \{\xi_{\mathcal{IJ}}^{v_{31}}\}$ of W_{12}, W_{23}, W_{31} respectively. Thirdly, let W_1, W_2, W_3 be the complement spaces of

$$W_0 \oplus W_{12} \oplus W_{31}, W_0 \oplus W_{23} \oplus W_{12}, W_0 \oplus W_{31} \oplus W_{23}$$

in $\text{rad}\mathcal{K}_1, \text{rad}\mathcal{K}_2, \mathcal{V}$, and take basis $\{\xi_{\mathcal{IJ}}^{v_1}\}, \{\xi_{\mathcal{IJ}}^{v_2}\}, \{\xi_{\mathcal{IJ}}^{v_3}\}$ of W_1, W_2, W_3 respectively. Thus the dual basis

$$\{(\xi_{\mathcal{IJ}}^{v_{123}})^*, (\xi_{\mathcal{IJ}}^{v_{12}})^*, (\xi_{\mathcal{IJ}}^{v_{23}})^*, (\xi_{\mathcal{IJ}}^{v_{31}})^*, (\xi_{\mathcal{IJ}}^{v_1})^*, (\xi_{\mathcal{IJ}}^{v_2})^*, (\xi_{\mathcal{IJ}}^{v_3})^*\}$$

can be regarded as the coefficient functions. Write

$$\begin{aligned} \{\xi_{\mathcal{IJ}}^{v'_1}\} &= \{\xi_{\mathcal{IJ}}^{v_{123}}, \xi_{\mathcal{IJ}}^{v_{12}}, \xi_{\mathcal{IJ}}^{v_{31}}, \xi_{\mathcal{IJ}}^{v_1}\}, \\ \{\xi_{\mathcal{IJ}}^{v'_2}\} &= \{\xi_{\mathcal{IJ}}^{v_{123}}, \xi_{\mathcal{IJ}}^{v_{12}}, \xi_{\mathcal{IJ}}^{v_{23}}, \xi_{\mathcal{IJ}}^{v_2}\}, \\ \{\xi_{\mathcal{IJ}}^{v'_3}\} &= \{\xi_{\mathcal{IJ}}^{v_{123}}, \xi_{\mathcal{IJ}}^{v_{23}}, \xi_{\mathcal{IJ}}^{v_{31}}, \xi_{\mathcal{IJ}}^{v_3}\}. \end{aligned}$$

We can define the representation category $\text{Mat}(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M})$ of a matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ similarly to 2.2. Given any size vector $\underline{n}_1$ of $(T_1, \sim_1)$, $\underline{n}_2$ of $(T_2, \sim_2)$, such that $(n_1)_i = (n_2)_i$ for any $i \in T_1 \cap T_2$, a triple $(N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2)$ is defined similarly to Formula (6) of 2.6, such that N, R_1, R_2 and V are $\underline{n}_1 \times \underline{n}_2$, $\underline{n}_1 \times \underline{n}_1$, $\underline{n}_2 \times \underline{n}_2$ and $\underline{n}_1 \times \underline{n}_2$ partitioned matrices respectively. More precisely,

$$\begin{aligned} N &= \sum_{(\mathcal{I}, \mathcal{J}) \in (T_1/\sim_1) \times (T_2/\sim_2)} \sum_{w=1}^{r_{\mathcal{IJ}}} \overline{(\chi_{\mathcal{IJ}}^w)^*} \otimes \chi_{\mathcal{IJ}}^w \\ R_1 &= \sum_{\mathcal{I} \in T_1/\sim_1} \overline{E_{\mathcal{I}}^*} \otimes E_{\mathcal{I}} + \sum_{(\mathcal{I}, \mathcal{J}) \in (T_1/\sim_1) \times (T_1/\sim_1)} \sum_{v'_1=1}^{r_{\mathcal{IJ}}^1} \overline{(\xi_{\mathcal{IJ}}^{v'_1})^*} \otimes \xi_{\mathcal{IJ}}^{v'_1} \\ R_2 &= \sum_{\mathcal{I} \in T_2/\sim_2} \overline{E_{\mathcal{I}}^*} \otimes E_{\mathcal{I}} + \sum_{(\mathcal{I}, \mathcal{J}) \in (T_2/\sim_2) \times (T_2/\sim_2)} \sum_{v'_2=1}^{r_{\mathcal{IJ}}^2} \overline{(\xi_{\mathcal{IJ}}^{v'_2})^*} \otimes \xi_{\mathcal{IJ}}^{v'_2} \\ V &= \sum_{(\mathcal{I}, \mathcal{J}) \in (T_1/\sim_1) \times (T_2/\sim_2)} \sum_{v'_3=1}^{r_{\mathcal{IJ}}^3} \overline{(\xi_{\mathcal{IJ}}^{v'_3})^*} \otimes \xi_{\mathcal{IJ}}^{v'_3} \end{aligned}$$

Then we start a reduction according to the matrix equation

$$R_1 N = V + N R_2, \quad \text{or} \quad X_{pp} N_{pq} = V_{pq} + N_{pq} X_{qq},$$

in the same way as in 2.6, where the second equation is the (p, q) -block of the first one.

Regularization. If the equation $x_{pq}^3 = 0$ is not a linear combination of the equations in equation system II' , let $\overline{N}_{pq} = \emptyset$. Then we add $x_{pq}^3 = 0$ into the equation system II' . Let R'_1, R'_2, V' be the restrictions of R_1, R_2, V given by $x_{pq}^3 = 0$ respectively.

Suppose now that $x_{pq}^3 = 0$ is a linear combination of the equations in equation system II'.

Edge reduction. If $\mathcal{P} \neq \mathcal{Q}$, let $\overline{N}_{pq} = \begin{pmatrix} 0 & I \\ 0 & 0 \end{pmatrix}$.

Loop reduction. If $\mathcal{P} = \mathcal{Q}$, let $\overline{N}_{pq} = W$ for some Weyr matrix W .

In the last two cases, R_1 and R_2 are restricted by the equation

$$X_{pp}\overline{N}_{pq} = \overline{N}_{pq}X_{qq}.$$

Denote by R'_1 and R'_2 the restricted algebras of R_1 and R_2 respectively, and

$$V' = V - R'_1(\overline{N}_{pq} \otimes \chi) + (\overline{N}_{pq} \otimes \chi)R'_2.$$

where χ is the first basis element of A' .

Let

$$N' = N - N_{pq} \otimes \rho$$

in all the 3 cases, then we obtain a new triple $(N', R'_1 \times R'_2, \underline{n}'_1 \times \underline{n}'_2)$ similarly to 2.6. Consequently we also obtain a new matrix problem $(\mathcal{K}'_1 \times \mathcal{K}'_2, \mathcal{M}', \mathcal{V}')$ with a new index set $(T', \sim')$ similarly to 3.1. Finally we have a matrix equation over $(\mathcal{K}'_1 \times \mathcal{K}'_2, \mathcal{M}', \mathcal{V}')$:

$$R'_1 N' = V' + N' R'_2.$$

Inductively we have a reduction sequence:

$$(\triangle) \quad (N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2), \dots, (N^r, R_1^r \times R_2^r, \underline{n}_1^r \times \underline{n}_2^r), \dots, (N^s, R_1^s \times R_2^s, \underline{n}_1^s \times \underline{n}_2^s).$$

The notions of parametrization, and free parametrization are still valid for matrix problems.

Proposition 8.1.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem. Then $(\mathcal{K} \times \mathcal{K}, \mathcal{M}, \mathcal{V})$ is a matrix problem, where $T_1 = T$, $T_2 = T$, $\mathcal{V} = \{V = SH - HS \mid \forall S \in \mathcal{K}\}$.

Proof. If $V = (v_{ij})_{t \times t}$, then

$$v_{ij} = \sum_{l=i+1}^t s_{il}h_{lj} - \sum_{l=1}^{j-1} h_{il}s_{lj}.$$

In fact $s_{ii}h_{ij} - h_{ij}s_{jj} = 0$ in both cases $i \sim j$ and $i \approx j$, and $v_{ij} = 0$ when $(i, j) \prec (p, q)$. Then we obtain the required equation system II'. $\square$

Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, and $\rho_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}$ be given in the beginning of the subsection, (p, q) be defined in 2.4. Suppose in addition that if

$$(p_{\mathcal{I} \mathcal{J}}^w, q_{\mathcal{I} \mathcal{J}}^w) \succ (p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}, q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}), \text{ then } p_{\mathcal{I} \mathcal{J}}^w < p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}.$$

Let $\mathcal{K}_1$ be a subalgebra of $\mathcal{K}$ consisting of the submatrices of the matrices of $\mathcal{K}$ with the (i, j) -th entries for

$$p \geq i, j \geq p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1};$$

$\mathcal{K}_2$ be that for

$$q \leq i, j \leq q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1};$$

let $\overline{\mathcal{M}}$ consist of the submatrices of the matrices of $\mathcal{M}$ with the (i, j) -entries for

$$p \geq i \geq p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1} \quad \text{and} \quad q \leq j \leq q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1};$$

and $\mathcal{V}$ consist of those of $\{SH - HS \mid \forall S \in \text{rad}\mathcal{K}\}$.

Proposition 8.1.2 $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$ defined above is a matrix problem, whose representation category $\text{Mat}(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}})$ is equivalent to $\text{Mat}(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$, as well as $R(\mathfrak{B})$.

Proof. I'. $T_1 = \{p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}, \dots, p\}$, $T_2 = \{q, \dots, q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}\}$.

II'. The equation system is given by Formula (18) of 3.6:

$$\begin{cases} x_{ij}^1 = \sum_w (\zeta_{\mathcal{I}\mathcal{J}}^w)^* b_{\mathcal{I}\mathcal{J}}^w(i, j) & \text{for } p \geq i, j \geq p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}; \\ x_{ij}^2 = \sum_w (\zeta_{\mathcal{I}\mathcal{J}}^w)^* b_{\mathcal{I}\mathcal{J}}^w(i, j) & \text{for } q \leq i, j \leq q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}; \\ x_{ij}^3 = \sum_l \left(\sum_w (\zeta_{\mathcal{I}\mathcal{L}}^w)^* b_{\mathcal{I}\mathcal{L}}^w(i, l) h_{lj} - \right. \\ \quad \left. h_{il} \sum_w (\zeta_{\mathcal{L}\mathcal{J}}^w)^* b_{\mathcal{L}\mathcal{J}}^w(l, j) \right) & \text{for } p \geq i \geq p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}, q \leq j \leq q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}. \end{cases}$$

III'. $\overline{\mathcal{M}}$ is a $\mathcal{K}_1$ - $\mathcal{K}_2$ -bimodule, and the equation system is given by Formula (19) of 3.6:

$$z_{ij} = \sum_w (\rho_{\mathcal{I}\mathcal{J}}^w)^* a_{\mathcal{I}\mathcal{J}}^w(i, j) \quad \text{for } p \geq i \geq p_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}, q \leq j \leq q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}. \quad \square$$

Corollary 8.1.1 Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem, $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$ be the matrix problem given by a partial bimodule problem $(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$. Suppose $(N_0, R, \underline{n})$ is a triple of $(\mathcal{K}, \mathcal{M}, H)$, and $(N, R_1 \times R_2, V)$ is the corresponding triple of $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$, such that $\forall i \in T_1, (n_1)_i = n_i$, and $\forall i \in T_2, (n_2)_i = n_i$. Then the reduction sequence (Δ) of $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$ starting from $(N, R_1 \times R_2, V)$ determines a reduction sequence $(*)$ of $(\mathcal{K}, \mathcal{M}, H)$ starting from $(N_0, R, \underline{n})$.

Proof. We use induction on the reduction step r . The case of $r = 0$ is clear.

Suppose that we have already a reduction sequence $(*)$ up to $(N_0^r, R^r, \underline{n}^r)$ determined by (Δ) , and χ^r is the first basis element of $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \overline{\mathcal{M}}^r, \mathcal{V}^r)$, ρ^r is that of $(\mathcal{K}^r, \mathcal{M}^r, H^r)$, then $\chi^r = \rho^r \mid_{\overline{\mathcal{M}}^r}$.

For step $(r + 1)$, let $\overline{N}_{p_r q_r}^r$ be obtained at the $(r + 1)$ -th reduction of (Δ) . Then the size vector $\underline{n}^{r+1}$ is given by $\underline{n}^r$ and $\overline{N}_{p_r q_r}^r$ according to 3.1; $H_{\underline{n}^{r+1}}^{r+1} = H_{\underline{n}^r}^r + \overline{N}_{p_r q_r}^r \otimes \rho^r$; $N_0^{r+1} = N_0^r - N_{p_r q_r}^r \otimes \rho^r$; and R^{r+1} is a restriction of R^r given in Formula (10) of 2.6. $\square$

8.2 One-sided differentials I

Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem with a triangular basis (A, B) . Let $\rho_{\mathcal{I}_1 \mathcal{J}_1}^w \in A$ be given in the beginning of 8.1, and $\rho \in A$ be the first basis element given in 2.4. We assume that

$$p_{\mathcal{I}_1 \mathcal{J}_1}^w = p,$$

and if

$$(p_{\mathcal{I} \mathcal{J}}^w, q_{\mathcal{I} \mathcal{J}}^w) \succ (p, q_{\mathcal{I}_1 \mathcal{J}_1}^{w_1}), \quad \text{then } p_{\mathcal{I} \mathcal{J}}^w < p.$$

Thus the partial bimodule problem $(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$ corresponds to a partial boc $\mathfrak{B}$, and determines a matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$, such that

$$T_1 = \{1\}, \quad T_2 = \{1, 2, \dots, t\};$$

and

$$\mathcal{K}_1 = \{(s)\}, \quad \mathcal{K}_2 = \{S = (s_{ij})_{t \times t}\},$$

where $s_{ij} = 0$ if $i > j$, $s_{ii} = s_{jj}$ if $i \sim j$, and $s_{ii} = s$ if $i \sim 1$,

$$\mathcal{V} = \{V = (v_1 \ v_2 \ \cdots \ v_t)\},$$

moreover s_{ij} ($i < j$) and v_j satisfy equation system II' of Definition 8.1.1;

$$\overline{\mathcal{M}} = \{M = (\cdots \ a \ \cdots \ b \ \cdots \ a \ \cdots \ b \ \cdots)_{1 \times t}\},$$

and the entries of M satisfy equation system III'.

Therefore the partial boc $\mathfrak{B}$ has a partial layer

$$L^p = (\Gamma'; \omega; a_1, \dots, a_n; b_1, \dots, b_m; u, v, w, q),$$


where the vertices $\mathcal{P}, \mathcal{I}_1, \dots, \mathcal{I}_r$ are all trivial; solid arrows

$$a : \mathcal{P} \rightarrow \mathcal{I}, \quad b : \mathcal{P} \rightarrow \mathcal{P};$$

and dotted arrows

$$u : \mathcal{P} \rightarrow \mathcal{P}, \quad v : \mathcal{P} \rightarrow \mathcal{I}, \quad w : \mathcal{I} \rightarrow \mathcal{I}, \quad q : \mathcal{I} \rightarrow \mathcal{P}.$$

Note that a, b stand for general solid arrows, and u, v, w, q stand for general dotted arrows.



(46)

It is obvious, that the summands of $\delta(a_i)$ involve only the monomials: (1) v , (2) $b^e v$, (3) aw ; and those of $\delta(b)$ involve: (1) u , (2) $b^e u$, (3) aq for some positive integer e . Such differentials of solid arrows are said to be *one-sided*.

Now we first consider the local partial boc $\mathfrak{B}_{\mathcal{P}}$ at $\mathcal{P}$ obtained from $\mathfrak{B}$ by deleting the vertices $\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_r$. Thus $\mathfrak{B}_{\mathcal{P}} = (\Gamma_{\mathcal{P}}, \Omega_{\mathcal{P}})$ has a partial layer

$$L_{\mathcal{P}}^p = (\Gamma'_{\mathcal{P}}; \omega_{\mathcal{P}}; b_1, \dots, b_m; u)$$

and the summands of $\delta(b_j)$ involve only the monomials of u or $b^e u$. Then we have the following possibilities.

$$\mathbf{P1} \quad \begin{cases} \delta(b_1)^0 &= g_{11}u_1 \\ \delta(b_2)^0 &= g_{21}u_1 + g_{22}u_2 \\ \dots &\dots \\ \delta(b_m)^0 &= g_{m1}u_1 + g_{m2}u_2 + \dots + g_{m,m}u_m \end{cases}$$

where $u_1, u_2, \dots, u_m$ are linearly independent, $g_{ll'} \in k$, $l \leq l'$, $g_{ll} \neq 0$. Then $\mathfrak{B}_{\mathcal{P}}$ is of finite type.

Otherwise we have the following formula.

$$\begin{cases} \delta(b_1)^0 &= g_{11}u_1 \\ \delta(b_2)^0 &= g_{21}u_1 + g_{22}u_2 \\ \dots &\dots \\ \delta(b_{j-1})^0 &= g_{j-1,1}u_1 + g_{j-1,2}u_2 + \dots + g_{j-1,j-1}u_{j-1} \\ \delta(b_j)^0 &= g_{j1}u_1 + g_{j2}u_2 + \dots + g_{j,j-1}u_{j-1} \end{cases} \quad (47)$$

where $u_1, u_2, \dots, u_{j-1}$ are linearly independent, $g_{ll'} \in k$, $l \leq l'$, $g_{ll} \neq 0$. If we set $b_l = \emptyset$, $u_l = 0, l = 1, \dots, j-1$, $b_j = \lambda$, then

$$\begin{cases} \delta(b_{j+1})^0 &= h_{j+1,j+1}(\lambda)u_{j+1} \\ \delta(b_{j+2})^0 &= h_{j+2,j+1}(\lambda)u_{j+1} + h_{j+2,j+2}(\lambda)u_{j+2} \\ \dots &\dots \\ \delta(b_{m'})^0 &= h_{m',j+1}(\lambda)u_{j+1} + h_{m',j+2}(\lambda)u_{j+2} + \dots + h_{m',m'}(\lambda)u_{m'} \end{cases} \quad (48)$$

where $u_{j+1}, \dots, u_{m'}$ are linearly independent, $h_{ll}(\lambda) \neq 0$.

P2 $m' = m$, $h_{ll}(\lambda)$ are all non-zero constants. Then $\mathfrak{B}_{\mathcal{P}}$ is of tame type.

P3 $m' = m$, $h_{11}(\lambda), \dots, h_{e-1,e-1}(\lambda)$ are non-zero constants but $h_{e,e}(\lambda)$ is not a constant. Then $\mathfrak{B}_{\mathcal{P}}$ is of wild type.

P4 $m' < m$, $\mathfrak{B}_{\mathcal{P}}$ is of wild type.

8.3 One-sided differentials II

Let us go back to the partial boc $\mathfrak{B}$ given at the beginning of 8.2. Suppose that the local partial boc $\mathfrak{B}_{\mathcal{P}}$ is in case P2 of 8.2. If

$$a_1 \prec \dots \prec a_{n_0} \prec b_j \prec a_{n_0+1},$$

then we have either a triangular formula:

$$\begin{cases} \delta(a_1)^0 &= f_{11}v_1 \\ \delta(a_2)^0 &= f_{21}v_1 + f_{22}v_2 \\ \dots &\dots \\ \delta(a_{n_0})^0 &= f_{n_01}v_1 + f_{n_02}v_2 + \dots + f_{n_0n_0}v_{n_0} \end{cases}$$

where $v_1, v_2, \dots, v_{n_0}$ are linearly independent, and $f_{ll'} \in k$, $f_{ll} \neq 0$, or we have a triangular formula:

$$\begin{cases} \delta(a_1)^0 &= f_{11}v_1 \\ \dots &\dots \\ \delta(a_{n(1)-1})^0 &= f_{n(1)-1,1}v_1 + \dots + f_{n(1)-1,n(1)-1}v_{n(1)-1} \\ \delta(a_{n(1)})^0 &= f_{n(1),1}v_1 + \dots + f_{n(1),n(1)-1}v_{n(1)-1} \end{cases} \quad (49)$$

where $v_1, v_2, \dots, v_{n(1)-1}$ are linearly independent, and $f_{ll'} \in k$, $f_{ll} \neq 0$. Set $a_l = \emptyset$, $v_l = 0$, then we obtain an induced boc by a series of regularizations. We continue the procedure, then obtain a sequence of positive integers $n(1), n(2), \dots, n(i)$, and finally reach to the first formula by induction. Let

$$I = \{1, 2, \dots, n_0\} \setminus \{n(1), n(2), \dots, n(i)\} \quad (50)$$

Let us write $\bar{a}_l = a_{n(l)}$, $l = 1, \dots, i$, $\bar{b} = b_j$ for simplicity. The differentials of $\bar{b}$ with respect to $\bar{a}_1, \dots, \bar{a}_i$ is either

$$\begin{aligned} \delta(\bar{b})^0 &= \bar{a}_1(h_{11}q_1) \\ &\quad + \bar{a}_2(h_{21}q_1 + h_{22}q_2) \\ &\quad + \dots \dots \\ &\quad + \bar{a}_i(h_{i1}q_1 + h_{i2}q_2 + \dots + h_{ii}q_i) \end{aligned} \quad (51)$$

where $q_1, q_2, \dots, q_i$ are linearly independent and $h_{11}h_{22} \cdots h_{ii} \neq 0$, or there exists some $p < i$, such that

$$\delta(\bar{b})^0 = \sum_{l=1}^i \bar{a}_l \left(\sum_{j=1}^p h_{lj} q_j \right). \quad (52)$$

Proposition 8.3.1 Suppose we are given an one-sided differential partial boc $\mathfrak{B}$, in diagram (46) of 8.2, such that $\mathfrak{B}_{\mathcal{P}}$ satisfies P2 of 8.2, and $\delta(\bar{b})$ satisfies Formula (52). Then $R(\mathfrak{B})$ is not homogeneous.

Proof. Since $p < i$, there exists at least one vertex $\mathcal{I} \in \{\mathcal{I}_1, \dots, \mathcal{I}_r\}$ such that the number of the solid arrows: $\mathcal{P} \rightarrow \mathcal{I}$ is more than that of the dotted arrows: $\mathcal{I} \rightarrow \mathcal{P}$ given in (52). By a suitable reordering, we may assume that the solid arrows are $\bar{a}_1, \dots, \bar{a}_n$; and the dotted arrows are $q_1, \dots, q_m$ respectively with $n > m$. Let $\mathfrak{B}'$ be the induced partial boc with two vertices $\mathcal{P}, \mathcal{I}$ obtained from $\mathfrak{B}$ by deletion and

$$\delta(\bar{b})^0 = \sum_{l=1}^n \bar{a}_l \left(\sum_{j=1}^m h_{lj} q_j \right) = \sum_{j=1}^m \left(\sum_{l=1}^n h_{lj} \bar{a}_l \right) q_j$$

in $\mathfrak{B}'$. Suppose that $R(\mathfrak{B})$ is homogeneous, then $R(\mathfrak{B}')$ is homogeneous by Corollary 7.2.1. Consider a system of equations in n indeterminates:

$$\sum_{l=1}^n h_{lj} x_l = 0, \quad j = 1, \dots, m$$

Because the rank of the coefficient matrix is smaller than n , there is a non-zero solution of the equation system, say $x_1 = \mu^1, x_2 = \mu^2, \dots, x_n = \mu^n$, where $\mu^1 = 0, \dots, \mu^{e-1} = 0$, but $\mu^e \neq 0$ for some $1 \leq e \leq n$.

Define an object $S_{\nu^0} \in R(\mathfrak{B}')$ for any $\nu^0 \in k$, such that

$$(S_{\nu^0})_{\mathcal{P}} = k, \quad (S_{\nu^0})_{\mathcal{I}} = 0, \quad S_{\nu^0}(\bar{b}) = (\nu^0), \quad S_{\nu^0}(b) = \emptyset \quad \text{for any loop } b \neq \bar{b}.$$

If $R(\mathfrak{B}')$ is homogeneous, there exists a cofinite subset $D_0 \subseteq k$ such that S_{ν^0} is homogeneous for any $\nu^0 \in D_0$. We fix an $\nu^0 \in D_0$, denote S_{ν^0} by S for simplicity, and suppose that $(e) : S \xrightarrow{\iota} E \xrightarrow{\pi} S$ is an almost split conflation in $R(\mathfrak{B}')$. Then (e) is also an almost split conflation in $R(\mathfrak{B}'_{\mathcal{P}})$ by Theorem 7.2.1. Thus

$$E_{\mathcal{P}} = k^2, \quad E_{\mathcal{I}} = 0, \quad E(\bar{b}) = \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix}, \quad \text{and} \quad E(b) = \emptyset \quad \text{for } b \neq \bar{b}$$

by Proposition 6.6.1. Now we construct an object $L \in R(\mathfrak{B}')$, such that

$$L_{\mathcal{P}} = k, \quad L_{\mathcal{I}} = k, \quad L(\bar{a}_l) = (\mu^l), \quad L(a_j) = \emptyset \quad \text{for } j \in I, \quad L(\bar{b}) = (\nu^0), \quad L(b) = \emptyset \quad \text{for } b \neq \bar{b};$$

and we also construct a morphism $\varphi : L \rightarrow S$, such that

$$\varphi_{\mathcal{P}} = 1, \quad \varphi_{\mathcal{I}} = 0, \quad \varphi(u) = 0, \quad \varphi(v) = 0, \quad \varphi(q) = 0.$$

Then φ is not a split epimorphism. In fact, if $\psi : S \rightarrow L$ with $\psi\varphi = id$, then $\psi_{\mathcal{P}}\varphi_{\mathcal{P}} = 1$, $\psi_{\mathcal{I}} = 1$, and

$$-\mu^e = S(\bar{a}_e)\psi_{\mathcal{I}} - \psi_{\mathcal{P}}L(\bar{a}_e) = \psi(\delta(\bar{a}_e)) = 0.$$

We have a contradiction. Thus there exists some $\varphi' : L \rightarrow E$, such that $\varphi'\pi = \varphi$, whenever (e) is an almost split conflation in $R(\mathfrak{B}')$. Then $\varphi'_{\mathcal{P}} = (1, a)$ for some $a \in k$. On the other hand,

$$L(\bar{b})\varphi'_{\mathcal{P}} - \varphi'_{\mathcal{P}}E(\bar{b}) = \varphi'(\delta(\bar{b})) = 0, \quad \text{i.e.} \quad \nu^0(1, a) = (1, a) \begin{pmatrix} \nu^0 & 1 \\ 0 & \nu^0 \end{pmatrix}.$$

We have again a contradiction. Therefore the assumption fails, and $R(\mathfrak{B})$ is not homogeneous. $\square$

The possibilities of the differentials of $a_{n_0+1}, \dots, a_n$ with respect to $\bar{b}$ are given in the following two formulae.

$$\begin{cases} \delta(a_{n_0+1})^0 &= \sum_{l \in I} f_{n_0+1,l}(\bar{b})v_l + f_{n_0+1,n_0+1}(\bar{b})v_{n_0+1} \\ \dots & \dots \\ \delta(a_n)^0 &= \sum_{l \in I} f_{n,l}(\bar{b})v_l + f_{n,n_0+1}(\bar{b})v_{n_0+1} + \dots + f_{n,n}(\bar{b})v_n \end{cases} \quad (53)$$

where $\{v_l \mid \forall l \in I\}$ and $\{v_{n_0+1}, \dots, v_n\}$ are linearly independent, $f_{ll}(\bar{b}) \neq 0$. Or

$$\begin{cases} \delta(a_{n_0+1})^0 &= \sum_{l \in I} f_{n_0+1,l}(\bar{b})v_l + f_{n_0+1,n_0+1}(\bar{b})v_{n_0+1} \\ \dots & \dots \\ \delta(a_{n_1-1})^0 &= \sum_{l \in I} f_{n_1-1,l}(\bar{b})v_l + f_{n_1-1,n_0+1}(\bar{b})v_{n_0+1} + \dots + f_{n_1-1,n_1-1}(\bar{b})v_{n_1-1} \\ \delta(a_{n_1})^0 &= \sum_{l \in I} f_{n_1,l}(\bar{b})v_l + f_{n_1,n_0+1}(\bar{b})v_{n_0+1} + \dots + f_{n_1,n_1-1}(\bar{b})v_{n_1-1} \end{cases}$$

where $\{v_l \mid \forall l \in I\}$ and $\{v_{n_0+1}, \dots, v_{n_1-1}\}$ are linearly independent, $f_{ll}(\bar{b}) \neq 0$.

Proposition 8.3.2 Suppose we are given an one-sided differential partial boc $\mathfrak{B}$ in diagram (46) of 8.2, such that $\mathfrak{B}_{\mathcal{P}}$ satisfies P2 of 8.2; $\delta(\bar{b})$ satisfies Formula (51); and $\delta(a)$'s with respect to $\bar{b}$ satisfy the preceding second formula. Then $\mathfrak{B}$ is not homogeneous.

Proof. We set $\bar{a}_l = 0$ for $l = 1, \dots, i$, $a_l = \emptyset$ for $l \in I$, $\bar{b} = \lambda$, $b_l = \emptyset$ for $l \neq j$, and $a_{n_0+1} = \emptyset, \dots, a_{n_1-1} = \emptyset$. If $a_{n_1} : \mathcal{P} \rightarrow \mathcal{I}$, then we delete $T/\sim \setminus \{\mathcal{P}, \mathcal{I}\}$ and obtain an induced partial boc $\lambda \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \bullet \xrightarrow{a_{n_1}} \bullet$, where $\delta(a_{n_1}) = 0$. The non-homogeneous property of $\mathfrak{B}$ follows from Proposition 7.3.2 and Corollary 7.2.1. $\square$

8.4 The structure of loop $\bar{b}$

Let the partial boc $\mathfrak{B}$ be given as in 8.2 such that the induced local partial boc $\mathfrak{B}_{\mathcal{P}}$ is in case P2 of 8.2, $\delta(\bar{b})$ satisfies Formula (51) of 8.3, and $\delta(a)$'s with respect to $\bar{b}$ satisfy Formula (53) of 8.3. We will prove in this subsection, that the block determined by loop $\bar{b}$ will not contain any parameter after any series of reductions, whenever the parameters appear in a local boc.

We first assume that the index set $I = \emptyset$ in Formula (50) of 8.3 and $m = 1$, $\bar{b} = b_1$ given in 8.2. For any size vector $\underline{n}$ of $(T, \sim)$, we have a size vector

$$\underline{n}_1 \times \underline{n}_2 \text{ of } (T_1, \sim_1) \times (T_2, \sim_2)$$

according to Corollary 8.1.1. Then we start a sequence of reductions from triple $(N, X \times R, \underline{n}_1 \times \underline{n}_2)$ given by the matrix equation:

$$\begin{aligned} & X(D_1 \cdots D_i B A_{i+1} \cdots A_{i+j}) \\ = & (0 \cdots 0 \quad 0 \quad V_1 \cdots V_j) + (D_1 \cdots D_i B A_{i+1} \cdots A_{i+j})R \end{aligned} \quad (54)$$

where matrices $D_1, \dots, D_i$ and B correspond to the solid arrows $\bar{a}_1, \dots, \bar{a}_i$ and $\bar{b}$ respectively, A_l correspond to the solid arrows $a_l : \mathcal{P} \rightarrow \mathcal{I}$ for $l \geq i+1$;

$$R = \left(\begin{array}{ccc|ccc} Y_1 & & W & Q_1 & & \\ & \ddots & & \vdots & & W \\ & & Y_i & Q_i & & \\ \hline & & & X & V_1 & \cdots & V_j \\ & & & & Y_{i+1} & & W \\ & & & & & \ddots & \\ & & & & & & Y_{i+j} \end{array} \right)$$

where X, Y_l are $n_{\mathcal{P}} \times n_{\mathcal{P}}, n_{\mathcal{I}} \times n_{\mathcal{I}}$ invertible matrices, V, W, Q correspond to the dotted arrows v, w, q respectively. In particular, $Q_1, \dots, Q_i$ are blocks with entries being linearly independent of all the other entries according to Formula (51). Denote $(D_1 \cdots D_i B A_{i+1} \cdots A_{i+j})$ by N .

Lemma 8.4.1 All the reductions sequence (Δ) starting from the triple $(N, X \times R, \underline{n}_1 \times \underline{n}_2)$ defined above must reach a step r , such that either

- (1) the r -th step is an edge reduction before A_{i+1} , and $(X_{p_{r-1}, p_{r-1}})' = X_{(p_{r-1})_1, (p_{r-1})_1}$ (see the edge reduction given in 3.1); or
- (2) the r -th step is a loop reduction with a Weyr matrix W^0 inside B before A_{i+1} . Where all the reductions before r -th step are edge reductions and regularizations.

Moreover, the induced triple $(N^r, X^r \times R^r, \underline{n}_1^r \times \underline{n}_2^r)$ has the property that $Q_l^r = Q_l$, and the entries of the blocks Q_l^r and the nilpotent blocks of X^r are linearly independent and linearly independent of all the other entries of X^r, R^r, V^r .

Proof. Assume that the number of the columns of matrix $(D_1 \cdots D_i B)$ equals m . Our assertion is shown by induction on m . If m_B stands for the size of B and $m = m_B$, then the first reduction is a loop reduction for B , we obtain the case (2) for $r = 1$. Otherwise $m > m_B$, $\bar{a}_1$ is an edge and after an edge reduction, X is restricted to


$$X' = \begin{pmatrix} X_{11} & X_{12} \\ 0 & X_{22} \end{pmatrix}.$$

If $X_{22} = 0$, then $X' = X_{11}$, we obtain the case (1) for $r = 1$.

If $X_{22} \neq 0$, then X_{22} is independent of X_{11} and Y 's, if $X_{11} \neq 0$ and $X_{22} \neq 0$, then X_{12} is independent of Y, Q, W, V respectively. Write

$$m_1 = m - (\text{the number of columns of } D_1),$$

then $m_1 \leq m$. Note that the first reduction for $\bar{a}_1$ must be an edge reduction, but we write here also a regularization for a unified statement in the further reductions. If it is a regularization, then we still have $m_1 < m$.




In case of edge reduction and $X_{22} \neq 0$, we remove the block-row and the block-column where X_{11} sits from X^1 ; and those where Y_1 (according to the partition of $(T_2, \sim_2)$) sits from R^1 ; as well as $\bar{D}_1$, and the upper rows parallel to X_{11} from N^1 . Then we obtain 3 matrices shown in the shadowed part of diagram (55) for example, which still satisfy the hypothesis given in (54), if we use X_{22} instead of X , B^1 instead of B , $D_1^1, \dots, D_{i_1}^1$

partitioned after the edge reduction instead of $D_1, \dots, D_i$ respectively; the block X_{12} goes to part Q^1 , and X_{11} goes to part Y^1 . In case of regularization, we remove the block-row and the block-column where Y_1 sits from R^1 , as well as $\bar{D}_1$, and we also obtain 3 matrices satisfying the hypothesis given in (54).

Note that the above procedure is an illustration in order to help to understand the induction. Thus by a sequence of edge reductions and regularizations, or possibly a loop reduction at last, we finally reach case (1) or (2), since m is a finite number. Consequently, we obtain an induced triple of matrices $(N^r, X^r \times R^r, \underline{n}_1^r \times \underline{n}_2^r)$ before A_{i+1} , (see the diagram (56) below for example), such that Q_i^r and the nilpotent blocks of X^r satisfy the conditions of the lemma. $\square$

Lemma 8.4.2 With the notations above. Let $N^r = (D^r, B^r, A^r)$ be obtained from $N = (D, B, A)$ after all the reductions given in Lemma 8.4.1. Then Q^r and the nilpotent blocks of X^r force the rest blocks of B^r equal to $\emptyset$ in any sequence of further reductions under the order of Definition 2.3.1.



Proof. For the sake of simplicity, we denote X^r, N^r, R^r by X', N', R' respectively.

We first prove case (2) of Lemma 8.4.1 according to the partition of

$$(T_1^r \times T_2^r, \sim_1^r \times \sim_2^r).$$

Assume that $I_1, I_2, \dots, I_\beta$ are identity matrices appearing in the preceding edge reductions. The assertion will be shown by induction on β . $\beta = 0$ is trivial since B' is then obtained by a loop reduction and there is no block left in B' . Denote by $X_1, X_2, \dots, X_\beta$ the diagonal blocks of X' parallel to $I_1, I_2, \dots, I_\beta$ and X_0 that at the lower right corner of

X' parallel to the Weyr matrix W^0 . If $I_1, \dots, I_\alpha$ are sitting in part D' , and $I_{\alpha+1}, \dots, I_\beta$ are sitting in B' , then we denote by $D^1, \dots, D^\alpha$ the blocks of D' parallel to $I_1, I_2, \dots, I_\alpha$, which remain in N' after the reductions in Lemma 8.4.1; by $A^1, \dots, A^\beta, A^0$ the blocks of A' parallel to $I_1, \dots, I_\beta, W^0$ which remain in N' after the reductions in Lemma 8.4.1 respectively. See the middle matrix of diagram (56) for example, where $\beta = 4, \alpha = 2$. On the other hand, we denote by $Q^1, \dots, Q^\alpha$ the row-blocks in part Q parallel to $I_1, \dots, I_\alpha$ of $(N')^T$, the transpose of N' , and by $X^{\alpha+1}, \dots, X^\beta$ those of X' parallel to $I_{\alpha+1}, \dots, I_\beta$ of $(N')^T$ respectively in R' . See the shadowed blocks Q^1, Q^2, X^3, X^4 of diagram (56) for example. Denote Y', V', W', Q' of R' still by Y, V, W, Q respectively for simplicity. It is obvious that X^l are higher than X_l in X' .

The differentials in part A^0 involve only the entries of X_0, Y, V, W . They do not involve Q and the nilpotent part of X' at all. Therefore those blocks are still linearly independent of all the other blocks. The differential of the block of B' parallel to I_β has a term $I_\beta X^\beta$, which leads to $\emptyset_\beta$ by regularization (see X^4 and $\emptyset_4$ in (56) for example). Suppose that we have already $\emptyset_\beta, \emptyset_{\beta-1}, \dots, \emptyset_{l+1}$ parallel to $I_\beta, I_{\beta-1}, \dots, I_{l+1}$ in B' given by regularizations involving $X^\beta, X^{\beta-1}, \dots, X^{l+1}$ for some $l \geq \alpha$. Now we consider the block parallel to I_l in B' . Since the differentials in parts

$$A^0; \emptyset_\beta, A^\beta; \emptyset_{\beta-1}, A^{\beta-1}; \dots; \emptyset_{l+1}, A^{l+1}$$

involve only the entries of Y, V, W , and some blocks parallel to or lower than X_{l+1} in X' ; on the other hand, X^l is higher than X_{l+1} (see diagram (56) for example), we conclude that the entries of X^l are still linearly independent after the reduction for A^{l+1} . The differential of the block of B' parallel to I_l has a term $I_l X^l$, which leads to $\emptyset_l$ by regularization (see X^3 and $\emptyset_3$ in (55) for example). Then we obtain a sequence of $\emptyset$'s up to the $(\alpha + 1)$ -th block row in B' by induction. Moreover all the reductions before D^α involve only X', Y, V, W , and the entries of Q remain linearly independent.

Next the differential in parts $A^{\alpha+1}$ and D^α involve the entries of X', Y, V, W . The differential of the row-block of B' parallel to I_α has a term $I_\alpha Q^\alpha$, which leads to $\emptyset_\alpha$ by regularization (see Q^2 and $\emptyset_2$ in (56) for example). Suppose we already have $\emptyset_\alpha, \emptyset_{\alpha-1}, \dots, \emptyset_{l+1}$ parallel to $I_\alpha, I_{\alpha-1}, \dots, I_{l+1}$ given by regularizations involving $Q^\alpha, Q^{\alpha-1}, \dots, Q^{l+1}$ for some integer $l > 0$. Now we consider the block parallel to I^l in B' . Since the differentials for

$$A^{\alpha+1}, D^\alpha; \emptyset_\alpha, A^\alpha, D^{\alpha-1}; \dots; \emptyset_{l+1}, A^{l+1}, D^l$$

involve only the entries of X', Y, V, W , and some entries of Q lower than Q^l , we conclude that the entries of Q^l are still linearly independent after the reduction for D^l . The differential of the block of B' parallel to I_l has a term $I_l Q^l$, which leads to $\emptyset_l$ by regularization

(see Q^1 and $\emptyset_1$ in (56) for example). Thus we obtain a sequence of $\emptyset$'s up to the block parallel to I_1 in B' by induction.

Therefore the rest blocks of B' are all equal to $\emptyset$ in any sequence of reductions under the order given by Definition 2.3.1 according to the partition $(T_1^r \times T_2^r, \sim_1^r \times \sim_2^r)$.

The proof for the case (1) of Lemma 8.4.1 is as follows: if the $(r+1)$ -th reduction is for a block in A_{i+1} , we start the procedure from A^β , where $\beta \neq \alpha$; if the $(r+1)$ -th is for a block in B' , we start the procedure from $\emptyset_\beta$, where $\beta \neq \alpha$ or $\beta = \alpha$; if the $(r+1)$ -th reduction is for a block before B' , we start the procedure from D^β , where $\beta = \alpha$.

The lemma still holds, if we use the usual partition step by step in the reduction sequence. In fact we only need to make some refinement for the block-rows. $\square$

Proposition 8.4.1 Let $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$ and $\mathfrak{B}$ be given in diagram (46) of 8.2. Suppose that the partial boc $\mathfrak{B}$ is wild and the induced local partial boc $\mathfrak{B}_{\mathcal{P}}$ is in case P2 of 8.2; $\delta(\overline{b})$ satisfies Formula (51) of 8.3; and the partial differentials of a 's after $\overline{b}$ satisfy Formula (53) of 8.3. Given any size vector $\underline{n}_1 \times \underline{n}_2$ we have a matrix equation:

$$X(\cdots A \cdots B \cdots A \cdots B \cdots) = (V_1 \ V_2 \ \cdots \ V_n) + (\cdots A \cdots B \cdots A \cdots B \cdots)R_0$$

Then the reductions given by this equation are completely determined by the reductions given by Formula (54), and A_l goes to $\emptyset$ for any $l \in I$ in Formula (50) of 8.2, B_l goes to $\emptyset$ for any $l \neq j$ in Formula (47) and (48) of 8.2.

Moreover, assume that $\mathfrak{B}$ is a partial boc of $\mathfrak{A}$ appearing in a sequence $(*)$ with the end term in case of MW5. Then block $\overline{B}$ corresponding to $\overline{b}$ will not contain any parameter after any reductions.

Proof. (1) Consider the differentials of b_l for any $l \neq j$ given by Formula (47) and (48) of 8.2 with $m' = m$, $h_{ll}(\lambda)$ being non-zero constants ensure that the reduction for any block in B_l (corresponding to b_l), must be a regularization determined by U_l (corresponding to u_l), which does not influence the reductions of any other blocks in $(\cdots A \cdots B \cdots A \cdots B \cdots)$.

(2) Consider the differentials of a_l , $l \in I$, given by Formula (49) of 8.3, and the partial differentials of $a_{l'}$ given by Formula (53). Since

$$\{v_l \mid l \in I\} \quad \text{and} \quad \{v_{n_0+1}, \cdots, v_n\}$$

are linearly independent, any reduction (in particular, regularization) of any part of block $A_{l'}$ for $l' > n_0$ (see the beginning of 8.3) does not influence the regularization of any part of block A_l for $l \in I$. On the other hand, the regularization of any block of A_l does not influence the reductions of any other blocks in $(\cdots A \cdots B \cdots A \cdots B \cdots)$.

(1) and (2) tell us that the reductions given by the matrix equation in the proposition are completely determined by those given by Formula (54).

(3) Finally we prove the last assertion of the proposition. If we are in case (1) of Lemma 8.4.1, then there is no loop reduction (even no loop and edge reductions) in $\overline{B}$, the assertion holds obviously. Otherwise the only possible loop reduction inside $\overline{B}$ yields a Weyr matrix W^0 in case (2) of Lemma 8.4.1. We claim that the induced triple $(N^{r-1}, X^{r-1} \times R^{r-1}, \underline{n}_1^{r-1} \times \underline{n}_2^{r-1})$ before W_0 appearing is not local. In fact, if there exist some diagonal blocks X_i of X^{r-1} with $i > 0$, then X_0 is independent of all the X_i 's. Otherwise, $X_0 = X^{r-1}$, there must exist some Y_j as a diagonal block of R^{r-1} by wildness of $\mathfrak{B}$ and tameness of $\mathfrak{B}_{\mathcal{P}}$. Then X_0 is independent of all the Y_j 's. In both cases the induced partial boc at this step is not local. We conclude that $\overline{B}$ will not contain any parameter after any reductions, since MW5 requires that any parameter appears for the first time in a local boc by Corollary 5.6.1. The proof is finished. $\square$

8.5 Some lemmas in algebra

This subsection is devoted to giving several lemmas in algebra which are needed in Section 9.

Let

$$\{x_1, \dots, x_m\} \quad \text{and} \quad \{y_1, \dots, y_m\}$$

be two sets of indeterminates. We say that

$$a_1x_1 + a_2x_2 + \dots + a_mx_m$$

and

$$a_1y_1 + a_2y_2 + \dots + a_my_m$$


are the same linear forms, where $a_1, \dots, a_m \in k$ are constants.

Lemma 8.5.1 Let $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ be a matrix problem, and $(N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2)$ be a triple. Suppose that $\mathcal{N}$ is a matrix with constant entries of size $\underline{n}_1 \times \underline{n}_2$. If we fix a row-index α and an index $l \in \mathcal{L} \in T_2/\sim_2$, then any linear forms at the (α, β) -entry of $V + \mathcal{N}R_2$ in the indeterminates

$$(\overline{\xi_{\mathcal{IL}}^{v_{123}}})^*, (\overline{\xi_{\mathcal{IL}}^{v_{ij}}})^*, (i, j) = \{(12), (23), (31)\} \quad \text{and} \quad (\overline{\xi_{\mathcal{IL}}^{v_i}})^*, \quad i = 1, 2, 3, \quad \forall \mathcal{I} \in T/\sim$$

given in 8.1 at β -th column of V and R_2 are the same for all $\beta \in l$, (see the figure below).

$\square$



For example, in the matrix

$$\begin{pmatrix} v_{11} & v_{12} & v_{13} \\ v_{21} & v_{22} & v_{23} \end{pmatrix} + \begin{pmatrix} 0 & 1 & \lambda^0 \\ 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{pmatrix}$$

the linear forms at the first row are

$$v_{11} + x_{21} + \lambda^0 x_{31}, \quad v_{12} + x_{22} + \lambda^0 x_{32}, \quad v_{13} + x_{23} + \lambda^0 x_{33}.$$

Lemma 8.5.2 Let $Q : \mathcal{P} \xrightarrow{c} \mathcal{Q}$ be a quiver, and the diagram

$$(e) \quad \begin{array}{ccccccc} 0 & \longrightarrow & k & \xrightarrow{\iota_{\mathcal{P}}} & k^2 & \xrightarrow{\pi_{\mathcal{P}}} & k \longrightarrow 0 \\ & & \alpha \downarrow & & \downarrow \beta & & \downarrow \alpha \\ 0 & \longrightarrow & k^m & \xrightarrow{\iota_{\mathcal{Q}}} & k^{2m} & \xrightarrow{\pi_{\mathcal{Q}}} & k^m \longrightarrow 0 \end{array}$$

be an exact sequence over the path algebra kQ .


(1) If $\alpha = (0 \ \cdots \ 0 \ 1)$, then $\beta = \begin{pmatrix} 0 & \cdots & 0 & 1 & 0 \\ 0 & \cdots & 0 & 0 & 1 \end{pmatrix}_{2 \times 2m}$, and $\iota_{\mathcal{P}} = (0 \ 1)$, $\pi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$

under some suitable basis.

(2) If $\alpha = (0 \ \cdots \ 0 \ 0)$, then $\beta = 0$ or $\beta = \begin{pmatrix} 0 & \cdots & 0 & 0 & 1 \\ 0 & \cdots & 0 & 0 & 0 \end{pmatrix}_{2 \times 2m}$, and $\iota_{\mathcal{P}} = (0 \ 1)$,

$\pi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ under some suitable basis. $\square$

Lemma 8.5.3 Let $Q = \lambda \begin{array}{c} \curvearrowright \\ \bullet \end{array} \xrightarrow{c} \begin{array}{c} \bullet \\ \curvearrowright \end{array} \mathcal{Q}$ be a quiver, and kQ be the corresponding path algebra. Let $\mathcal{C}$ be a full subcategory of $\text{mod } kQ$ consisting of modules M , such that the eigenvalue of $M(\lambda)$ is λ^0 .



The above diagram in $\mathcal{C}$ consists of the modules S, E, L , where

$$S_{\mathcal{P}} = k, S_{\mathcal{Q}} = k^m, S(\lambda) = (\lambda^0), S(c) = 0;$$

$$E_{\mathcal{P}} = k^2, E_{\mathcal{Q}} = k^{2m}, E(\lambda) = \lambda^0 I_2 \text{ or } J_2(\lambda^0), \text{ and } E(c) = \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 0 & 0 \end{pmatrix};$$

$$L_{\mathcal{P}} = k^2, L_{\mathcal{Q}} = k^{2m}, L(\lambda) = J_2(\lambda^0), L(c) = 0;$$

and morphisms

$$\pi = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} I_m \\ 0 \end{pmatrix} \right) : E \rightarrow S, \varphi = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} I_m \\ 0 \end{pmatrix} \right) : L \rightarrow S$$

in kQ -mod. Then

- (1) the morphism φ is not a split epimorphism.
- (2) there does not exist any morphism $\varphi' : L \rightarrow M$, such that $\varphi'\pi = \varphi$.

Proof. (1) is obvious by considering $\begin{pmatrix} 1 \\ 0 \end{pmatrix} : J_2(\lambda) \rightarrow J_1(\lambda)$.


(2) Suppose we have the contrary. Then $\varphi'_{\mathcal{P}}\pi_{\mathcal{P}} = \varphi_{\mathcal{P}}$ gives $\varphi'_{\mathcal{P}} = \begin{pmatrix} 1 & b \\ 0 & d \end{pmatrix}$. On the other hand

$$0 = L(c)\varphi'_{\mathcal{Q}} = \varphi'_{\mathcal{P}}E(c) = \begin{pmatrix} 1 & b \\ 0 & d \end{pmatrix} \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 0 & 0 \end{pmatrix},$$

which means a contradiction. $\square$

8.6 Some lemmas in category theory

Lemma 8.6.1 Let $Q_0 =$



be a quiver consisting of vertices $\mathcal{P}, \mathcal{Q}_1, \dots, \mathcal{Q}_r$; a loop λ at $\mathcal{P}$, and n_j arrows $a_{j1}, \dots, a_{jn_j}$ from $\mathcal{P}$ to $\mathcal{Q}_j$. Let $\Gamma_0 = kQ_0$ be the path algebra. We define a full subcategory $\mathcal{C}_0$ of Γ_0 -mod consisting of modules M , such that $M_{\mathcal{P}} = k^m$, $M_{\mathcal{Q}_j} = k^{mn_j}$; $M(\lambda) = W$, which is a Weyr matrix of eigenvalue λ^0 of size m under a fixed basis B of $M_{\mathcal{P}}$; and

$$M(a_{jl}) = (0 \ \cdots \ 0 \ \underbrace{I_m \ 0 \ \cdots \ 0}_l)_{1 \times n_j}$$

under a basis $B_{j1}, \dots, B_{jn_j}$ of M_{Q_j} . Then for any $M, L \in \mathcal{C}_0$, a morphism $\varphi = (\varphi_{\mathcal{P}}; \varphi_{Q_1}, \dots, \varphi_{Q_r}) : M \rightarrow L$ is given by

$$\varphi_{\mathcal{P}} = Z, \quad \varphi_{Q_j} = \begin{pmatrix} Z & & & \\ & Z & & \\ & & \ddots & \\ & & & Z \end{pmatrix}_{n_j \times n_j}$$

under the basis B and $B_{j1}, \dots, B_{jn_j}$ for $j = 1, \dots, r$, where $M(\lambda)Z = ZL(\lambda)$. $\square$

Lemma 8.6.2 Let Q be a quiver obtained from Q_0 by adding one arrow $c : \mathcal{P} \rightarrow Q$, where either

$$(i) \ Q \notin \{Q_1, \dots, Q_r\}, \text{ or } (ii) \ Q = Q_{j_0} \text{ for some } 1 \leq j_0 \leq r,$$

and $\Gamma = kQ$. We define a full subcategory $\mathcal{C}$ of $\Gamma\text{-mod}$ as follows.

Let $M_{\mathcal{P}} = k^m$ have a k -basis B , and $M_{Q_j} = k^{m \cdot n_j}$ have a k -basis $(B_{j1}, \dots, B_{jn_j})$ given by Lemma 8.6.1, where $1 \leq j \leq r$ in case (i), or $j \neq j_0$ in case (ii). And let $M_Q = k^n$ in case (i), $M_{Q_{j_0}} = k^n \oplus k^{m \cdot n_{j_0}}$ in case (ii), where $k^{m \cdot n_{j_0}}$ has a k -basis $(B_{j_01}, \dots, B_{j_0n_{j_0}})$. Moreover $M(\lambda) = W$ of eigenvalue λ^0 , and $M(a_{jl})$ are given by Lemma 8.6.1, except

$$M(a_{j_0l}) = (0 \mid 0 \cdots 0 \underbrace{I_m \ 0 \cdots 0}_l)_{1 \times (n_{j_0}+1)}$$

in case (ii), where 0 before the vertical line is an $m \times n$ matrix. And

$$M(c) = P \text{ in case (i), } M(c) = (P \mid 0 \cdots 0)_{1 \times (n_{j_0}+1)} \text{ in case (ii).}$$

Then

(1) $\forall M, L \in \mathcal{C}$, a morphism

$$\varphi = (\varphi_{\mathcal{P}}; \varphi_1, \dots, \varphi_r, \varphi_Q) \text{ or } (\varphi_{\mathcal{P}}, \varphi_1, \dots, \varphi_r)$$

from M to L under the fixed basis is given by

$$\varphi_{\mathcal{P}} = Z \text{ satisfying } M(\lambda)Z = ZL(\lambda); \quad \varphi_{Q_l} = \begin{pmatrix} Z & & & \\ & Z & & \\ & & \ddots & \\ & & & Z \end{pmatrix}$$

where $1 \leq l \leq r$ in case (i), $l \neq j_0$ in case (ii); and

$$\varphi_Q = U \text{ in case (i); } \varphi_{j_0} = \left(\begin{array}{c|ccc} U & * & \cdots & * \\ \hline & Z & & \\ & & \ddots & \\ & & & Z \end{array} \right) \text{ in case (ii),}$$

where $L(c) = P$ and $PU = ZP'$ in case (i), $L(c) = (P' \mid 0 \cdots 0)$ with $PU = ZP'$ and $P* = 0$ in case (ii).

(2) $\mathcal{C}$ is an exact category having the exact structure inherited from $\Gamma\text{-mod}$.

Proof. (1) is obvious.

(2) We only prove (E2) of 6.1. If $(e) \quad 0 \longrightarrow M \xrightarrow{\iota} E \xrightarrow{\pi} L \longrightarrow 0$ is a conflation of $\mathcal{C}$, and $\varphi : L' \rightarrow L$ is a morphism of $\mathcal{C}$, then a pullback

$$\begin{array}{ccccccccc} 0 & \longrightarrow & M & \xrightarrow{\iota'} & E' & \xrightarrow{\pi'} & L' & \longrightarrow & 0 \\ & & \downarrow id & & \downarrow \varphi' & & \downarrow \varphi & & \\ 0 & \longrightarrow & M & \xrightarrow{\iota} & E & \xrightarrow{\pi} & L & \longrightarrow & 0 \end{array}$$

of $\Gamma\text{-mod}$ gives the desired deflation π' of $\mathcal{C}$. In fact the eigenvalue of $E'(\lambda)$ must be λ^0 , since E' is a submodule of $E \oplus L'$. $\square$

Lemma 8.6.3 Let $Q = \lambda \curvearrowright \bullet \xrightarrow{c} \bullet \xrightarrow{Q}$ be a quiver, $\Gamma = kQ$ be the path algebra, and

$$(e) \quad 0 \rightarrow S \rightarrow E \rightarrow S \rightarrow 0$$

be the exact sequence given by Lemma 7.6.2. We define a full subcategory $\mathcal{D}$ of $\Gamma\text{-mod}$ consisting of modules M , such that $M_{\mathcal{P}} = k^{m_{\mathcal{P}}}$; $M_Q = k^{m_Q}$; and the eigenvalue of $M(\lambda)$ is a constant λ^0 . Then

(1) $\mathcal{D}$ is an exact category having the exact structure inherited from $\Gamma\text{-mod}$.

(2) (e) is not an almost split sequence in $\mathcal{D}$.

Proof. (1) is a special case of Lemma 8.6.2.

(2) The almost split sequences in $\mathcal{D}$ has also the property of item (2) of Lemma 7.6.1. In fact if

$$0 \longrightarrow M \xrightarrow{\iota} N \xrightarrow{\pi} L \longrightarrow 0$$

is an almost split conflation of $\mathcal{D}$, K is a non-zero submodule of M , then K and M/K are contained in $\mathcal{D}$. The following diagram of $\Gamma\text{-mod}$

$$\begin{array}{ccccccccc} 0 & \longrightarrow & M & \xrightarrow{\iota} & N & \xrightarrow{\pi} & L & \longrightarrow & 0 \\ & & \downarrow \psi & & \downarrow \psi & & \downarrow id & & \\ 0 & \longrightarrow & M/K & \longrightarrow & N/\iota(K) & \longrightarrow & L & \longrightarrow & 0 \end{array}$$

is still in $\mathcal{D}$. Thus the second sequence splits in $\mathcal{D}$. Therefore Lemma 7.6.2 tells us that (e) is not an almost split sequence of $\mathcal{D}$. The proof is completed. $\square$

Lemma 8.6.4 Let $\mathcal{C}$ and $\mathcal{D}$ be two exact categories. Suppose there exists a functor $F : \mathcal{C} \rightarrow \mathcal{D}$, which is a representation equivalence (i.e. dense, full, and reflects isomorphisms). If $(e) \quad M \xrightarrow{\iota} N \xrightarrow{\pi} L$ is an almost split conflation of $\mathcal{C}$ such that $F(M)$ (resp. $F(L)$) is non-injective (resp. non-projective), then $F(e)$ is an almost split conflation in $\mathcal{D}$.

Proof. Since F reflects isomorphisms, $F(\pi)$ is not a retraction. Suppose that $\varphi' : K' \rightarrow F(L)$ is a morphism in $\mathcal{D}$, which is not a split epimorphism. Then there exists some morphism $\varphi : K \rightarrow L$ in $\mathcal{C}$ such that $F(K) = K'$, $F(\varphi) = \varphi'$, since F is dense and full. And φ is not a split epimorphism in $\mathcal{C}$, otherwise φ' would be so in $\mathcal{D}$. Then there exists some $\psi : K \rightarrow L$ with $\psi\pi = \varphi$, so $F(\psi)F(\pi) = \varphi'$, thus $F(\pi)$ is a right almost split morphism. Similarly, we can prove that $F(\iota)$ is a left almost split morphism. And $F(e)$ is an almost split conflation of $\mathcal{D}$ by the definition given in 6.1. $\square$

9 Added columns

This section is still devoted to presenting some technical preparations in order to prove our main theorem in case MW5 of Theorem 5.6.1.

9.1 The structure of added columns

Let $(\mathcal{K}, \mathcal{M}, H)$ be a bimodule problem with a reduction sequence $(*)$ of freely parameterized triples starting from $(N_0, R, \underline{n})$, such that the end term of the corresponding sequence $(***)$ is in the case of MW5 of Theorem 5.6.1. Then λ appears for the first time in $\mathfrak{A}^s$ by corollary 5.6.1, and we have Formula (42) of 5.4. If $h(\lambda) \neq 0$ in the Formula (43), then a sequence of regularizations and a loop reduction for a_i gives a parameter $\lambda_\gamma = \nu$.

Definition 9.1.1 Denote the matrix index of λ in $H_{\underline{n}}^s$ by $(\alpha(0), \beta(0))$; the index of ν in $H_{\underline{n}^{s+i}}^{s+i}$ by (α^0, β^0) . Suppose that the parameter ν is contained in a block G partitioned by $(T, \sim)$, the index of the block-column of G is l^0 , and $l^0 \in \mathcal{L}^0 \in T/\sim$. $\square$

Let $(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$ be a partial bimodule problem, $\mathfrak{B}$ be the corresponding partial boc. Suppose that $(\mathcal{K}, \mathcal{M}/\mathcal{M}', d)$ determines a matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$, for example given by Proposition 8.1.1 or 8.1.2.

Remark. Let us take

$$\mathcal{N}^r = \sum_{l=0}^{r-1} \overline{N_{p_l q_l}^l} \otimes \chi^l \text{ for } 1 \leq r \leq s,$$

where $(\chi^l)^*$ is the first arrow of $(\mathcal{K}_1^l \times \mathcal{K}_2^l, \overline{\mathcal{M}}^l, \mathcal{V}^l)$, $\overline{N_{p_0 q_0}^0} = \overline{N_{pq}}$, $\chi^0 = \chi$. Now we fix a k -basis B' for the matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$ given in 8.1, and we regard the coefficient functions $(B')^*$ as the indeterminates. In particular, $\overline{(\xi_{\mathcal{I}\mathcal{J}}^v)^*}$, $\overline{(\xi_{\mathcal{I}\mathcal{J}}^{v_{ij}})^*}$, $(ij) = (12), (23)$ or (31) , $\overline{(\xi_{\mathcal{I}\mathcal{J}}^{v_i})^*}$, $i = 1, 2, 3$, and $\overline{(E_{\mathcal{I}})^*}$ are regarded as $n_{\mathcal{I}} \times n_{\mathcal{J}}$ and $n_{\mathcal{I}} \times n_{\mathcal{I}}$ matrices with entries being indeterminates (see 8.1). Fix an index r , R_1^r, R_2^r, V^r can be obtained from R_1, R_2, V restricted by the matrix equation:

$$R_1 \mathcal{N}^r \equiv_{\leq(p_{r-1}, q_{r-1})} V + \mathcal{N}^r R_2.$$

the equation system Π' is given by the above matrix equation and the equation $x_{p_r q_r}^3 = 0$ is given by (the (p_r, q_r) -block of $\mathcal{N}^r R_2 - R_1 \mathcal{N}^r = 0$).

Definition 9.1.2 Consider the matrix equation $R_1 \mathcal{N}^r = V + \mathcal{N}^r R_2$. Denote by $eq_{(\alpha, \beta)}$ the (α, β) -entry, and by $E_{q_{(\alpha, \beta)}}$ the set of entries indexed before (α, β) under the matrix order of Definition 2.3.1. $\square$

Lemma 9.1.1 Suppose that $\underline{n}_1^r \times \underline{n}_2^r$ has all the components to be 1, and $N_{p_r q_r}$ is the first non-zero block of N^r with the matrix index (α_r, β_r) . Then

(1) $eq_{(\alpha_r, \beta_r)}$ is the equation $x_{p_r q_r}^3 = 0$ in $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \overline{\mathcal{M}}^r, H^r)$; $Eq_{(\alpha_r, \beta_r)}$ consists of the equation system Π' of Definition 8.1.1 and the condition on the diagonal elements determined by $(T_1^r \times T_2^r, \sim_1^r \times \sim_2^r)$.

(2) $eq_{(\alpha_r, \beta_r)}$ is a linear combination of some equations in $E_{q(\alpha_r, \beta_r)}$, if and only if the equation $x_{p_r q_r}^3 = 0$ is a linear combination of some equations in Π' of 8.1.1 in $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \overline{\mathcal{M}}^r, \mathcal{V}^r)$.

Let (Δ) be a reduction sequence of 8.1 starting from $(N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2)$, with G being also a block partitioned by $(T_1 \times T_2, \sim_1 \times \sim_2)$, such that

S01 the end term $\mathfrak{B}^s$ of the corresponding sequence of partial bocses is in the case of MW5 of Theorem 5.6.1;

S02 $l^0 \in L^0 \in T_2 / \sim_2$, such that $\mathcal{L}^0 \cap T_1 = \emptyset$.

Note that since $\mathfrak{A}^s$, as well as $\mathfrak{B}^s$, are both local, we have that $\forall \mathcal{I} \in T / \sim, \mathcal{I} \upharpoonright_{T_1} \neq \emptyset$ or $\mathcal{I} \upharpoonright_{T_2} \neq \emptyset$.

Under the assumptions S01 and S02, we define a new size vector $\tilde{n}$ of $(T, \sim)$, with

$$n_{\mathcal{I}} = n_{\mathcal{I}} \quad \forall \mathcal{I} \in T / \sim \setminus \{\mathcal{L}^0\} \text{ and } \tilde{n}_{\mathcal{L}^0} = n_{\mathcal{L}^0} + 1.$$

Let $\tilde{\mathcal{N}}^s$ be obtained from $\mathcal{N}^s$ by adding a column to the left of each l -th block column consisting of 0's for all $l \in \mathcal{L}^0$. Suppose $\mathcal{L}^0 = \{l_1 < l_2 < \cdots < l_{\mathcal{L}^0}\}$, we denote by

$$o_1 < o_2 < \cdots < o_{l_{\mathcal{L}^0}}$$

the indices of the added columns. Let

$$\tilde{T} = T \bigcup \{o_1, o_2, \cdots, o_{l_{\mathcal{L}^0}}\}.$$

In fact, $\tilde{\mathcal{N}}^s$ is just a direct sum of $\mathcal{N}^s$ and a zero representation of size vector $\underline{m}$, where $m_{\mathcal{L}^0} = 1, m_{\mathcal{I}} = 0, \forall \mathcal{I} \in T / \sim \setminus \{\mathcal{L}^0\}$ according to the partition of $(T, \sim)$. We will prove in the next subsection, that there exists a reduction sequence $(\tilde{\Delta})$ based on (Δ) :

$$(\tilde{\Delta}) \quad (\tilde{N}, \tilde{R}_1 \times \tilde{R}_2, \tilde{\underline{n}}_1 \times \tilde{\underline{n}}_2), \cdots, (\tilde{N}^r, \tilde{R}_1^r \times \tilde{R}_2^r, \tilde{\underline{n}}_1^r \times \tilde{\underline{n}}_2^r), \cdots, (\tilde{N}^s, \tilde{R}_1^s \times \tilde{R}_2^s, \tilde{\underline{n}}_1^s \times \tilde{\underline{n}}_2^s)$$

Next we may compare the entries in the following two matrix equations:

$$R_1 N^r = V + N^r R \text{ and } \tilde{R}_1 \tilde{\mathcal{N}}^r = \tilde{V} + \tilde{\mathcal{N}}^r \tilde{R}_2.$$

Given any pair of indices (α, β) of N^r , denote by $\tilde{e}q_{(\alpha, \beta)}$ the equation given by the (α, β) -entry and by $\tilde{e}q_{(\alpha, o_l)}$ the equation given by the (α, o_l) -entry in the second matrix equation.

Lemma 9.1.2 (1) $eq_{(\alpha, \beta)}$ and $\tilde{e}q_{(\alpha, \beta)}$ are the same; their right hand sides given by $V + \mathcal{N}^r R_2$ and $\tilde{V} + \tilde{\mathcal{N}}^r \tilde{R}_2$ contain only the indeterminates at the β -th column of R_2, V and $\tilde{R}_2, \tilde{V}$ respectively.

(2) The left side of the equation $\tilde{e}q_{(\alpha, o_l)}$ given by $\tilde{R}_1 \tilde{\mathcal{N}}^r$ equals zero; and the right side of $\tilde{e}q_{(\alpha, o_l)}$ given by $\tilde{V} + \tilde{\mathcal{N}}^r \tilde{R}_2$ contains only the indeterminates at the o_l -th column of $\tilde{R}_2$ and $\tilde{V}$. $\square$

9.2 The reduction sequence $(\tilde{\Delta})$

We may write the matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ instead of $(\mathcal{K}_1 \times \mathcal{K}_2, \overline{\mathcal{M}}, \mathcal{V})$ for simplicity.

Proposition 9.2.1 There exists a reduction sequence $(\tilde{\Delta})$ given in 9.1, such that the added columns of $\tilde{\mathcal{N}}^s$ consist of zero's and $\emptyset$'s.




Figure 1

(i, l) -block:




Figure 2




Figure 3




Figure 4

Proof. When $r = 0$, we have $\tilde{T} = T \cup \{o_1, o_2, \dots, o_{l_{\mathcal{L}^0}}\}$ and the numbers of equivalent classes of $T/\sim$ and of $\tilde{T}/\sim$ are the same, and the equation system II' at $(\mathcal{P}_0, \mathcal{Q}_0)$ for $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ and at $(\tilde{\mathcal{P}}_0, \tilde{\mathcal{Q}}_0)$ for $(\tilde{\mathcal{K}}_1 \times \tilde{\mathcal{K}}_2, \tilde{\mathcal{M}}, \tilde{\mathcal{V}})$ are also the same. Suppose the r -th reduction has been done and we obtain a triple $(\tilde{N}^r, \tilde{R}_1^r \times \tilde{R}_2^r, \tilde{n}_1^r \times \tilde{n}_2^r)$, such that $\tilde{N}^r$ is obtained from N^r by adding a column at the left of the l -th block column consisting of 0's and $\emptyset$'s for all $l \in \mathcal{L}^0$.

(1) If the number of the equivalent classes of $\tilde{T}^r/\sim^r$ equals that of $T^r/\sim^r$, the equation system II at $(\mathcal{P}_r, \mathcal{Q}_r)$ in Definition 8.1.1 and the equation $x_{p_r q_r}^3 = 0$ for $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \mathcal{M}^r, \mathcal{V}^r)$ and those at $(\tilde{\mathcal{P}}_r, \tilde{\mathcal{Q}}_r)$ for $(\tilde{\mathcal{K}}_1^r \times \tilde{\mathcal{K}}_2^r, \tilde{\mathcal{M}}^r, \tilde{\mathcal{V}}^r)$ are the same. Thus the reduction type of both of them are the same. Suppose $N_{p_r q_r}^r$, the first block of N^r , belongs to the (i, j) -block according to the partition of $(T, \sim)$. If $j \notin \mathcal{L}^0$, or $j \in \mathcal{L}^0$ but $N_{p_r q_r}^r$ does not intersect the first column of the j -th block column of N^r , then $\tilde{n}_{p_r}^r = n_{p_r}^r$, $\tilde{n}_{q_r}^r = n_{q_r}^r$, and we set $\overline{\tilde{N}_{p_r q_r}^r} = \overline{N_{p_r q_r}^r}$ (see figure 2). If $j = l \in \mathcal{L}^0$ and $N_{p_r q_r}^r$ intersects the first column of the l -th block column of N^r , then $\tilde{n}_{p_r}^r = n_{p_r}^r$, but $\tilde{n}_{q_r}^r = n_{q_r}^r + 1$ (see figure 3).

Then $p_r \approx q_r$ in $(T^r, \sim^r)$, otherwise we would have $n_{p_r}^r = n_{q_r}^r$ and $\tilde{n}_{p_r}^r = \tilde{n}_{q_r}^r$, which is a contradiction. Now we give the $(r+1)$ -th reduction for $(\tilde{N}^r, \tilde{R}_1^r \times \tilde{R}_2^r, \tilde{\underline{n}}_1^r \times \tilde{\underline{n}}_2^r)$ based on that for $(N^r, R_1^r \times R_2^r, \underline{n}_1^r \times \underline{n}_2^r)$.

Regularization. we set $\overline{\tilde{N}_{p_r q_r}^r} = \emptyset$;

Edge reduction. we set $\overline{\tilde{N}_{p_r q_r}^r} = \begin{pmatrix} 0 & 0 & I_d \\ 0 & 0 & 0 \end{pmatrix}$;

Loop reduction. it can not occur, because of $p_r \approx q_r$.

Therefore $\tilde{\mathcal{N}}^{r+1}$ is obtained from $\mathcal{N}^{r+1}$ by adding a column at the left of each l -th block column consisting of 0's or $\emptyset$'s for all $l \in \mathcal{L}^0$ in this case.

(2) If $d = n_{q_{r_0}}^{r_0}$ in the preceding edge reduction for some step r_0 , and

$$\overline{\tilde{N}_{p_r q_r}^r} = \begin{pmatrix} 0_{d \times 1} & \vdots & I_d \\ 0 & & 0 \end{pmatrix},$$

then a new equivalent class $\mathcal{Q} = \{o_1, o_2, \dots, o_{l_{\mathcal{Q}}}\}$ takes place at this stage. Thus

$$\tilde{T}^{r_0+1}/\sim^{r_0+1} = (T^{r_0+1}/\sim^{r_0+1}) \cup \{\mathcal{Q}\}.$$

We stress that the situation must occur, since $\mathfrak{B}^s$ is local.

(3) Suppose $r > r_0$. If $j \notin \mathcal{L}^0$, or $j \in \mathcal{L}^0$ but $N_{p_r q_r}^r$ does not intersect the first column of the j -th block column of N^r , then $\tilde{n}_{p_r}^r = n_{p_r}^r$, $\tilde{n}_{q_r}^r = n_{q_r}^r$. We set $\overline{\tilde{N}_{p_r q_r}^r} = \overline{N_{p_r q_r}^r}$, since the equation systems II' are the same at $(\mathcal{P}_r, \mathcal{Q}_r)$ for both $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \mathcal{M}^r, \mathcal{V}^r)$ and $(\tilde{\mathcal{K}}_1^r \times \tilde{\mathcal{K}}_2^r, \tilde{\mathcal{M}}^r, \tilde{\mathcal{V}}^r)$ by Lemma 9.1.2. Otherwise $\tilde{n}_{p_r}^r = n_{p_r}^r$, $\tilde{n}_{q_r}^r = n_{q_r}^r$, $\tilde{n}_{o_l} = \tilde{n}_{\mathcal{Q}} = 1$ (see figure 4). The $(r+1)$ -th reduction for $\tilde{N}_{p_r q_r}^r$ is divided into two reductions based on the reduction for $N_{p_r q_r}^r$. Since equation system II' at $(\mathcal{P}_r, \mathcal{Q}_r)$ are the same in both $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \mathcal{M}^r, \mathcal{V}^r)$ and $(\tilde{\mathcal{K}}_1^r \times \tilde{\mathcal{K}}_2^r, \tilde{\mathcal{M}}^r, \tilde{\mathcal{V}}^r)$, we set

$$\overline{\tilde{N}_{p_r q_r}^r} = (0 \mid \overline{N_{p_r q_r}^r}) \text{ or } \overline{\tilde{N}_{p_r q_r}^r} = (\emptyset \mid \overline{N_{p_r q_r}^r})$$

according to the equation $x_{p_r o_l}^3 = 0$ is, or is not a linear combination of the equations in II' of 8.1.1 at $(\mathcal{P}_r, \mathcal{Q})$ for $(\tilde{\mathcal{K}}_1^r \times \tilde{\mathcal{K}}_2^r, \tilde{\mathcal{M}}^r, \tilde{\mathcal{V}}^r)$.

Thus $\tilde{\mathcal{N}}^{r+1}$ is obtained from $\mathcal{N}^{r+1}$ by adding a column at the left of the l -th block column consisting of 0's and $\emptyset$'s for all $l \in \mathcal{L}^0$ in this case.

Finally, $(\tilde{\Delta})$ is a reduction sequence by induction on r . The proof is completed. $\square$

Corollary 9.2.1 $(\tilde{T}^s, \sim^s)$ possesses two equivalent classes: one is the unique class $\mathcal{P}$ of $(T^s, \sim^s)$, another one is $\mathcal{Q} = \{o_1, o_2, \dots, o_{l_{\mathcal{Q}}}\}$.

Suppose that for any $i, j \in T$, R_{ij}^s stands for the (i, j) -block of R_2^s partitioned by $T_2/\sim_2$, and $\tilde{R}_{ij}^s$ for the (i, j) -block of $\tilde{R}_2^s$ partitioned of by $\tilde{T}_2/\sim_2$.

Corollary 9.2.2 (1) $\tilde{R}_1^r = R_1^r$ for $r = 0, 1, \dots, s$.

(2) $\forall i, j \in T_2 \setminus \mathcal{L}^0$, $\tilde{R}_{ij}^s = R_{ij}^s$; and $\forall l_1, l_2 \in \mathcal{L}^0$, $\tilde{R}_{l_1 j}^s$ has a subblock $R_{l_1 j}^s$ at the lower $n_{\mathcal{L}^0} \times n_{\mathcal{J}}$ part, $\tilde{R}_{i l_2}^s$ has a subblock $R_{i l_2}^s$ at the right $n_{\mathcal{I}} \times n_{\mathcal{L}^0}$ part, $\tilde{R}_{l_1 l_2}^s$ has a subblock $R_{l_1 l_2}^s$ at the lower-right $n_{\mathcal{L}^0} \times n_{\mathcal{L}^0}$ part (see figure 5).

(3) The diagonal block $\tilde{R}_{ll}^s$ shown in figure 6 has free entries at the top-row and zero's at the left-column below the top-row.




figure 5: from R_2^s to $\tilde{R}_2^s$

figure 6

Proof. We first prove the assertions (1) and (2). By lemma 9.1.2, $\emptyset$'s, appearing in the added column, only influence the added column of $\tilde{R}_2$, do not influence $R_1 \times R_2$ at all.

(3) The elements $*$ in $\tilde{R}_{ll}^s$ of figure 6 are not involved in any equations, therefore they are independent of all the others. The elements 0 at the left column of $\tilde{R}_{ll}^s$ are given by case (2) of the proof of Proposition 9.2.1, since the $*$ at the upper-left corner of figure 6 must determine a single equivalent class $\mathcal{Q}$ at this stage. $\square$

Example 1. Consider Example of 7.6. Let $\mathcal{K}_1 = \tilde{\Lambda}, \mathcal{K}_2 = \tilde{\Lambda}, \mathcal{M} = \text{rad} \tilde{\Lambda}$ and $\mathcal{V} = 0$. We obtain a matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ given by Proposition 8.1.1 with an index set $(T, \sim)$, such that $T/\sim = \{\mathcal{I}_1, \mathcal{I}_2\}$, where $\mathcal{I}_1 = \{1, \dots, 5\}$, $\mathcal{I}_2 = \{6, \dots, 10\}$. Let $T_1 = \mathcal{I}_1$, $T_2 = \mathcal{I}_2$, then $\mathcal{K}_1$ is partitioned by T_1 , $\mathcal{K}_2$ by T_2 respectively. Define a size vector $\underline{n}$, such that $n_1 = \dots = n_5 = 2, n_6 = \dots = n_{10} = 2$. Then after a sequence of reductions, we obtain a boc $\mathfrak{B}^s = \mathfrak{A}^s$ (not properly partial!) given by

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad C = \begin{pmatrix} c_3 & c_4 \\ \lambda & c_2 \end{pmatrix}, \quad D = \begin{pmatrix} d_3 & d_4 \\ d_1 & d_2 \end{pmatrix},$$

which satisfies MW5 of Theorem 5.6.1. Continue the reductions, then $C = \begin{pmatrix} \emptyset & \emptyset \\ \lambda & \emptyset \end{pmatrix}$ yields a local boc $\mathfrak{A}^{s'}$. Let $D = \begin{pmatrix} \emptyset & \emptyset \\ \eta & \emptyset \end{pmatrix}$. Thus $G = D$, and 10, the column index of G , belongs to $\mathcal{I}_2$, such that $\mathcal{I}_2 \cap T_1 = \emptyset$. Therefore the matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ satisfies S01 and S02 of 9.1. Define a new size vector $\tilde{\underline{n}}$, such that $\tilde{n}_1 = \dots = \tilde{n}_5 = 2$, $\tilde{n}_6 = \dots = \tilde{n}_{10} = 3$, and we add a column to the left of the l -th block column for

$l = 6, \dots, 10$ as follows:

$$\tilde{A} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \tilde{B} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad \tilde{C} = \begin{pmatrix} c_0 & c_3 & c_4 \\ 0 & \lambda & c_2 \end{pmatrix}, \quad \tilde{D} = \begin{pmatrix} d'_0 & d_3 & d_4 \\ d_0 & d_1 & d_2 \end{pmatrix}.$$

Then we obtain a boc $\tilde{\mathfrak{B}}^s$. If we continue the reductions up to $\tilde{C}' = \begin{pmatrix} \emptyset & \emptyset & \emptyset \\ 0 & \lambda & \emptyset \end{pmatrix}$, and obtain a layered boc $\tilde{\mathfrak{B}}^{s'}$ having two vertices :

$$\lambda \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \bullet \xrightarrow{d_0} \bullet \quad \mathcal{Q}.$$

Example 2 . Consider Example 4 in 2.1 and 4.3. A sequence of reductions leads to a minimal boc below in figure 7, where λ^0 is a fixed eigenvalue. Let us set the shadowed part $(1, \nu)$ free, then we obtain a matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$ in figure 8, where $T = \{1, 2, 3\}$ and $2 \sim 3$, $T_1 = \{1\}$, $T_2 = \{2, 3\}$;

$$K_1 = \{(s) \mid \forall s \in k\}, \quad K_2 = \left\{ \begin{pmatrix} u & 0 \\ 0 & u \end{pmatrix} \mid \forall u \in k \right\};$$

$\mathcal{M} = \{(m_1, m_2) \mid \forall m_1, m_2 \in k\}$; $\mathcal{V} = \{(0, 0)\}$. Denote the equivalent class $\{1\}$ by $\mathcal{P}$, and $\{2, 3\}$ by $\mathcal{I}$. Define a size vector $\underline{n}$, such that $n_{\mathcal{P}} = 1$, $n_{\mathcal{I}} = 1$, then $\underline{n}_1 = (1)$, $\underline{n}_2 = (1, 1)$. Let

$$N = (f_1, f_2), \quad R_1 = (x), \quad R_2 = \begin{pmatrix} y & 0 \\ 0 & y \end{pmatrix}, \quad V = (0, 0),$$

then $(N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2)$ is a triple of matrix problem $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, \mathcal{V})$, which possesses an equation

$$(x)(f_1, f_2) = (f_1, f_2) \begin{pmatrix} y & 0 \\ 0 & y \end{pmatrix}.$$

Let us define a new size vector $\tilde{\underline{n}}$, such that $\tilde{n}_{\mathcal{P}} = n_{\mathcal{P}} = 1$, $\tilde{n}_{\mathcal{I}} = n_{\mathcal{I}} + 1 = 2$. Then we add a column to the left at each block-column of the matrix $\mathcal{N} = (0, 0)$. The structure in 9.1 gives a triple of matrices $(\tilde{N}, \tilde{R}_1 \times \tilde{R}_2, \tilde{\underline{n}}_1 \times \tilde{\underline{n}}_2)$, where $\tilde{N} = (f_{11} \ f_{12} \vdots f_{21} \ f_{22})$, $\tilde{R}_1 = R_1 = (x)$,

$$\tilde{R}_2 = \left(\begin{array}{cc|cc} y_{11} & y_{12} & & \\ y_{21} & y_{22} & & \\ \hline & & y_{11} & y_{12} \\ & & y_{21} & y_{22} \end{array} \right).$$

After an edge reduction $(\overline{f_{11} \ f_{12}}) = (0 \ 1)$ in case of $\overline{f_1} = (1)$, we have $\tilde{N}' = (0 \ 1 \vdots 0 \ 0)$, $\tilde{N}' = (0 \ 0 \vdots c \ *)$, where $c = f_{21}$, $* = f_{22}$, $\tilde{R}'_1 = (x)$,

$$\tilde{R}'_2 = \left(\begin{array}{cc|cc} y_{11} & y_{12} & & \\ 0 & x & & \\ \hline & & y_{11} & y_{12} \\ & & 0 & x \end{array} \right).$$

Figure 7

Figure 8

Figure 9

Figure 9


9.3 The differentials in partial boc's $\tilde{\mathfrak{B}}^s$

Lemma 9.3.1 Let $(\alpha(0), \beta(0))$ be the matrix index of λ , and (α^0, β^0) that of ν given by Definition 9.1.1. Then $\alpha(0) > \alpha^0$ under the order of Definition 2.3.1.

Suppose that the partial bocs $\widetilde{\mathfrak{B}}^s$ has a partial layer

$$\tilde{L}^s = (\tilde{\Gamma}', \tilde{\omega}; a_1, \dots, a_j, a_{j+1}, \dots, a_{j+m}; c_1, \dots, c_r, c^0, d_1, \dots, d_n; w, u, v),$$

where $\text{ind}\tilde{\Gamma}' = \{\mathcal{P}, \mathcal{Q}\}$, $\tilde{\Gamma}'(\mathcal{P}, \mathcal{P}) = k[\lambda]$ and λ appears for the first time in $\tilde{\mathfrak{B}}^s$, $\tilde{\Gamma}'(\mathcal{Q}, \mathcal{Q}) = k$; $a_l : \mathcal{P} \longrightarrow \mathcal{P}$, $l = 1, 2, \dots, j$, are solid loops appearing before the α^0 -th row of G according to the matrix order of Definition 2.3.1, and $a_{j+1}, \dots, a_{j+m}$ are solid loops at the α^0 -th row of G ; $c_1, \dots, c_r, c^0 : \mathcal{P} \rightarrow \mathcal{Q}$ are solid edges sitting at the added columns up to the α^0 -th row of G , and the last one c^0 is sitting at the intersection of the o_{j^0} -th column and the α^0 -th row; $d_1, \dots, d_n$ are all the solid arrows after a_{j+m} ; $w : \mathcal{P} \rightarrow \mathcal{P}$, $u : \mathcal{P} \rightarrow \mathcal{Q}$ are dotted arrows (see the figures below).



Where $l, l' \in \mathcal{L}^0$, $a_{j+1}, \dots, a_{j+m}$ belong to the area A .

Lemma 9.1.1 and 9.1.2 enables us to calculate the differentials of the solid arrows of $\tilde{\mathfrak{B}}^s$. Let

$$\begin{cases} \delta(a_1)^0 &= h_{11}(\lambda, \mu)w_1 \\ \dots & \dots \\ \delta(a_j)^0 &= h_{j1}(\lambda, \mu)w_1 + \dots + h_{jj}(\lambda, \mu)w_j \end{cases} \quad (57)$$

be given by Formula (42) of 5.4 up to j , then $h_{ll}(\lambda, \lambda) \neq 0$, (see part A in the figure).

On the other hand, part C in the figure gives

$$\begin{cases} \delta(c_1)^0 &= f_{11}(\lambda)u_1 \\ \delta(c_2)^0 &= f_{21}(\lambda)u_1 + f_{22}(\lambda)u_2 \\ \dots & \dots \\ \delta(c_r)^0 &= f_{r1}(\lambda)u_1 + f_{r2}(\lambda)u_2 + \dots + f_{rr}(\lambda)u_r \end{cases} \quad (58)$$

where $u_{r(1)} = 0, \dots, u_{r(\beta)} = 0$ for some indices $1 \leq r(1) < \dots < r(\beta) \leq r$, and the dotted arrows u_l , $\forall l \in I^0$, are linearly independent, where $I^0 = \{1, 2, \dots, r\} \setminus \{r(1), \dots, r(\beta)\}$.

If

$$\tilde{R}_{ll}^s = \left(\begin{array}{c|cccc} x_{00} & x_{01} & x_{02} & \dots & x_{0m} \\ \hline 0 & & R_{ll}^s & & \end{array} \right), \text{ then } \begin{cases} \delta(a_{j+1}) &= c^0 x_{01} + \dots \\ \delta(a_{j+2}) &= c^0 x_{02} + \dots \\ \dots & \dots \\ \delta(a_{j+m}) &= c^0 x_{0m} + \dots \end{cases} \quad (59)$$

Define a polynomial:

$$h^0(\lambda, \mu) = \prod_{l=1}^j c_l(\lambda) h_{ll}(\lambda, \mu) \cdot \prod_{l \in I^0} f_l(\lambda). \quad (60)$$

(see Formula (30) of 4.3 and Formula (43) of 5.4.) Let $h^0(\lambda, \lambda) \neq 0$, we set $a_l = \emptyset$, for $l = 1, \dots, j$; $c_l = \emptyset$ for $l \in I^0$, $c_{r(l)} = 0$ for $l = 1, \dots, \beta$, then we obtain an induced partial bocs $\tilde{\mathfrak{B}}^{s+j+r}$. We will prove in the next subsection that $\delta(c^0) = 0$ in $\tilde{\mathfrak{B}}^{s+j+r}$. Let $c^0 = 1$; $a_{j+l} = \emptyset$, and $x_{o_l} + \dots = 0$ in Formula (59) for $l = 1, \dots, m$. Then we obtain an induced partial bocs $\tilde{\mathfrak{B}}^{s+J}$ with $J = j + r + 1 + m$, which is parameterized, but not necessarily

freely parameterized in general, because there is no guarantee that $h^0(\lambda, \mu)$ satisfies item (3) of Definition 4.2.1.


Lemma 9.3.2 $\tilde{\mathfrak{B}}^{s+J} = (\tilde{\Gamma}^{s+J}, \tilde{\Omega}^{s+J})$ is an induced boc of $\tilde{\mathfrak{B}} = (\tilde{\Gamma}, \tilde{\Omega})$, such that

- (1) $\tilde{\mathfrak{B}}^{s+J}$ is local;
- (2) $\tilde{\mathfrak{B}}^{s+J}$ possess a unique parameter λ with the domain $h^0(\lambda, \mu) \neq 0$ given in Formula (60), which is not necessarily freely parameterized;
- (3) $\tilde{\Gamma}^{s+J}$ is freely generated by $d_1, d_2, \dots, d_n$ over $k[\lambda, \mu, h^0(\lambda, \mu)^{-1}]$;
- (4) $\tilde{\Omega}^{s+J}$ is obtained from $\tilde{\Omega}$ by the restriction $w_l = 0$ for $1 \leq l \leq j$, (Formula(57)), $u_l = 0$ for any $l \in I^0$ (Formula(58)), and $x_{ol} + \dots = 0$ for $1 \leq l \leq m$ (Formula(59)). $\square$

Let T_1, T_2 be the index set of $\mathfrak{B}$, and let $\mathcal{S} = \{1, 2, \dots, n_1\}$, if the number of the rows of matrix N is n_1 . We define a set of pairs of indices

$$\{(\alpha, l) \mid \alpha \in \mathcal{S}, l \in T_2\}.$$

And we give an order in the set, such that $(\alpha_1, l_1) \prec (\alpha_2, l_2)$, provided $\alpha_1 > \alpha_2$, or $\alpha_1 = \alpha_2$ but $l_1 < l_2$, (see the following two examples).



We specify a pair of indices (α^0, l^0) for ν given by Definition 9.1.1. And we also specify a pair $(\alpha(0), l(0))$ for λ , where $\alpha(0)$ is the row-index of λ in the matrix; $l(0)$ the index of the block-column partitioned by $(T, \sim)$, such that λ sits in that block.

Next we use the procedure of Lemma 5.3.1 once more for the local partial boc $\tilde{\mathfrak{B}}^{s+J}$ to get a sequence of parameters $\lambda = \lambda_{10}, \lambda_{11}, \dots, \lambda_{1\gamma^1}$.

Proposition 9.3.1 (1) If $\gamma^1 = 0$, we have

$$\begin{cases} \delta(d_1)^0 &= g_{11}(\lambda, \mu)v_1 \\ \delta(d_2)^0 &= g_{21}(\lambda, \mu)v_1 + g_{22}(\lambda, \mu)v_2 \\ \dots &\dots \dots \\ \delta(d_n)^0 &= g_{n1}(\lambda, \mu)v_1 + g_{n2}(\lambda, \mu)v_2 + \dots + g_{nn}(\lambda, \mu)v_n \end{cases} \quad (61)$$

where $g_{ll}(\lambda, \mu) \neq 0$ (see Formula (40) and (41) in 5.4, and we use λ, μ instead of ν, κ).

(2) If $\gamma^1 = 1$, denote λ_{11} by ν_1 . Then $(\alpha^1, l^1) \succ (\alpha^0, l^0)$.

(3) If $\gamma^1 \geq 2$, denote λ_{1,γ^1-1} by λ_1 , λ_{1,γ^1} by ν_1 respectively. Then $(\alpha(1), l(1)) \succ (\alpha(0), l(0))$.


Proof. (2) If $\gamma^1 = 1$, then ν_1 is located outside of G or above the α^0 -row of G by Formula (59). Thus $(\alpha^1, l^1) \succ (\alpha^0, l^0)$.

(3) If $\gamma^1 \geq 2$, then λ_1 is located outside of G or above the α^0 -row of G still by Formula (59), thus $(\alpha(1), l(1)) \succ (\alpha^0, l^0)$. On the other hand, $(\alpha^0, l^0) \succ (\alpha(0), l(0))$ by Lemma 9.3.1. Consequently $(\alpha(1), l(1)) \succ (\alpha(0), l(0))$. $\square$

9.4 The calculation of $\delta(c^0)$

In this subsection we will prove that $\delta(c^0) = 0$ in $\tilde{\mathfrak{B}}^{s+j+r}$ under the assumption S01 and S02 of structure 9.1.

Given any $l \in \mathcal{L}^0$, we define a matrix index $\beta_l \in l$, such that the distance from the β_l -th column to the left edge of the l -th block-column is the same as the distance from the β^0 -th column to the left edge of l^0 -th block-column. And $\beta_{l^0} = \beta^0$.



Recall the matrix equation $R_1 \mathcal{N}^s = V + \mathcal{N}^s R_2$ from 9.1. If eq is an equation, eq^R stands for the right-hand side of eq given by $V + \mathcal{N}^s R_2$, and if Eq is a set of equations, Eq^R stands for the set of the right-hand side of the equations in Eq . Recall from Lemma 9.1.2, that $eq_{(\alpha,\beta)}^R$ contains only the indeterminates at the β -th column of R_2 and V for any pair of indices (α, β) .

Lemma 9.4.1 Let (α^0, β^0) be the index of ν given in Definition 9.2.1. Then $eq_{(\alpha^0, \beta^0)}^R$ is a linear combination of the forms of Eq^R , where $Eq = Eq_{(\alpha^0, \beta_0)}$ is given by Definition 9.1.2.

Proof. (1) We set $\lambda = \mu$, then $eq_{(\alpha^0, \beta^0)}^c$ is a linear combination of the equations of Eq^c in $\mathfrak{B}^{s+i-1}$ by Lemma 9.1.1 and Formula (42) of 5.4, where eq^c stands for the equation eq when we do not distinguish the multiplication of the parameters from left or right.

(2) Let X_1 be the set of indeterminates at the l -th block column of R_2 and V for any $l \in \mathcal{L}^0$, and X_2 be that at the j -th block column of R_1 , R_2 and V for any $j \in T \setminus \mathcal{L}^0$.

Thus $X_1 \dot{\cup} X_2$ is the complete set of the indeterminates. Deleting X_2 from $eq_{(\alpha^0, \beta^0)}^c$ and Eq^c , we obtain an equation $eq'_{(\alpha^0, \beta^0)}^c$ and a set of equations Eq'^c . It is straightforward that $eq'_{(\alpha^0, \beta^0)}^c$ is the same linear combination of the equations in Eq'^c .

(3) The left-hand sides of $eq'_{(\alpha^0, \beta^0)}^c$ and the equations in Eq'^c are all zero by assumption S02 of 9.1. And the right-hand side of $eq'_{(\alpha^0, \beta^0)}^c$ is just $eq_{(\alpha^0, \beta^0)}^R$, and that of any equation in Eq'^c is either a form in Eq^R containing the indeterminates at the β -th column for some $\beta \in l \in \mathcal{L}^0$ or zero. Therefore $eq_{(\alpha^0, \beta^0)}^R$ is a linear combination of the forms of Eq^R . The proof is completed. $\square$

Lemma 9.4.2 In the matrix equation $\tilde{R}_1 \tilde{\mathcal{N}}^s = \tilde{V} + \tilde{\mathcal{N}}^s \tilde{R}_2$, $\tilde{eq}_{(\alpha^0, o_{l^0})}^R$ is a linear combination of the forms in $\tilde{Eq}^R$, where $\tilde{Eq} = \tilde{Eq}_{(\alpha^0, o_{l^0})}$ is given by Definition 9.1.2.

Proof. (1) Since for any pair of indices (α, β) of $\mathcal{N}^s$, $eq_{(\alpha, \beta)}$ is the same as $\tilde{eq}_{(\alpha, \beta)}$ by Lemma 9.1.2, it is also the case for the right-hand sides of the equations. Because $eq_{(\alpha^0, \beta^0)}^R$ is a linear combination of the forms in Eq^R by Lemma 9.4.1, so is $\tilde{eq}_{(\alpha^0, \beta^0)}^R$ in $\tilde{Eq}^R$.

(2) For any $\alpha \geq \alpha^0$, and $l \in \mathcal{L}^0$, the linear form of the (α, o_l) -th entry of $\tilde{V} + \tilde{\mathcal{N}}^s \tilde{R}_2$ in the indeterminates at the o_l -th column of $\tilde{V}$ and $\tilde{R}_2$ is the same as the linear form of the (α, β_l) -th entry in the indeterminates at the β_l -th column by Lemma 8.5.1, where $\beta_l \in l$ given by the above diagram.

Combining (1) and (2), $\tilde{eq}_{(\alpha^0, o_{l^0})}^R$ is a linear combination of the forms in $\tilde{Eq}^R$ in the indeterminates of the o_l -th column of $\tilde{R}_2$ and $\tilde{V}$ for $l \in \mathcal{L}^0$. $\square$

Proposition 9.4.1 $\tilde{eq}_{(\alpha^0, o_{l^0})}$ is a linear combination of the equations in $\tilde{Eq}$.

Proof. Since the left-hand sides of all the equations $\tilde{eq}_{(\alpha, o_l)}$ for $\alpha \geq \alpha^0$ and $l \in \mathcal{L}^0$ equal zero, $\tilde{eq}_{(\alpha^0, o_{l^0})}$ is a linear combination of equations of $\tilde{Eq}$ by Lemma 9.4.2. $\square$

Corollary 9.4.1 $\delta(c^0) = 0$ in $\tilde{\mathfrak{B}}^{s+j+r}$.

Proof. By Proposition 9.4.1 and Lemma 9.1.1. $\square$

9.5 The induction on (α^η, l^η)

Suppose we are given $\mathfrak{B}^s, \tilde{\mathfrak{B}}^s, \tilde{\mathfrak{B}}^{s+J}$ in 9.3, then $\tilde{\mathfrak{B}}^{s+J}$ is local by Lemma 9.3.2. Assume that when we perform the procedure 5.3 starting from $\tilde{\mathfrak{B}}^{s+J}$, $\gamma^1 = 1$ in Proposition 9.3.1, and $\tilde{\mathfrak{B}}^{s+J}$ possesses the following formulae of 5.4

$$\begin{cases} \delta(a_1^1)^0 &= h_{11}^1(\lambda, \mu)w_1^1 \\ \dots & \dots \dots \\ \delta(a_{i^1-1}^1)^0 &= h_{i^1-1,1}^1(\lambda, \mu)w_1^1 + \dots + h_{i^1-1,i^1-1}^1(\lambda, \mu)w_{i^1-1}^1 \\ \delta(a_{i^1}^1)^0 &= h_{i^1,1}^1(\lambda, \mu)w_1^1 + \dots + h_{i^1,i^1-1}^1(\lambda, \mu)w_{i^1-1}^1 + \tilde{h}^1(\lambda, \mu)\tilde{w}^1 \end{cases} \quad (62)$$

where $a_l^1 = d_l$ for $l = 1, \dots, i^1$, $\tilde{w}^1$ is linearly independent of $w_1^1, \dots, w_{i^1-1}^1$, and $(\lambda - \mu) \mid \tilde{h}^1(\lambda, \mu)$, $(\lambda - \mu)^2 \nmid \tilde{h}^1(\lambda, \mu)$. Let

$$h^1(\lambda, \mu) = h^0(\lambda, \mu) \prod_{l=1}^{j^1} c_l^1(\lambda) h_{ll}^1(\lambda, \mu) \cdot \prod_{l \in I^1} f_l^1(\lambda) \quad (63)$$

For any fixed $\lambda^0 \in k$ with $h^1(\lambda^0, \lambda^0) \neq 0$, set $a_l^1 = \emptyset$, $l = 1, \dots, (i^1 - 1)$, $a_{i^1}^1 = \nu_1$, then

$$\begin{cases} \delta(d_1^1)^0 &= g_{11}^1(\nu_1, \kappa_1)v_1^1 \\ \dots & \dots \dots \\ \delta(d_{n^1-1}^1)^0 &= g_{n^1-1,1}^1(\nu_1, \kappa_1)v_1^1 + \dots + g_{n^1-1,n^1-1}^1(\nu_1, \kappa_1)v_{n^1-1}^1 \\ \delta(d_{n^1}^1)^0 &= g_{n^1,1}^1(\nu_1, \kappa_1)v_1^1 + \dots + g_{n^1,n^1-1}^1(\nu_1, \kappa_1)v_{n^1-1}^1 + g_{n^1,n^1}^1(\nu_1, \kappa_1)v_{n^1}^1 \end{cases}$$

where $d_l^1 = d_{i^1+l}$, $n^1 = n - i^1$, (see Formula (40) of 5.4), and $\sigma_l^1(\nu_1)$ are given by (41) of 5.4, $g_{ll}^1(\nu_1, \kappa_1) \in k[\nu_1, \kappa_1, \sigma_l^1(\nu_1)^{-1}, \sigma_l^1(\kappa_1)^{-1}]$ are all invertible.

Suppose ν_1 is contained in a block G^1 partitioned by $(\tilde{T}, \sim)$, and l^1 , the index of the block-column of G^1 , belongs to $\mathcal{L}^1$. We propose the following definition and assumption.

Definition 9.5.1 A local boc $\mathcal{A}$ (not necessarily layered) is said to have generalized MW5, if $\mathcal{A}$ has a unique parameter λ with a domain $g(\lambda, \mu) \neq 0$, such that Formula (42) of Lemma 5.4.1 holds, and for any fixed $\lambda^0 \in k$ with $g(\lambda^0, \lambda^0) \neq 0$, all the $g_l(\nu, \kappa) \in k[\nu, \kappa, \sigma_l(\nu)^{-1}, \sigma_l(\kappa)^{-1}]$ are invertible in Formula (41). $\square$

Our assumptions are:

S11 $\gamma^1 = 1$, the end term $\tilde{\mathfrak{B}}^{s+J}$ of the sequence of partial bocses is in the case of generalized MW5.

S12 $l^1 \in \mathcal{L}^1 \in \tilde{T}_2/\sim_2$, such that $\mathcal{L}^1 \cap \tilde{T}_1 = \emptyset$.

From now on, we write $({}^1N^r, {}^1R_1^r \times {}^1R_2^r, {}^1\underline{n}_1^r \times {}^1\underline{n}_2^r)^r$ instead of $(\tilde{N}^r, \tilde{R}_1^r \times \tilde{R}_2^r, \tilde{\underline{n}}_1^r \times \tilde{\underline{n}}_2^r)$ and sequence $({}^1\Delta)$ instead of $(\tilde{\Delta})$. We will construct a new sequence $({}^2\Delta)$ based on $({}^1\Delta)$ similarly to structure 9.1 as follows.

First we define a new size vector ${}^2\underline{n}$ of $({}^1T, {}^1\sim)$, such that ${}^2n_{\mathcal{L}^1} = {}^1n_{\mathcal{L}^1} + 1$, and ${}^2n_{\mathcal{I}} = {}^1n_{\mathcal{I}}$, $\forall I \in {}^1T/{}^1\sim \setminus \{\mathcal{L}^1\}$. Then we add a column consisting of 0's and $\emptyset$'s at the left of the l -th block-column for each $l \in \mathcal{L}^1$ into ${}^1\mathcal{N}^s$. Thus ${}^2T = {}^1T \cup \mathcal{Q}^1$ where $\mathcal{Q}^1 =$

$\{o_1^1, o_2^1, \dots, o_{l_{\mathcal{L}^1}}^1\}$ is the set of the indices of the added columns of $\mathcal{L}^1 = \{l_1^1, l_2^1, \dots, l_{\mathcal{L}^1}^1\}$. And ${}^2T^s$ has either equivalent classes $\mathcal{P}, \mathcal{Q}, \mathcal{Q}_1$ with $Q_1 = Q^1$ when $\mathcal{L}^1 \neq \mathcal{L}^0$; or $\mathcal{P}, \mathcal{Q}_1$ with $Q_1 = \mathcal{Q} \cup Q^1$ when $\mathcal{L}^1 = \mathcal{L}^0$. The formula (57) and (59) of 9.3 do not change in ${}^2\mathfrak{B}^s$. When $\mathcal{L}^1 \neq \mathcal{L}^0$, Formulae (58) remains, (see Figure 1 below); when $\mathcal{L}^1 = \mathcal{L}^0$, $c_l, u_l, l = 1, \dots, r$ and c^0 change to 1×2 matrices with the unchanged differentials of c_l given in Formulae (58), (see Figure 2).




figure 1




figure 2

Denote by c^1 the $(\alpha^1, o_{l_1}^1)$ -entry; and by C^1 the set of entries at $o_{l_1}^1$ -th column before c^1 (and after c^0 in the case of $\mathcal{L}^1 = \mathcal{L}^0$). Let $A^1 \subset D$ up to the (α^1, l^1) -th row block. The differentials of the solid arrows before the (α^1, l^1) -th row in A^1 , are given by Formula (62) and we may denote it by $(57)^1$; Those at the (α^1, l^1) -th row of A^1 are parallel to Formula (59) of 9.3, we may denote it by $(59)^1$. And we also have Formula $(58)^1$ for the entries in the area C^1 . Let $h^1(\lambda, \lambda) \neq 0$, we set $a \in A \cup A^1$ before c^1 to be $\emptyset$; $c \in C \cup C^1$ to be $\emptyset$ or 0; $c^0 = (1)$ when $\mathcal{L}^1 \neq \mathcal{L}^0$, or $c^0 = (0 \ 1)$ when $\mathcal{L}^1 = \mathcal{L}^0$. We conclude that $\delta(c^1) = 0$ in the induced partial boc ${}^2\mathfrak{B}^{s+J+j^1+r^1}$ under the assumption S11, S12 similarly to Corollary 9.4.1. Let $c^1 = (1)$, set the solid arrows at the (α^1, l^1) -th row of A^1 to be ϕ , then we obtain an induced boc ${}^2\mathfrak{B}^{s+J+J^1}$, where $J^1 = j^1 + r^1 + 1 + m^1$. The local boc ${}^2\mathfrak{B}^{s+J+J^1}$ may not be freely parameterized in general, because $h^1(\lambda, \mu)$ does not necessarily satisfy the item (3) of Definition 4.2.1. Then we perform procedure 5.3 starting from the local boc ${}^2\mathfrak{B}^{s+J+J^1}$ once again, and obtain a sequence of parameters $\lambda = \lambda_{20}, \lambda_{21}, \dots, \lambda_{2\gamma^2}$. If $\gamma^2 = 1$ and ${}^2\mathfrak{B}^{s+J+J^1}$ satisfies the assumption S21, S22 parallel to S11, S12, we may continue to use the structure of 9.1 once more.

Now suppose we have a chain of indices:

$$(\alpha^0, l^0) \prec (\alpha^1, l^1) \prec \dots \prec (\alpha^{p-1}, l^{p-1}) \prec (\alpha^p, l^p),$$

such that $\gamma^1 = 1, \dots, \gamma^{p-1} = 1$, and we have a chain of index sets

$$T \subseteq {}^1T \subseteq {}^2T \subseteq \dots \subseteq {}^pT$$

such that ${}^qT = T \cup \mathcal{Q} \cup \mathcal{Q}^1 \cup \dots \cup \mathcal{Q}^{q-1}$ for $0 \leq q \leq p$, where $\mathcal{Q}^{q-1} = \{o_1^{q-1}, o_2^{q-1}, \dots, o_{\mathcal{L}^{q-1}}^{q-1}\}$ is the set of the indices of the added columns if $\mathcal{L}^{q-1} = \{l_1^{q-1}, l_2^{q-1}, \dots, l_{\mathcal{L}^{q-1}}^{q-1}\}$. And ${}^qT^s$ possesses the equivalent classes $\mathcal{P}$ and $\mathcal{Q}_{q1}, \dots, \mathcal{Q}_{qr_q}$, which are unions of some $\mathcal{Q}, \mathcal{Q}^1, \dots, \mathcal{Q}^{q-1}$. Moreover we have a chain of reduction sequences:

$$(\Delta), ({}^1\Delta), ({}^2\Delta), \dots, ({}^p\Delta);$$

a chain of the corresponding sequence of partial bocses

$$(***), ({}^1***), ({}^2***), \dots, ({}^p***);$$

and a chain of the end terms of the preceding sequences:

$$\mathfrak{B}^s, {}^1\mathfrak{B}^s, {}^2\mathfrak{B}^s, \dots, {}^p\mathfrak{B}^s;$$

such that all the local partial bocses

$$\mathfrak{B}^s, {}^1\mathfrak{B}^{s+J}, {}^2\mathfrak{B}^{s+J+J^1}, \dots, {}^{(p-1)}\mathfrak{B}^{s+\sum_{\eta=0}^{p-2} J^\eta}$$

satisfy the assumption S01, S02; S11, S12; S21, S22; $\dots$; S(p-1)1, S(p-1)2 inductively, where $J^\eta = j^\eta + r^\eta + 1 + m^\eta$. We perform the procedure 5.3 for the local boc ${}^p\mathfrak{B}^{s+\sum_{\eta=0}^{p-1} J^\eta}$ induced from ${}^p\mathfrak{B}^s$, and obtain a sequence of parameters $\lambda = \lambda_{p0}, \lambda_{p1}, \dots, \lambda_{p\gamma^p}$. If $\gamma^p = 1$, let

$$h^p(\lambda, \mu) = h^{p-1}(\lambda, \mu) \prod_{l=1}^{j^p} c_l^p(\lambda) h_{ll}^p(\lambda, \mu) \prod_{l \in I^p} f_l^p(\lambda); \quad (64)$$

If ${}^p\mathfrak{B}^{s+\sum_{\eta=0}^{p-1} J^\eta}$ still satisfies the assumption SP1, SP2, and we set $\nu_p = \lambda_{p1}$, then $(\alpha^p, l^p) \succ (\alpha^{p-1}, l^{p-1})$. Adding a column to the left of the l -th block column of ${}^p\mathcal{N}^s$ for all $l \sim l^p$, we obtain a boc ${}^{(p+1)}\mathfrak{B}^s$. Such a procedure must stop at some stage since the index set $\{(\alpha^q, l^q)\}$ is bounded by (n_1, t_2) , (see the explanation below Lemma 9.3.2).

Proposition 9.5.1 We perform the procedure of 5.3 starting from boc ${}^p\mathfrak{B}^{s+\sum_{\eta=0}^{p-1} J^\eta}$. If $\gamma^p \neq 1$, or $\gamma^p = 1$ but ${}^p\mathfrak{B}^{s+\sum_{\eta=0}^{p-1} J^\eta}$ does not satisfy the assumption SP1 and SP2, then we must have one of the following possibilities.

Q1. $\gamma^p = 0$, then Formula (60) of 9.3 becomes

$$\begin{cases} \delta(d_1^{p-1})^0 &= g_{11}^{p-1}(\lambda, \mu) v_1^{p-1} \\ \delta(d_2^{p-1})^0 &= g_{21}^{p-1}(\lambda, \mu) v_1^{p-1} + g_{22}^{p-1}(\lambda, \mu) v_2^{p-1} \\ \dots &\dots \dots \\ \delta(d_{n^{p-1}}^{p-1})^0 &= g_{n^{p-1}1}^{p-1}(\lambda, \mu) v_1^{p-1} + g_{n^{p-1}2}^{p-1}(\lambda, \mu) v_2^{p-1} + \dots + g_{n^{p-1}n^{p-1}}^{p-1}(\lambda, \mu) v_{n^{p-1}}^{p-1} \end{cases}$$

and

$$\sigma_l^{p-1}(\lambda) = h^{p-1}(\lambda, \lambda) \prod_{l \in I^p} f_l^p(\lambda) \prod_{q=1}^l c_q^{p-1}(\lambda), \text{ write } \sigma_{n^{p-1}}^{p-1}(\lambda) = \sigma^{p-1}(\lambda), \quad (65)$$

(see (41) of 5.3), where $v_1^{p-1}, v_2^{p-1}, \dots, v_{n^{p-1}}^{p-1}$ are linearly independent, and $g_{ll}^{p-1}(\lambda, \lambda) \neq 0$.

(1) There exists some $\lambda^0 \in k$ with $\sigma_l^{p-1}(\lambda^0) \neq 0$ such that $g_{ll}^{p-1}(\lambda, \mu) \in k[\lambda, \mu, \sigma_l^{p-1}(\lambda)^{-1} \sigma_l^{p-1}(\mu)^{-1}]$ for $1 \leq l \leq e-1$ are all invertible, but $g_{ee}^{p-1}(\lambda, \mu) \in k[\lambda, \mu, \sigma_e^{p-1}(\lambda)^{-1} \sigma_e^{p-1}(\mu)^{-1}]$ is not invertible.

(2) $\forall \lambda^0 \in k$ with $\sigma_l^{p-1}(\lambda^0) \neq 0$, $g_{ll}^{p-1}(\lambda, \mu) \in k[\lambda, \mu, \sigma_l^{p-1}(\lambda)^{-1} \sigma_l^{p-1}(\mu)^{-1}]$ are all invertible for $1 \leq l \leq n^{p-1}$.

Q2. $\gamma^p = 1$, then Formula (62) of 9.5 becomes

$$\begin{cases} \delta(a_1^p)^0 &= h_{11}^p(\lambda, \mu) w_1^p \\ \dots & \dots \dots \\ \delta(a_{i^p-1}^p)^0 &= h_{i^p-1,1}^p(\lambda, \mu) w_1^p + \dots + h_{i^p-1,i^p-1}^p(\lambda, \mu) w_{i^p-1}^p \\ \delta(a_{i^p}^p)^0 &= h_{i^p,1}^p(\lambda, \mu) w_1^p + \dots + h_{i^p,i^p-1}^p(\lambda, \mu) w_{i^p-1}^p + \tilde{h}^p(\lambda, \mu) \tilde{w}^p \end{cases}$$

where $\tilde{w}^p$ is a linear combination of $w_1^p, \dots, w_{i^p-1}^p$ or $\tilde{w}^p$ is linearly independent of $w_1^p, \dots, w_{i^p-1}^p$, but $(\lambda - \mu)^2 \mid \tilde{h}^p(\lambda, \mu)$.

Moreover for any fixed $\lambda^0 \in k$ with $h^p(\lambda^0, \lambda^0) \neq 0$, if we set $a_l^p = \emptyset$, $l = 1, \dots, i^p - 1$, and $a_{i^p}^p = \nu_p$, and write $d_l^p = a_{i^p+l}^p$, then we have the following formula:

$$\begin{cases} \delta(d_1^p)^0 &= g_{11}^p(\nu_p, \kappa_p) v_1^p \\ \dots & \dots \dots \\ \delta(d_{n^p}^p)^0 &= g_{n^p,1}^p(\nu_p, \kappa_p) v_1^p + \dots + g_{n^p,n^p}^p(\nu_p, \kappa_p) v_{n^p}^p \end{cases}$$

Let

$$\sigma_l^p(\nu_p) = h^p(\nu_p, \nu_p) \prod_{q=1}^l c_q^p(\nu_p), \text{ write } \sigma^p(\nu_p) = \sigma_{n^p}^p(\nu_p)$$

be given by Formula (64) of 9.5 and (41) of 5.4, then $\forall \lambda^0 \in k$ with $h^p(\lambda^0, \lambda^0) \neq 0$, $g_{ll}^p(\nu_p, \kappa_p) \in k[\nu_p, \kappa_p, \sigma_l^p(\nu_p)^{-1} \sigma_l^p(\kappa_p)^{-1}]$ are all invertible.

Q3. $\gamma^p \geq 2$, then we have a sequence of parameters $\lambda = \lambda_{p0}, \lambda_{p1}, \dots, \lambda_{p\gamma^p}$. If we set $\lambda_1 = \lambda_{p,\gamma^p-1}$ having a pair of indices $(\alpha(1), l(1))$, then $(\alpha(1), l(1)) > (\alpha(0), l(0))$.

9.6 Non-homogeneous partial boc ${}^p\mathfrak{B}^s$

We will prove that the partial boc ${}^p\mathfrak{B}^s$ given in Q1 and Q2 of Proposition 9.5.1 is not homogeneous in this subsection.

In case of Q1, let $\lambda^0 \in k$ with $\sigma_e^{p-1}(\lambda^0) \neq 0$ in item (1), or $\sigma^{p-1}(\lambda^0) \neq 0$ in item (2), (see Formula (65)). We define an object $S \in R({}^p\mathfrak{B}^s)$, such that $S_{\mathcal{P}} = k$; $S_{Q^q} = k$ for $0 \leq q < p$; $S(\lambda) = (\lambda^0)$, $S(a) = \emptyset$, $\forall a \in \bigcup_{\eta=0}^{p-1} A^\eta$; $S(c) = \emptyset$ or 0 , $\forall c \in \bigcup_{\eta=0}^{p-1} C^\eta$; $S(c^\eta) = (0 \cdots 1)$ for $0 < \eta \leq p-1$; $S(d) = \emptyset$, $\forall d \in D^{p-1}$.

Lemma 9.6.1 Suppose we are given a partial boc ${}^p\mathfrak{B}^s$ satisfying Q1 of Proposition 9.5.1 and the object $S \in R({}^p\mathfrak{B}^s)$ defined above. If $(e) : S \xrightarrow{\iota} E \xrightarrow{\pi} S$ is an almost split conflation in $R({}^p\mathfrak{B}^s)$, then $E_{\mathcal{P}} = k^2$; $E_{\mathcal{Q}^q} = k^2$;

$$E(\lambda) = J_2(\lambda^0); E(a) = \emptyset; E(c) = \emptyset \text{ or } 0, E(c^\eta) = (0 \cdots 0 I_2),$$

and ι, π are given by Corollary 6.4.1.

Proof. The eigenvalue of E is λ^0 , since the sequence $0 \longrightarrow S \xrightarrow{\iota_0} E \xrightarrow{\pi_0} S \longrightarrow 0$ is exact over $k[\lambda, \sigma_e^{p-1}(\lambda)^{-1}]$ or $k[\lambda, \sigma^{p-1}(\lambda)^{-1}]$

Suppose we have some $c_i = c^q$ or $c_i \in C^q$ with $S(c_i) \neq \emptyset$, such that $E(a) = \emptyset$, $\forall a \in \bigcup_{\eta=0}^{p-1} A^\eta$ before c_i ; $E(c) = \emptyset$ or 0 , $\forall c \in \bigcup_{\eta=0}^{p-1} C^\eta$ before c_i ; $E(c^\eta) = (0 \cdots 0 I_2)$ for any c^η before c_i . We claim that $E(c_i) = (0 \cdots 0 I_2)$ when $S(c_i) = (0 \cdots 0 1)$ by item (1) of Lemma 8.5.2; and

$$E(c_i) = 0 \text{ or } \begin{pmatrix} 0 & \cdots & 0 & 0 & 1 \\ 0 & \cdots & 0 & 0 & 0 \end{pmatrix}$$

when $S(c_i) = 0$ by item (2) of Lemma 8.5.2. In the second case, we define a full subcategory $\mathcal{C}$ of $R({}^p\mathfrak{B}^s)$ consisting of the objects M such that $M_{\mathcal{P}} = k^m$, $M_{\mathcal{Q}^q} = k^m$, $0 \leq q < p$, the eigenvalue of $M(\lambda)$ is λ^0 , $M(a) = \emptyset$, $M(c) = \emptyset$ or 0 , $M(c^\eta) = (0, \cdots, 0, I_m)$ for all the a, c, c^η before c_i . We take an object $L \in \mathcal{C}$, such that $L_{\mathcal{P}} = k^2$, $L_{\mathcal{Q}^q} = k^2$, $0 \leq q < p$; $L(\lambda) = J_2(\lambda^0)$; $L(a) = \emptyset$, $\forall a \in \bigcup_{\eta=0}^{p-1} A^\eta$ before c_i ; $L(c) = \emptyset$ or 0 , $\forall c \in \bigcup_{\eta=0}^{p-1} C^\eta$ before c_i ; $L(c^\eta) = (0, \cdots, 0, I_2)$ for any c^η before c_i ; and $L(c_i) = 0$. We also define a morphism $\varphi : L \rightarrow S$, such that $\varphi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\varphi_{\mathcal{Q}^q} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ for $0 \leq q < p$. If (e) is an almost split conflation in $R({}^p\mathfrak{B}^s)$, it is so in $\mathcal{C}$. But Lemma 8.5.3 leads to a contradiction in case of

$$E(c_i) = \begin{pmatrix} 0 & \cdots & 0 & 0 & 1 \\ 0 & \cdots & 0 & 0 & 0 \end{pmatrix}.$$

Thus $E(c_i) = 0$. Our conclusion on $E(a)$, $E(c)$ and $E(c^\eta)$ follows by induction.

Finally, $E(\lambda)$ must equal to $J_2(\lambda^0)$, otherwise $E(\lambda) = \lambda^0 I_2$ would lead to (e) to be split. The proof is finished. $\square$

Proposition 9.6.1 The partial boc ${}^p\mathfrak{B}^s$ given by condition Q1 of Proposition 9.5.1 is not homogeneous.

Proof. We take an infinite list of objects S_{λ^0} defined in the beginning of 9.6. If $R({}^p\mathfrak{B}^s)$ is homogeneous, there exists a cofinite subset $D_0 \subset k \setminus \{\text{the roots of } \sigma^{p-1}(\lambda)\}$ (see Formula (65) of 9.5), such that S_{λ^0} is homogeneous for any $\lambda^0 \in D_0$. Fix such a λ^0 we denote S_{λ^0} by S for simplicity. If $(e) S \xrightarrow{\iota} E \xrightarrow{\pi} S$ is an almost split conflation starting and ending at S in $R({}^p\mathfrak{B}^s)$, then Lemma 9.6.1 gives the structure of E .

(1) In case of item (1) of Q1, we define an object $L \in R({}^p\mathfrak{B}^s)$, such that $L_{\mathcal{P}} = k^2$, $L_{\mathcal{Q}^q} = k^2$;

$$L(\lambda) = \begin{pmatrix} \lambda^0 & 0 \\ 0 & \mu^0 \end{pmatrix} \quad \text{with } g_{ee}^{p-1}(\lambda^0, \mu^0) = 0, \quad \sigma^{p-1}(\lambda^0)\sigma_e^{p-1}(\mu^0) \neq 0;$$

$L(a) = 0, \forall a \in \cup_{\eta=0}^{p-1} A^\eta$; $L(c) = 0, \forall c \in \cup_{\eta=0}^{p-1} C^\eta$; $L(c^\eta) = (0, \dots, 0, I)$ for $0 \leq \eta \leq p-1$; $L(d) = 0, \forall d \in D^{p-1}$. We also define a morphism $\varphi : L \rightarrow S$, such that $\varphi_{\mathcal{P}} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $\varphi_{\mathcal{Q}^q} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. Then a contradiction appears similarly to the proof of Proposition 7.4.1.

(2) In case of item (2) of Q1, we recall that ${}^pT^s$ has the equivalent classes $\mathcal{P}; \mathcal{Q}_1, \dots, \mathcal{Q}_r$, where $\mathcal{P}$ is the unique equivalent class of $\mathfrak{B}^s$, $\mathcal{Q}_1, \dots, \mathcal{Q}_r$ are some unions of $\mathcal{Q}, \mathcal{Q}^1, \mathcal{Q}^2, \dots, \mathcal{Q}^{p-1}$ respectively. Suppose $\mathcal{Q}_j = \cup_{i=1}^{n_j} \mathcal{Q}^{j_i}$. We define a full subcategory $\overline{\mathcal{C}}_0$ of $R({}^{(p-1)}\mathfrak{B}^s)$ consisting of objects M (see Lemma 8.6.1) such that $M_{\mathcal{P}} = k^m, M_{\mathcal{Q}^q} = k^m, 0 \leq q < p$; $M(\lambda)$ has eigenvalue λ^0 with $\sigma^{p-1}(\lambda^0) \neq 0$; $M(a) = \emptyset, \forall a \in \cup_{\eta=0}^{p-1} A^\eta$; $M(c) = \emptyset$ or $0, \forall c \in \cup_{\eta=0}^{p-1} C^\eta$; $M(c^\eta) = (0, \dots, 0, I^m)$ for $0 \leq \eta \leq p-2$. Moreover we define a full subcategory $\overline{\mathcal{C}}$ of $R({}^p\mathfrak{B}^s)$ (see Lemma 8.6.2) according to the following two cases:

- (i) $\mathcal{Q}^{p-1} \in {}^pT^s / {}^p \sim^s$ is an equivalent class;
- (ii) $\mathcal{Q}^{p-1} \subset \mathcal{Q}_{j_0}$ in ${}^pT^s / {}^p \sim^s$ for some $1 \leq j_0 \leq r$.

Then $\forall M \in \overline{\mathcal{C}}, M_{\mathcal{P}} = k^m; M_{\mathcal{Q}_j} = k^{mn_j}$ for $1 \leq j \leq r$ in case (i), or $1 \leq j \leq r, j \neq j_0$ in case (ii); $M_{\mathcal{Q}^{p-1}} = k^n$ in case (i) or $M_{\mathcal{Q}_{j_0}} = k^{n+mn_{j_0}}$ in case (ii). And $M(\lambda)$ has eigenvalue λ^0 with $\sigma^{p-1}(\lambda^0) \neq 0$; $M(a) = \emptyset, \forall a \in \cup_{\eta=0}^{p-1} A^\eta$; $M(c) = \emptyset$ or $0, \forall c \in \cup_{\eta=0}^{p-1} C^\eta$; $M(c^\eta) = (0, \dots, 0, I^m)$ for $0 \leq \eta \leq p-2$; $M(c^{p-1}) = P$.

Let $\mathcal{C}$ be the category defined in Lemma 8.6.2, we define a functor

$$F_1 : \overline{\mathcal{C}} \rightarrow \mathcal{C} \quad \text{with } F_1(M) = M'$$

such that $M'_{\mathcal{P}} = k^m; M'_{\mathcal{Q}_j} = k^{mn_j}$ for $1 \leq j \leq r$ in case (i) or $1 \leq j \leq r, j \neq j_0$ in case (ii); $M'_{\mathcal{Q}^{p-1}} = k^n$ in case (i) or $M'_{\mathcal{Q}_{j_0}} = k^{n+mn_{j_0}}$ in case (ii). And $M'(\lambda) = M(\lambda)$; $M'(a_{j_l}) = (M(c^{j_l}) \mid 0)_{1 \times n_{j_l}}$ for $1 \leq j \leq r$ in case (i) or $1 \leq j \leq r, j \neq j_0$ in case (ii); $M'(c) = P$ in case (i), or $M'(c) = (P \mid 0)_{1 \times (n_{j_0}+1)}$ in case (ii). It is obvious that F_1 is a representation equivalence. In fact, the morphisms in $\overline{\mathcal{C}}$ go to the diagonal parts according to the partition of $({}^pT^s / {}^p \sim^s)$ under the action of F_1 .

Let $\mathcal{D}$ be the category defined in Lemma 8.6.3. Then we have a representation equivalence

$$F_2 : \mathcal{C} \rightarrow \mathcal{D},$$

such that $F_2(M') = M''$ with $M''_{\mathcal{P}} = k^m$ and $M''_{\mathcal{Q}} = k^n$; $M''(\lambda) = M(\lambda)$; $M''(c) = P$. In fact, the morphisms in $\mathcal{C}$ go to the diagonal blocks under the action of F_2 , i.e. “*” of φ_{j_0} in case (ii) go to zero in Lemma 8.6.2.

Thus

$$F = F_2 F_1 : \overline{\mathcal{C}} \rightarrow \mathcal{D}$$

is a representation equivalence. If (e) given in the beginning of the proof was an almost split conflation in $R({}^p\mathfrak{B}^s)$, then so was in $\overline{\mathcal{C}}$. Therefore $F(e)$ would be an almost split conflation in $\mathcal{D}$ by Lemma 8.6.4. Which contradicts to Lemma 8.6.3.

The proof is finished, i.e. ${}^p\mathfrak{B}^s$ is not homogeneous in both cases of Q1 as desired. $\square$

Proposition 9.6.2 The partial boc ${}^p\mathfrak{A}^s$ given by condition Q2 of Proposition 9.5.1 is not homogeneous.

Proof. We fix some $\lambda^0 \in k$ with $h^p(\lambda^0, \lambda^0) \neq 0$ given in Formula (64) of 9.5. Let $\lambda = \lambda^0$, $a_l^p = \emptyset$, $l = 1, \dots, i^p - 1$, $a_{i^p}^p = \nu^p$ then the partial boc ${}^p\mathfrak{B}_{(\lambda^0)}^{s'}$ with $s' = s + (\sum_{\eta=0}^{p-1} J^\eta) + (i^p + n^p)$ induced from ${}^p\mathfrak{B}_{(\lambda^0)}^{s + \sum_{\eta=0}^{p-1} J^\eta}$ is minimal. We construct an infinite list of the objects $\{S'_{\nu_p^0} \mid \nu_p^0 \in k \text{ with } \sigma^p(\nu_p^0) \neq 0\} \subset R({}^p\mathfrak{B}_{(\lambda^0)}^{s'})$, such that $(S'_{\nu_p^0})_{\mathcal{P}} = k$; $(S'_{\nu_p^0})(\nu_p) = (\nu_p^0)$; $(S'_{\nu_p^0})(d_l^p) = \emptyset$ for $1 \leq l \leq n_p$. If

$$\vartheta : R({}^p\mathfrak{B}_{(\lambda^0)}^{s'}) \rightarrow R({}^p\mathfrak{B}^s)$$

is the reduction functor, then $S_{\nu_p^0} = \vartheta(S'_{\nu_p^0})$ provide an infinite list of the indecomposables over ${}^p\mathfrak{B}^s$. If ${}^p\mathfrak{B}^s$ is homogeneous, there exists a cofinite subset $D_0 \subseteq k \setminus \{\text{the roots of } \sigma^p(\nu_p)\}$, such that $S_{\nu_p^0}$ is homogeneous for any $\nu_p^0 \in D_0$. Denote $S'_{\nu_p^0}$ by S' for simplicity. Let $(e') : S' \rightarrow E' \rightarrow S'$ be the almost split conflations starting and ending at S' given by Proposition 6.6.1. Then $(e) : S \rightarrow E \rightarrow S$, the image of (e') under ϑ , is an almost split conflation of $R({}^p\mathfrak{B}^s)$ by Theorem 7.2.1. Construct an object $L \in R({}^p\mathfrak{B}^s)$ such that $L_{\mathcal{P}} = k^2$, $L_{\mathcal{Q}^q} = k^2$; $L(\lambda) = J_2(\lambda^0)$; $L(a) = \emptyset$ for any $a \in \cup_{\eta=0}^{p-1} A^\eta$; $L(c) = \emptyset$ or 0, $\forall c \in \cup_{\eta=0}^{p-1} C^\eta$; $L(c^\eta) = (0 \cdots 0 I_2)$ for $\eta = 1, \dots, p-1$; $L(a_l^p) = \emptyset$ for $l = 1, \dots, (i^p - 1)$, $L(a_{i^p}^p) = \nu_p^0 I_2$; and $L(d_l^p) = \emptyset$ for $l = 1, \dots, n^p$, then L is indecomposable.

$$\begin{array}{ccc}
 & \begin{array}{c} L(\lambda) \quad L(\nu_p) \\ \curvearrowright \quad \curvearrowright \\ k^2 \end{array} & \\
 & \swarrow \varphi' \quad \searrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \\
 \begin{array}{c} E(\lambda) \quad E(\nu_p) \\ \curvearrowright \quad \curvearrowright \\ k^2 \end{array} & \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} & \begin{array}{c} S(\lambda) \quad S(\nu_p) \\ \curvearrowright \quad \curvearrowright \\ k \end{array}
 \end{array}$$

A contradiction follows similarly to the proof of Proposition 7.5.1 for $\mathfrak{B}_4$. Therefore ${}^p\mathfrak{B}^s$ is not homogeneous as desired. $\square$

10 Bipartite bimodule Problems

10.1 Bipartite property

We stress that according to Proposition 7.3.1, 7.3.2, 7.4.1 and 7.5.1, the only possible exceptional situation for a minimally wild bimodule problem being homogeneous is given by MW5 of Theorem 5.6.1. On the other hand Example (4) of 6.5 shows that $Mat(\mathcal{K}, \mathcal{M})$ can be strongly homogeneous in this case. Thus we are forced to focus on some special class of bimodule problems, in which $P_1(\Lambda)$ are included.

Definition 10.1.1 A bimodule problem $(\mathcal{K}, \mathcal{M}, H)$ is said to be *bipartite*, if

- I. $T = T_1 \cup T_2$, where $T_1 = \{1, 2, \dots, t_1\}$, $T_2 = \{t_1 + 1, t_1 + 2, \dots, t_1 + t_2\}$, $T/\sim = T_1/\sim_1 \cup T_2/\sim_2$.
- II. $\mathcal{K} = \left\{ \begin{pmatrix} S_1 & 0 \\ 0 & S_2 \end{pmatrix} \right\}$, where S_r are $t_r \times t_r$ upper triangular matrices, such that $s_{ii}^r = s_{jj}^r$, if $i \sim_r j$, and if $i < j$, s_{ij}^r satisfies the following r -th equation for $r = 1, 2$,

$$\sum_{\mathcal{I}^1 \ni i < j \in \mathcal{J}^1} c_{ij}^{1l_1} x_{ij} = 0,$$

$$0 = \sum_{\mathcal{I}^2 \ni i < j \in \mathcal{J}^2} c_{ij}^{2l_2} y_{ij},$$

where $1 \leq l_r \leq q_{\mathcal{I}\mathcal{J}}^r$ for some $q_{\mathcal{I}\mathcal{J}}^r \in \mathbb{N}$, and for each pair

$$(\mathcal{I}^r, \mathcal{J}^r) \in (T_r/\sim_r) \times (T_r/\sim_r).$$

- III. $\mathcal{M} = \left\{ \begin{pmatrix} 0 & A \\ 0 & 0 \end{pmatrix} \right\}$ where A are $t_1 \times t_2$ matrices satisfying the equations

$$\sum_{(i,j) \in \mathcal{I}^1 \times \mathcal{J}^2} d_{ij}^l z_{ij} = 0,$$

$1 \leq l \leq q_{\mathcal{I}\mathcal{J}}$ for some $q_{\mathcal{I}\mathcal{J}} \in \mathbb{N}$, and for each pair $(\mathcal{I}^1, \mathcal{J}^2) \in (T_1/\sim_1) \times (T_2/\sim_2)$.

- IV. $H = 0$.

- V. The row-indices of the free entries of N_0 (see Formula (19) of 3.6) are pairwise different.

For the sake of convenience we sometimes denote $(\mathcal{K}, \mathcal{M}, H = 0)$ by $(\mathcal{K}_1 \times \mathcal{K}_2, \mathcal{M}, H = 0)$, where $\mathcal{K}_1 \times \mathcal{K}_2 = \{(S_1) \times (S_2)\}$ and $\mathcal{M} = \{(A)\}$. And we denote a triple

$$\left(\begin{pmatrix} 0 & N \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} R_1 & 0 \\ 0 & R_2 \end{pmatrix}, \underline{n} \right)$$

given in Formula (6) of 2.6 by $(N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2)$.

The differential biquiver corresponding to a bipartite bimodule problem is as follows. Let

$$T_1/\sim_1 = \{\mathcal{I}_1, \dots, \mathcal{I}_{s_1}\}, \quad T_2/\sim_2 = \{\mathcal{J}_1, \dots, \mathcal{J}_{s_2}\},$$

then we draw s_1 vertices on the top and s_2 vertices on the bottom. The solid arrows are all from top to bottom, and the dotted arrows given by $\mathcal{K}_1$ are sitting at the top, and those by $\mathcal{K}_2$ are sitting at the bottom, see Examples in 4.3.

Given a finite-dimensional algebra Λ , the bimodule problem $P_1(\Lambda)$, or equivalently $(\mathcal{K}, \mathcal{M}, H = 0)$ with $\mathcal{K} = \tilde{\Lambda} \times \tilde{\Lambda}$, $\mathcal{M} = \text{rad} \tilde{\Lambda}$ is obviously bipartite. Namely, let $T_1 = \{1', 2', \dots, t'\}$ according to the row-indices of $\tilde{\Lambda}$ from top to bottom, and let $T_2 = \{1, 2, \dots, t\}$ according to the column-indices of $\tilde{\Lambda}$ from left to right. Then $\forall \mathcal{J} \in T_2/\sim_2$, with

$$\mathcal{J} = \{j_1 < j_2 < \dots < j_{\mathcal{J}}\},$$

the $j_{\mathcal{J}}$ -th column of N_0 consists of free entries and zeros (see examples of 2.1), conversely any free entries of N_0 must sit at $j_{\mathcal{J}}$ -column for some $\mathcal{J} \in T_2/\sim_2$. In particular, all the row-indices of the free entries are pairwise different.

We claim that for the simplicity of the proof, we stick to have the hypothesis V in the definition, but it may not be essential for our purpose.

10.2 Locations

Assume that our original bimodule problem $(\mathcal{K}, \mathcal{M}, H = 0)$ is wild and bipartite, which has a reduction sequence $(*)$, such that the end term of $(***)$ is in case MW5 of Theorem 5.6.1. Then Lemma 9.3.1 shows the following possibilities of the locations of λ and ν with respect to the block $\overline{G}$ containing ν partitioned by $(T, \sim)$.




figure 1




figure 2




figure 3

Definition 10.2.1 [XZ]. Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bimodule problem having a reduction sequence $(*)$ of parameterized triples, (not necessarily freely). If

$$\overline{N_{p_r q_r}^r} = \begin{pmatrix} 0 & I \\ 0 & 0 \end{pmatrix} \quad \text{or} \quad W$$

given by edge or loop reduction respectively, where $W = \oplus_{j=1}^{\alpha} W_{\lambda^j}$ is a Weyr matrix defined in 2.5 such that

$$W_{l-1,l}^j = \begin{pmatrix} I_{m_l^j} \\ 0 \end{pmatrix}_{m_{l-1}^j \times m_l^j}.$$

Then all the 1's appearing in I and $I_{m_l^j}$, $l = 2, \dots, d_j$, $j = 1, \dots, \alpha$, are called links of $(\mathcal{K}^s, \mathcal{M}^s, H^s)$.

Location 1. There is no link in $\overline{G}$, or all the links in $\overline{G}$ are lower than ν (see figure 1 and 2).

Location 2. There exists at least one link in $\overline{G}$, which is parallel to or higher than ν (see figure 3).

Examples. Consider again the Examples in 9.2. Then Example 1 gives

$$\overline{G} = G = \begin{pmatrix} \emptyset & \emptyset \\ \nu & \emptyset \end{pmatrix}$$

which is in the case of Location 1.

The figure 7 of Example 2 may be used to illustrate the Location 2, but it has only one parameter ν .

10.3 Links

Recall that for any size vector $\underline{n}$ of $(T, \sim)$ its dimension $d = \dim \underline{n} = \sum_{\mathcal{I} \in T/\sim} n_{\mathcal{I}}$ (see 2.2). Denote by $\tau(H_{\underline{n}})$ the number of links in $H_{\underline{n}}$. Then the following proposition can be regarded as a generalization of Theorem A in [XZ].

Proposition 10.3.1 Let

$$(N^r, R^r, \underline{n}^r), \dots, (N^s, R^s, \underline{n}^s)$$

be a reduction sequence of parameterized triples with $r \geq 1$. Then for any $s \geq r$,

$$\dim \underline{n}^r + \tau(H_{\underline{n}^r}^r) = \dim \underline{n}^s + \tau(H_{\underline{n}^s}^s)$$

Proof. We use induction on s . $s = r$ is trivial. Suppose the proposition is true for s , now we consider the $(s+1)$ -th reduction.

Regularization. Clearly, $\dim \underline{n}^s = \dim \underline{n}^{s+1}$ and $\tau(H^s) = \tau(H^{s+1})$ by 3.1.

Edge reduction. Suppose $\overline{N_{p_s q_s}^s} = \begin{pmatrix} 0 & I_d \\ 0 & 0 \end{pmatrix}$, then $\dim \underline{n}^{s+1} = \dim \underline{n}^s - d$ by 3.1, but $\tau(H^{s+1}) = \tau(H^s) + d$, hence the formula holds for $(s+1)$.

Loop reduction. Assume that $\overline{N_{psqs}^s} = W$ is similar to $\bigoplus_{l=1}^{\alpha} \bigoplus_{i=1}^{d_l} J_i(\lambda_l)^{e_l^i}$, then $\tau(W) = \sum_{l=1}^{\alpha} \sum_{i=1}^{d_l} e_l^i(i-1)$. On the other hand, $\dim \underline{n}^{s+1} = \dim \underline{n}^s - \tau(W)$ by 3.1 (see also [XZ, Lemma 5]). On the other hand $\tau(H^{s+1}) = \tau(H^s) + \tau(W)$. Therefore the formula still holds for $(s+1)$. $\square$

We write $\mathcal{N}_r^s = \sum_{l=r}^{s-1} \overline{N_{plql}^l} \otimes \rho^l$, and $\tau(\mathcal{N}_r^s)$ stands for the number of links in $\mathcal{N}_r^s$. It is clear that

$$\tau(\mathcal{N}_r^s) = \tau(H_{\underline{n}^s}^s) - \tau(H_{\underline{n}^r}^r).$$

Corollary 10.3.1 Let $(N^r, R^r, \underline{n}^r), \dots, (N^s, R^s, \underline{n}^s)$ be a reduction sequence of parameterized triples given by one of the three reductions of 3.1. Then $(N^s, R^s, \underline{n}^s)$ is local if and only if $\dim \underline{n}^r = \tau(\mathcal{N}_r^s) + 1$.

Proof. R^s is a local ring, if and only if $\dim \underline{n}^s = 1$, if and only if $\dim \underline{n}^r + \tau(H_{\underline{n}^r}^r) = 1 + \tau(H_{\underline{n}^s}^s)$, if and only if $\dim \underline{n}^r = \tau(\mathcal{N}_r^s) + 1$. $\square$


10.4 The non-homogeneous property in case of MW5

Theorem 10.4.1 Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bipartite bimodule problem. If $(\mathcal{K}, \mathcal{M}, H)$ has a reduction sequence $(*)$ of minimal size, such that the end term $\mathfrak{A}^s$ of the corresponding sequence $(***)$ belongs to MW5 of Theorem 5.6.1, then $\text{Mat}(\mathcal{K}, \mathcal{M})$ is not homogeneous.

Proof. In case of Location 1. We use the structure of adding column given in 9.1 starting from the beginning term $(N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2)$ (see Definition 10.1.1). Since all the indices of the block-columns of N belong to T_2 , and $T_1 \cap T_2 = \emptyset$, the assumptions S02 of Structure 9.1 and S12, $\dots$, S(p-1)2 of 9.5 always take place. We stress that the triple $(N, R_1 \times R_2, \underline{n}_1 \times \underline{n}_2)$ corresponds to the whole boc $\mathfrak{B} = \mathfrak{A}$ (not properly partial!) and $\overline{G} = G$ in this case. We use induction both on the index pairs (α^η, l^η) of ν_η and the index pairs $(\alpha(\zeta), l(\zeta))$ of λ_ζ .

Going back to 9.5, a sequence of layered bocses $\mathfrak{B}^s = \mathfrak{A}^s$, ${}^1\mathfrak{B}^s = {}^1\mathfrak{A}^s$, ${}^2\mathfrak{B}^s = {}^2\mathfrak{A}^s$, $\dots$, constructed under the assumptions Sq1 and Sq2 for $q = 0, 1, \dots$ must stop at some stage, say p since the pairs (α^η, l^η) are bounded by (n_1, t_2) , where n_1 stands for the number of the rows of matrix N , and t_2 the number of the elements of T_2 . If ${}^p\mathfrak{A}^s$ satisfies Q1 or Q2 of Proposition 9.5.1, then ${}^p\mathfrak{A}^s$ is not homogeneous by Proposition 9.6.1 or 9.6.2 respectively. If ${}^p\mathfrak{A}^s$ is in case Q3 of Proposition 9.5.1, and we set $\lambda_1 = \lambda_{p, \gamma^{p-1}}$ with a pair of indices $(\alpha(1), l(1)) \succ (\alpha(0), l(0))$, then we start a new round of the preceding procedure from λ_1 . Inductively, such round must stop at some stage, since the pairs $(\alpha(\zeta), l(\zeta))$ are also bounded by (n_1, t_2) . Our conclusion for Location 1 follows from double induction.

In case of Location 2. $\overline{G}$ is of the following shape, and we will focus on the shadowed rectangle. $\overline{G} \supsetneq G$ in this case.



Suppose that a triple $(N_0^r, R^r, \underline{n}^r)$ (see the notation in the proof of Corollary 8.1.1) is the term of the sequence $(*)$ having $N_{p_r q_r}^r$ as the first block of the shadowed part in Figure (66). Then Proposition 8.1.2 and Condition V of Definition 10.1.1 determines a matrix problem $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \overline{\mathcal{M}}^r, \mathcal{V}^r)$ where

$$\mathcal{K}_1^r = \{(s) | \forall s \in k\}; \quad \mathcal{K}_2^r = \{S_{jj}^r\},$$

here S_{jj}^r is the j -th diagonal block of $\mathcal{K}^r$ partitioned by $(T, \sim)$ with j being the column index of $\overline{G}$; $\mathcal{V} =$ the shadowed part of $\{SH_{\underline{n}^r} - H_{\underline{n}^r}S | \forall S \in \mathcal{K}^r\}$. Furthermore the matrix problem $(\mathcal{K}_1^r \times \mathcal{K}_2^r, \overline{\mathcal{M}}^r, \mathcal{V}^r)$ corresponds to a partial boc $\mathfrak{B}^r$ of $\mathfrak{A}^r$, which must be one sided.

We first consider the induced local boc $\mathfrak{A}_{\mathcal{P}}^r$ at $\mathcal{P}$ obtained by deletion with a layer

$$L_{\mathcal{P}} = (\Gamma'_{\mathcal{P}}; \omega_{\mathcal{P}}; b_1, \dots, b_m, b_{m+1}, \dots, b_n; u)$$

such that $b_1, \dots, b_m$ belong to $\mathfrak{B}^r$ but $b_{m+1}, \dots, b_n$ do not. Then a partial boc $\mathfrak{B}_{\mathcal{P}}^r$ is obtained from $\mathfrak{A}_{\mathcal{P}}^r$ restricted to the shadowed part.

(1) If any reduction sequence starting from the induced boc $\mathfrak{A}_{\mathcal{P}}^r$ meets configuration (1) of Theorem 5.6.1, then the size of the local triple at $\mathcal{P}$ is small than that of $(N^r, R^r, \underline{n}^r)$ since the latter one is not local. We have a contradiction to the minimal size assumption. Therefore $\mathfrak{B}_{\mathcal{P}}^r$ can not satisfy P3 or P4 and must be in case P1 or P2 of 8.2.

(2) If $\mathfrak{B}_{\mathcal{P}}^r$ is in case P2, then $\mathfrak{B}^r$ is given by Figure (46) of 8.2. Suppose that $\delta(\overline{b})$ satisfies Formula (52) of 8.3, then the conflation (e) constructed in Proposition 8.3.1 can also be regarded as an almost split conflation of $R(\mathfrak{A}^r)$. Therefore the same proposition tells that $\mathfrak{A}^r$ is not homogeneous. If $\delta(\overline{b})$ satisfies Formula (51) and $\delta(a)^0$ with respect to $\overline{b}$ are given by Proposition 8.3.2, then $\mathfrak{A}^r$ is not homogenous. In fact the induced partial boc of $\mathfrak{B}^r$ constructed in the same proposition determines a global induced boc of $\mathfrak{A}^r$, which is also in the case of Proposition 7.3.2.

(3) If $\mathfrak{B}_{\mathcal{P}}^r$ is in case P2, and $\delta(\bar{b})$ satisfies Formula (51) of 8.3, moreover $\delta(a)^0$ with respect to $\bar{b}$ satisfy Formula (53) of 8.3. Then going back to Structure 9.1, let N^r be the shadowed part of N_0^r . Let $p \in \mathcal{P} \in T^r / \sim^r$ be the block-row index of N^r , then $T_1^r / \sim_1^r = \{\mathcal{P}\}$ and $T_2^r / \sim_2^r = \{\mathcal{P}, \mathcal{I}_1, \dots, \mathcal{I}_i\}$. N^r is partitioned by $(n_{\mathcal{P}}) \times \underline{n}_2$, $R_1 = (x_{ij})_{n_{\mathcal{P}} \times n_{\mathcal{P}}}$, R_2 is partitioned by $\underline{n}_2 \times \underline{n}_2$. Therefore the assumptions S02 of the structure 9.1 and S12, $\dots$, S(p-1)2 of 9.5 take place by Proposition 8.4.1, since $\bar{B}$, the block determined by $\bar{b}$, will not contain any parameter after any reductions.

(4) If $\mathfrak{B}_{\mathcal{P}}^r$ is in case P1 of 8.2, then the assumption S02 of 9.1 and S12, $\dots$, S(p-1)2 of 9.5 always take place.

Next we perform the structure 9.1 of adding columns in cases (3) and (4) starting from the triple $(N^r, R_1^r \times R_2^r, \underline{n}_1^r \times \underline{n}_2^r)$. Going back to 9.1, 9.2 and 9.3, it is clear that λ, ν both belong to $\mathfrak{B}^s$ and $\Gamma'(\mathcal{P}, \mathcal{P}) = k[\lambda]$. Then we add columns to the partial boc $\mathfrak{B}^s$ and obtain a partial boc ${}^1\mathfrak{B}^s$. ${}^1\mathfrak{B}^s$ determines a global boc ${}^1\mathfrak{A}^r$, which contains ${}^1\mathfrak{B}^s$ as the first part. Since ${}^1\mathfrak{B}^{s+J}$ given in Lemma 9.3.1 is local, ${}^1\mathfrak{A}^{s+J}$, having ${}^1\mathfrak{B}^{s+J}$ as the first part, must be local. In fact let $\mathcal{N}_r^s = \sum_{l=r}^{s-1} \overline{N_{p_l q_l}^l} \otimes \rho^l$ in the sequence (Δ) of 9.1, and ${}^1\mathcal{N}_r^{s+J} = \sum_{l=r}^{s+J-1} \overline{{}^1\mathcal{N}_{p_l q_l}^l} \otimes \rho^l$ in the sequence $({}^1\Delta)$. Then $\dim_k \underline{n}^r = \tau(\mathcal{N}_r^s) + 1$ according to Corollary 10.3.1 since $\mathfrak{A}^s$ is local. On the other hand, $\tau({}^1\mathcal{N}_r^{s+J}) = \tau(\mathcal{N}_r^s) + 1$, and $\dim_k {}^1\underline{n}^r = \dim_k \underline{n}^r + 1$, thus $\dim_k {}^1\underline{n}^r = \tau({}^1\mathcal{N}_r^{s+J}) + 1$. Then ${}^1\mathfrak{A}^{s+J}$ being local follows from Corollary 10.3.1 once again. Therefore we are able to perform the procedure 5.3 for ${}^1\mathfrak{A}^{s+J}$ and obtain parameters $\lambda = \lambda_{10}, \lambda_{11}, \dots, \lambda_{1\gamma^1}$. If $\gamma^1 = 0$, we have the case Q1 of Proposition 9.5.1 for the global boc ${}^1\mathfrak{A}^s$, which is not homogeneous by proposition 9.6.1. If $\gamma^1 > 0$ and $\lambda_{1\gamma^1}$ locates outside of the shadowed part, then we reach the case of Location 1. Now we assume that

(i) $\lambda_{1\gamma^1}$ locates inside of the shadowed part and

(ii) ${}^1\mathfrak{A}^s$ (not only ${}^1\mathfrak{B}^s$!) satisfies S11 of 9.5.

we stress that ${}^1\mathfrak{A}^s$ also satisfies S12 in both cases (3) and (4) as ${}^1\mathfrak{B}^s$ does. Continue to use the structure of adding columns under the assumptions (i) and (ii), we obtain by induction a sequence of partial bocses

$$\mathfrak{B}^s, {}^1\mathfrak{B}^s, {}^2\mathfrak{B}^s, \dots, {}^p\mathfrak{B}^s,$$

(see 9.5), as well as a sequence of global bocses

$$\mathfrak{A}^s, {}^1\mathfrak{A}^s, {}^2\mathfrak{A}^s, \dots, {}^p\mathfrak{A}^s.$$

Since the pairs of indices (α^η, l^η) are bounded by $(n_{\mathcal{P}}^r, t_2^r)$, where t_2^r stands for the number of elements of T_2^r , the procedure must stop at some stage, say p . If $\lambda_{p\gamma^p}$ locates outside of the shadowed part, we reach the case of Location 1. Now suppose $\lambda_{p\gamma^p}$ locates inside the shadowed part. If ${}^p\mathfrak{A}^s$ is in the condition Q1 or Q2 of Proposition 9.5.1, then ${}^p\mathfrak{A}^s$ is not homogeneous by Proposition 9.6.1 or 9.6.2. If ${}^p\mathfrak{A}^s$ is in the condition of Q3 of Proposition 9.5.1, then we set $\lambda_1 = \lambda_{p, \gamma^p-1}$. If λ_1 is outside of the shadowed part, we obtain the case of Location 1. Otherwise, the pair of indices of λ_1 , $(\alpha(1), l(1)) \succ (\alpha(0), l(0))$ in the shadowed part. Thus we start a new round of the above procedure for λ_1 . Inductively such round must stop at some stage, since the pairs $(\alpha(\zeta), l(\zeta))$ are also bounded by $(n_{\mathcal{P}}^r, t_2^r)$. Finally our conclusion for Location 2 follows from double induction. $\square$

Remark. All the discussion in this subsection also applies to MW4 of Theorem 5.6.1, which allows us to treat MW4 and MW5 in a unified way. But because of the particularity of MW5, we prefer to treat them separately. On the other hand we have already presented an easy proof for MW4 in Proposition 7.5.1.

10.5 The main theorem

Key Theorem 10.5.1 Let $(\mathcal{K}, \mathcal{M}, H = 0)$ be a bipartite bimodule problem of wild representation type. Then $\text{Mat}(\mathcal{K}, \mathcal{M})$ is not homogeneous.

Proof. Since $(\mathcal{K}, \mathcal{M}, H = 0)$ is of wild representation type, there must exist a reduction sequence $(*)$ of minimal size, such that the end term $\mathfrak{A}^s$ of the corresponding sequence $(***)$ is in one of the cases MW1—MW5 by Theorem 5.6.1. $\mathfrak{A}^s$ is not homogeneous in case MW1, MW2, MW3, MW4 by Proposition 7.3.1, 7.3.2, 7.4.1, 7.5.1 respectively. Thus $\text{Mat}(\mathcal{K}, \mathcal{M})$ is not homogeneous by Corollary 7.2.1. And $\text{Mat}(\mathcal{K}, \mathcal{M})$ is not homogeneous if $\mathfrak{A}^s$ is in case MW5 by Theorem 10.4.1. $\square$

Corollary 10.5.1 If $P_1(\Lambda)$ is of wild type, then $P_1(\Lambda)$ is not homogeneous.

Theorem 10.5.2 Let Λ be a finite-dimensional k -algebra of representation wild type, then $\Lambda\text{-mod}$ is not homogeneous.

Proof. Since there exists an one-to-one correspondence between the almost split sequences of $\Lambda\text{-mod}$ and the almost split conflations of $P_1(\Lambda)$ except finitely many by Theorem 6.2.1, the non-homogeneous property of $P_1(\Lambda)$ implies the same property of $\Lambda\text{-mod}$. $\square$

Our Main theorem 1.3.2 mentioned in the introduction follows from Theorem 10.5.2 immediately.

Acknowledgement The first author expresses her thanks to R. Bautista. She learnt this problem from him and received his preprint on the problem 14 years ago. She would like to thank C.M. Ringel, D. Simson and S. Liu for their concern for this work. She also thanks Professor Huang Weiming, Hu Yongjian and Zhao Xu'an in Beijing Normal University for their helpful conversations. Some ideas of the manuscript has been discussed with Drozd, Crawley-Boevey and Assashiba during ICRA XI held in Mexico in August 2004. Especially Crawley-Boevey proposed many valuable suggestions at the Asia-link Conference in Beijing in May 2005. In particular, both authors are grateful to Chen Bo, Zhao Deke and Pan Jun, the students of Beijing Normal University, for detailed discussions on this paper at our algebra seminars.

The first author is supported by the Important Project 19331030 of NNSFC, the Research Found for the Doctoral Program of Ministry of Education of China, the Cultivation Fund of the key Scientific and Technical innovation project of Ministry of Education of China, China-UK joint project of the Royal Society (No. 15262), and AsiaLink project of the European Community (ASI/B7-301/98/679-11); the second author is supported by NNSFC 10201004 and 10426014.

References

- [AR] M. Auslander and I. Reiten, Representation theory of artin algebras III, *Comm. alg.* 3(3)(1975), 239-294.
- [ARS] M. Auslander, I. Reiten, Representation theory of artin algebras, *Cambridge studies in advanced mathematics* 36, Cambridge: Cambridge University Press, 1995.
- [B1] R. Bautista, A characterization of finite-dimensional algebra of tame representation type, preprint, UNAM, 1989.
- [B2] R. Bautista, The category of morphisms between projectives. *Comm. Algebra* 32 (2004), no.11, 4303-4331.
- [BBP] R. Bautista, J. Boza and E. Pérez, Reduction functors and Exact structures for Bocses, *Bol.Soc.Mat.Mexicana* (3) 9, 2003, 21-60.
- [BB] W.L. Burt and M.C.R. Butler, Almost split sequences for bocses, *Canadian Math. Soc. Conference proceedings Vol. II*, (1991), 89-121.
- [BCLZ] R. Bautista, W.W. Crawley-Boevey, Tiangang Lei and Yingbo Zhang, On homogeneous exact categories, *J. Alg.* 230, (2000), 665-675.
- [BK] R. Bautista and M. Kleiner, Almost split sequences for relatively projective modules, *J. Alg.* 135(1), (1990), 19-56.
- [CB1] W.W. Crawley-Boevey, On tame algebras and bocses, *Proc. London Math. Soc.*(3), 56(1998), 451-483.
- [CB2] W.W. Crawley-Boevey, Matrix problems and Drozd's theorem, *Topics in algebra*, 26(1990), 199-222.
- [CB3] W.W. Crawley-Boevey. Matrix reductions for Artinian rings, and an application to rings of finite representation type, *J. Alg.* 157(1), 1993.
- [CB4] W.W. Crawley-Boevey, Tame algebras and generic modules, *Proc. London Math. Soc.*(3), 63(1991), 241-165.
- [D] Yu.A.Drozd, On tame and wild matrix problems, *Matrix problems*, Kiev (1977), 39-74.
- [DRSS] P.Dr  xler, I.Reiten, S.Smal   and   .Solberg, Exactcategories and vector space categories, *Transactions of AMS*, Vol. 351, No. 2, Feb. 1999, 647-682.

- [DS] P. Dowbor and A. Skowroński, On the representation type of locally bounded categories, Tsukuba J. Math. Vol. 10 No. 1 (1986), 63-72.
- [GR] P. Gabriel and A.V. Roiter, Representation of finite-dimensional algebras. Encyclopaedia of Math. Sci. Vol 73, Algebra VIII, Springer Verlag, 1992.
- [H] Han Yang, Controlled wild algebras, Proc. London Math. Soc. 83(3), 2001, 279-298.
- [HPR] D. Happel, U. Preiser and C.M. Ringel, Vinberg's characterization of Dynkin diagrams using subadditive function with application to DTr -periodic modules, LNM 832 (1979), 280-294.
- [K] M. Kleiner, Induced modules and comodules and representations of bocses and DGC's. LMN 903, (1980), 168-185.
- [Kr] H. Krause, Generic modules over artin algebras, Proc. London. Math. Soc. (3) 76 (1998), 276-306.
- [N1] H. Nagase, τ -wild algebras, Rep.Alg. Vol.2, Beijing Normal University Press, Beijing, 2002, 365-372.
- [N2] H. Nagase, Non-strictly wild algebras, J.London Math. Soc.(2) 67 (3003), 57-72.
- [O] S.A. Ovsienko, Generic representations of free bocses, preprint 93-010, Univ. Bielefeld.
- [PS] P. Dräxler and Ø. Solberg. Exact factors and exact categories. Boletin Sociedad de Matematica Mexicana 7 (2001), 59-72.
- [Rie] Riedtmann Chr., Algebren, Darstellungsköcher, Ueberlagerungen und zurück, Comm. Math. Helv. 55, 1980, 199-224.
- [R1] C.M. Ringel, The representation type of local algebras, Springer Lecture Notes 488 (1975), 282-305.
- [R2] C.M. Ringel, Tame algebras and integral quadratic forms, Springer LNM 1099 (1984).
- [R3] C.N. Ringel, The development of the representation theory of finite-dimensional algebras 1968-1975, Representation theory and algebraic geometry, 89-115, London Math. Soc. Lecture Note Soc. 238, Cambridge Univ. Press, Cambridge, 1997.
- [Ro] A.V. Rojter, Matrix problems and representations of BOCS's, in "Representation theory I", LNM 831, Springer, Berlin, 1980.

- [S] V.V. Sergeichuk, Canonical matrices for basic matrix problems, Linear algebra and its applications, Vol.317, No.1/3, (2000), 53-102.
- [XZ] Xu Yunge and Zhang Yingbo, Indecomposability and the number of links, Science in China (Series A). Vol.31, No.5, (2001)
- [V] D. Vossieck, A construction of homogeneous matrix problems, Bol. Soc. Mex. III Ser 5, No. 2 (1999), 301-305.
- [Z] Zhang Yingbo, The structure of stable components, Can. J. Math. Vol.43 (3) (1991), 652-672.
- [ZL] Zhang Yingbo and Lei Tiangang, A matrix description of a wild category, Science in China (Series A), Vol.41, No.5, 1998, 461-475.
- [ZLB] Zhang Yingbo, Lei Tiangang and R. Bautista, The representation category of a bocs I-IV, J. BJ. Nor. Univ. (Nat. Sci.), 1995, 31(3), 313-316, 31(4), 440-445, 1996, 32(2), 143-148, 32(3), 289-295.
- [ZX] Zhang Yingbo and Xu Yunge, On tame and wild bocses, Science in China, Ser.A, Mathematics, 2005, Vol.48, No.4, 456-468.
- [ZZ] Zeng Xiangyong and Zhang Yingbo, A correspondence of almost split sequences between some categories, Comm. alg. 29(2), (2001), 1-26.

Zhang Yingbo

Department of Mathematics, Beijing Normal University, Beijing 100875, P.R.China

Email: Zhangyb@bnu.edu.cn

Xu Yunge

Faculty of Mathematics & Computer Science, Hubei University, Wuhan 430062, P.R.China

Email: xuy@hubu.edu.cn